

# Planck Scale Unification in a Supersymmetric Standard Model

R. Howl<sup>1</sup> and S.F. King<sup>2</sup>

*School of Physics and Astronomy, University of Southampton,  
Southampton, SO17 1BJ, U.K.*

## Abstract

We show how gauge coupling unification near the Planck scale  $M_p \sim 10^{19}$  GeV can be achieved in the framework of supersymmetry, facilitating a full unification of all forces with gravity. Below the conventional GUT scale  $M_{GUT} \sim 10^{16}$  GeV physics is described by a Supersymmetric Standard Model whose particle content is that of three complete **27** representations of the gauge group  $E_6$ . Above the conventional GUT scale the gauge group corresponds to a left-right symmetric Supersymmetric Pati-Salam model, which may be regarded as a “surrogate SUSY GUT” with all the nice features of  $SO(10)$  but without proton decay or doublet-triplet splitting problems. At the TeV scale the extra exotic states may be discovered at the LHC, providing an observable footprint of an underlying  $E_6$  gauge group broken at the Planck scale. Assuming an additional low energy  $U(1)_X$  gauge group, identified as a non-trivial combination of diagonal  $E_6$  generators, the  $\mu$  problem of the MSSM can be resolved.

---

<sup>1</sup>E-mail: rhowl@phys.soton.ac.uk.

<sup>2</sup>E-mail: sfk@hep.phys.soton.ac.uk.

# 1 Introduction

Gauge coupling unification and the cancellation of quadratic divergences are two of the most appealing features of supersymmetric (SUSY) extensions of the standard model (SM) [1]. It is well known that the electroweak and strong gauge couplings extracted from LEP data and extrapolated to high energies using the renormalisation group (RG) evolution do not meet within the SM. However, in the framework of the minimal supersymmetric standard model (MSSM) [2] the couplings converge to a common value at some high energy scale. This allows one to embed SUSY extensions of the SM into Grand Unified Theories (GUTs), leading to SUSY GUTs based on  $SU(5)$  or  $SO(10)$ .

However, despite their obvious attractions, SUSY GUTs face some serious challenges from the experimental limits on proton decay on the one hand, and the theoretical requirement of Higgs doublet-triplet splitting on the other as recently discussed for example in [3]. Furthermore the unification of gauge couplings near a conventional GUT scale  $M_{GUT} \sim 10^{16}$  GeV leaves open the question of a full unification of all the forces with gravity, although this may be achieved in the framework of string unification, including high energy threshold effects [4].

It was suggested some time ago that one should consider replacing the SUSY GUT theory by a Pati-Salam gauge group above  $M_{GUT} \sim 10^{16}$  GeV, which plays the role of a “surrogate SUSY GUT” [5], since there is no proton decay or doublet-triplet splitting problem in such a theory. In this scheme the gauge couplings meet at  $M_{GUT} \sim 10^{16}$  GeV, as in the MSSM, and are then held together up to the Planck scale by a combination of left-right symmetry and carefully selected matter content chosen so that the  $SU(4)_{PS}$  gauge group has the same beta function as the  $SU(2)_L \times SU(2)_R$  gauge couplings [5]. However such “theoretical tuning” of the  $SU(4)_{PS}$  and  $SU(2)_L \times SU(2)_R$  beta functions appears to be somewhat contrived.

Recently a so-called Exceptional Supersymmetric Standard Model (ESSM) has been proposed [6, 7], in which the low energy particle content consists of three **27** representations of the gauge group  $E_6$ , plus in addition a pair of non-Higgs doublets  $H', \overline{H}'$  arising from incomplete **27',  $\overline{27}'$**  representations. In the ESSM, gauge coupling unification works even better than in the MSSM [8]. Although the ESSM solves the usual  $\mu$  problem via a singlet coupling to two Higgs doublets, the presence of the non-Higgs doublets  $H', \overline{H}'$  introduces a new  $\mu'$  problem since in this case a singlet coupling generating  $\mu'$  is not readily achieved [6, 7]. However the only purpose of including the non-Higgs states  $H', \overline{H}'$  is to help achieve gauge coupling unification at  $M_{GUT} \sim 10^{16}$  GeV. This allows the possibility of removing the non-Higgs states  $H', \overline{H}'$  from the spectrum. Of course the question of gauge coupling unification must then be addressed, which is the subject of the present paper.

In this paper we consider a similar model to the ESSM but without the additional non-Higgs doublets  $H', \overline{H}'$ . Clearly, without the additional non-Higgs doublets  $H', \overline{H}'$ , the gauge couplings will no longer converge at  $M_{GUT} \sim 10^{16}$  GeV, or any other scale, so at first sight this possibility looks unpromising. However we shall show that, if the theory is embedded into a Pati-Salam theory at  $M_{GUT} \sim 10^{16}$  GeV, then, remarkably,

this leads to a unification of all forces with gravity close to the Planck scale. In the region between  $M_{GUT}$  and  $M_p$  there is not a SUSY GUT but a “surrogate SUSY GUT” based on the Pati-Salam gauge group which resolves the proton decay and doublet-triplet splitting problems of SUSY GUTs, with Planck scale unification achieved in a more natural way than in [5].

Unification in supersymmetric models containing one or three **27** representations of the gauge group  $E_6$  has recently been considered in the literature [9]. Assuming an intermediate Pati-Salam gauge group at the scale  $10^{15}$  GeV at which the Standard Model (SM) couplings satisfy  $\alpha_1 = \alpha_2$ , it was claimed that the resulting Pati-Salam gauge couplings could subsequently meet at a higher scale about  $10^{18}$  GeV [9]. However the condition  $\alpha_1 = \alpha_2$  cannot be consistently applied at the Pati-Salam breaking scale. Instead we find the Pati-Salam breaking scale to be about an order of magnitude larger than the crossing point  $\alpha_1 = \alpha_2$ , close to  $M_{GUT} \sim 10^{16}$  GeV, with full unification close to  $M_p \sim 10^{19}$ . Planck scale unification has also been considered in non-supersymmetric models in [10]. In our analysis we shall naively extrapolate the two-loop RGEs up to  $M_p$ , although in reality we expect new physics effects arising from quantum gravity to set in about an order of magnitude below this. For example, although the Planck scale is usually equated with the Planck mass energy scale  $M_p$  given by  $M_p = \sqrt{\hbar c/G} \approx 1.2 \times 10^{19}$  GeV/ $c^2$  where  $G$  is Newton’s constant, the scale at which quantum gravity becomes relevant may be considered to be  $(8\pi G)^{-1/2} \approx 2.4 \times 10^{18}$  GeV, where the factor of  $8\pi$  comes from the Einstein field equation  $G^{\mu\nu} = 8\pi G T^{\mu\nu}$ , which is sometimes referred to as the reduced Planck scale. It is around this energy scale that an effective quantum field theory of gravity is expected to break down and some new physics takes over since effective quantum field theories of gravity contain corrections to the predictions of General Relativity proportional to powers of  $E^2/M_p^2$  where  $E$  is the energy scale of interest. A more precise estimate of the energy scale at which new physics associated with quantum gravity takes over, based on unitarity violation, may be found in [11], and we return to this point later.

The layout of the rest of this paper is as follows. In the section 2 we consider the pattern of symmetry breaking assumed in this paper. In section 3 we consider the two loop RG evolution of gauge couplings in this model from low energies, through the Pati-Salam breaking scale at  $M_{GUT} \sim 10^{16}$  GeV, assuming various Pati-Salam breaking Higgs sectors, and show that the Pati-Salam gauge couplings converge close to the Planck scale  $M_p \sim 10^{19}$  GeV. In section 4 we shall construct an explicit supersymmetric model of the kind we are considering. Finally we conclude the paper in section 5.

## 2 Pattern of Symmetry Breaking

The two step pattern of gauge group symmetry breaking we analyse in this paper is:

$$E_6 \xrightarrow{M_p} G_{422} \otimes D_{LR} \xrightarrow{M_{GUT}} G_{321} \tag{1}$$

where the gauge groups are defined by:

$$G_{422} \equiv SU(4) \otimes SU(2)_L \otimes SU(2)_R, \quad G_{321} \equiv SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \quad (2)$$

and we have assumed that the first stage of symmetry breaking happens close to the Planck scale and the second stage happens close to the conventional GUT scale. The first stage of symmetry breaking is based on the maximal  $E_6$  subgroup  $SO(10) \otimes U(1)_\psi$  and the maximal  $SO(10)$  subgroup  $G_{422} \otimes D_{LR}$  corresponding to a Pati-Salam symmetry with  $D_{LR}$  being a discrete left-right symmetry.<sup>3</sup>

The pattern of symmetry breaking assumed in this paper is different from that commonly assumed in the literature based on the maximal  $SO(10)$  subgroup  $SU(5) \otimes U(1)_\chi$  [6, 7, 13]. In particular the Pati-Salam subgroup does not contain the Abelian gauge group factor  $U(1)_\chi$ . The only Abelian gauge group factor involved in this pattern of symmetry breaking is  $U(1)_\psi$ , and in the present analysis we assume that this is broken at  $M_p$ . However, as discussed in section 4, there are good phenomenological motivations, related to the solution to the  $\mu$  problem, for preserving a low energy  $U(1)'$  gauge group, and this would require the  $U(1)_\psi$  gauge group to be preserved. In the present paper we do not consider the effect of including the  $U(1)_\psi$  gauge group factor in the RG analysis, however we have checked that Planck scale unification would still be possible, so the results presented here would not be much affected by its inclusion.<sup>4</sup>

The first stage of symmetry breaking close to  $M_p$  will not be considered in this paper. We only remark that the Planck scale theory may or may not be based on a higher dimensional string theory. Whatever the quantum gravity theory is, it will involve some high energy threshold effects, which will depend on the details of the high energy theory, and which we do not consider in our analysis.

The second stage of symmetry breaking close to  $M_{GUT}$  is within the realm of conventional quantum field theory, and requires some sort of Higgs sector, in addition to the assumed matter content of three  $\mathbf{27}$  representations of the gauge group  $E_6$ . In order to break the Pati-Salam symmetry  $G_{422}$  to  $G_{321}$  at  $M_{GUT}$  the minimal Higgs sector required are the  $G_{422}$  representations  $H_R = (4, 1, 2)$  and  $\overline{H}_R = (\overline{4}, 1, \overline{2})$  [14]. When these particles obtain VEVs in the right-handed neutrino directions they break the  $SU(4) \otimes SU(2)_R$  symmetry to  $SU(3)_c \otimes U(1)_Y$  with the desired hypercharge assignments, as discussed later.

Although a Higgs sector consisting of  $H_R$  and  $\overline{H}_R$  is perfectly adequate for breaking Pati-Salam symmetry, it does not satisfy  $D_{LR}$ . We must therefore also consider an extended Higgs sector including their left-right symmetric partners. A minimal left-right symmetric Higgs sector capable of breaking Pati-Salam symmetry consists of the  $SO(10)$  Higgs states  $\mathbf{16}_H$  and  $\overline{\mathbf{16}}_H$ . If complete  $E_6$  multiplets are demanded in the entire theory below  $M_p$ , then the Pati-Salam breaking Higgs sector at  $M_{GUT}$  may be

---

<sup>3</sup>Under  $D_{LR}$  the matter multiplets transform as  $q_L \rightarrow q_L^c$ , and the gauge groups  $SU(2)_L$  and  $SU(2)_R$  become interchanged [12].

<sup>4</sup>With  $U(1)_\psi$  included in the RG analysis for the  $\mathbf{27}_H + \overline{\mathbf{27}}_H$  graph (right panel of figure 1) it may be necessary to increase the effective MSSM threshold to 350 GeV to ensure Planck scale unification for the larger experimental values of the strong coupling constant.

assumed to be  $\mathbf{27_H}$  and  $\overline{\mathbf{27_H}}$ . Therefore in our analysis we shall consider two possible Higgs sectors which contribute to the SUSY beta functions in the region between  $M_{GUT}$  and  $M_p$ , namely either  $\mathbf{16_H} + \overline{\mathbf{16_H}}$  or  $\mathbf{27_H} + \overline{\mathbf{27_H}}$ , where it is understood that only the Pati-Salam gauge group exists in this region, and these Higgs representations must be decomposed under the Pati-Salam gauge group. No such Higgs sectors were included in the analysis in [9].

When  $H_R$  and  $\overline{H}_R$  (contained in either  $\mathbf{16_H} + \overline{\mathbf{16_H}}$  or  $\mathbf{27_H} + \overline{\mathbf{27_H}}$ ) develop VEVs in the right-handed neutrino directions they break the  $SU(4) \otimes SU(2)_R$  symmetry to  $SU(3)_c \otimes U(1)_Y$  with the desired hypercharge assignments. Six of the  $SU(4)$  and two of the  $SU(2)_R$  fields are then given masses related to the VEV of the Higgs bosons and the gauge bosons associated with the  $T^{15}$  and  $T_R^3$  generators are rotated by the Higgs bosons to create one heavy gauge boson and the bino gauge boson associated with  $U(1)_Y$ . In breaking  $SU(4) \otimes SU(2)_R$  to  $SU(3)_c \otimes U(1)_Y$  the SM hypercharge generator is a combination of the diagonal generator  $T^{15} = \sqrt{\frac{3}{2}} \text{diag}(\frac{1}{6}, \frac{1}{6}, \frac{1}{6}, -\frac{1}{2})$  of  $SU(4)$  and the diagonal generator of  $SU(2)_R$ ,  $T_R^3 = \frac{1}{2} \text{diag}(1, -1)$ .  $T^{15} = \sqrt{\frac{3}{2}}(B - L)/2$  where  $B$  and  $L$  are the baryon and lepton number assignments of each standard model particle. Comparing these diagonal generators to the hypercharge values we must have  $Y = T_R^3 + (B - L)/2$ . Then one finds the following relation between the hypercharge gauge coupling constant  $g_Y$  and the  $SU(4)$  and  $SU(2)_R$  gauge coupling constants  $g_4$  and  $g_{2R}$  respectively [15]:

$$\frac{1}{\alpha_Y} = \frac{1}{\alpha_{2R}} + \frac{1}{\frac{3}{2}\alpha_4} \quad (3)$$

where  $\alpha_Y \equiv \frac{g_Y^2}{4\pi}$ ,  $\alpha_{2R} \equiv \frac{g_{2R}^2}{4\pi}$  and  $\alpha_4 \equiv \frac{g_4^2}{4\pi}$ .

Because the Pati-Salam symmetry, and hence the standard model, is assumed to come from an  $E_6$  group, then all the charges and generators should be correctly normalized.<sup>5</sup> In this case the conventional standard model hypercharge assignments must be modified by a factor of  $\sqrt{\frac{5}{3}}$ . Therefore Eq.3 should be rewritten in terms of the ‘GUT’ normalized hypercharge  $g_1 \equiv \sqrt{\frac{5}{3}}g_Y$ :

$$\frac{5}{\alpha_1} = \frac{3}{\alpha_{2R}} + \frac{2}{\alpha_4} \quad (4)$$

where  $\alpha_1 \equiv \frac{g_1^2}{4\pi}$ . Eq.4 is the boundary condition for the gauge couplings at the Pati-Salam symmetry breaking scale, in this case  $M_{GUT}$ . Due to left-right symmetry, at the Pati-Salam symmetry breaking scale we have the additional boundary condition  $\alpha_{2L} = \alpha_{2R}$ . In [9] it was assumed that at the Pati-Salam symmetry breaking scale  $\alpha_1 = \alpha_{2L} = \alpha_{2R}$  which disagrees with Eq.4, since  $\alpha_4 \neq \alpha_{2L} = \alpha_{2R}$  at this scale, as discussed in the next section.

<sup>5</sup>We choose to normalize the  $E_6$  generators  $G^a$  by  $\text{Tr}(G^a G^b) = 3\delta^{ab}$ . It then follows that the Pati-Salam and standard model operators are conventionally normalized by  $\text{Tr}(T^a T^b) = \frac{1}{2}\delta^{ab}$ .




Figure 1: Two-loop Planck Scale Unification in a supersymmetric model which contains three generations of SUSY  $\mathbf{27}$  particles, above an assumed threshold scale of 1.5 TeV. Below this scale the MSSM is assumed with a threshold scale of 250 GeV, below which the SM is assumed. At the scale  $M_{GUT} = 10^{16.44(2)}$  GeV the spectrum is embedded into a left-right symmetric Pati-Salam theory, with  $\alpha_4 = \alpha_3$  at  $M_{GUT} = 10^{16.44(2)}$  GeV, and  $\alpha_{2L} = \alpha_{2R}$  above this scale. The left panel contains the additional the SUSY Higgs contained in  $\mathbf{16}_H + \overline{\mathbf{16}}_H$  while the right panel contains the SUSY Higgs in  $\mathbf{27}_H + \overline{\mathbf{27}}_H$  which enter above  $M_{GUT} = 10^{16.44(2)}$  GeV. The Pati-Salam gauge couplings  $\alpha_4$  and  $\alpha_{2L} = \alpha_{2R}$  converge at  $10^{18.83(7)}$  GeV and  $10^{18.97(9)}$  GeV for the left and right panels respectively, close to the Planck scale, leading to a unified coupling of  $\alpha_P = 0.166(7)$  or  $\alpha_P = 0.321(46)$ . The numbers in parentheses represent the error resulting from the experimental error in the coupling constants.

### 3 Two-loop RG Analysis: Planck Scale Unification

In this section we perform a SUSY two-loop RG analysis of the gauge couplings, corresponding to the pattern of symmetry breaking discussed in the previous section. According to our assumptions there are three complete  $\mathbf{27}$  SUSY representations of the gauge group  $E_6$  in the spectrum which survive down to low energies, but, unlike the original ESSM, there are no additional  $H', \overline{H}'$  states so the gauge couplings are not expected to converge at  $M_{GUT}$ . Instead, we envisage the pattern of symmetry breaking shown in Eq.1, where above the Pati-Salam symmetry breaking scale  $M_{GUT}$  we assume, in addition to the three  $\mathbf{27}$  representations, a Pati-Salam symmetry breaking Higgs sector of either  $\mathbf{16}_H + \overline{\mathbf{16}}_H$  or  $\mathbf{27}_H + \overline{\mathbf{27}}_H$  which are assumed to gain masses of order the Pati-Salam symmetry breaking scale  $M_{GUT}$ , leaving only the three  $\mathbf{27}$  matter representations below this scale.

For the present RG analysis, we run the couplings up from low energies to high energies, using as input the SM couplings measured on the Z-pole at LEP, which are as follows [16]:  $\alpha_1(M_Z) = 0.016947(6)$ ,  $\alpha_2(M_Z) = 0.033813(27)$  and  $\alpha_3(M_Z) = 0.1187(20)$ . The general two-loop beta functions used to run the gauge couplings can be found in [17]. From  $M_Z$  up to an assumed MSSM threshold energy of 250 GeV we consider only the non-SUSY SM spectrum including a top quark threshold at 172 GeV. From 250 GeV to 1.5 TeV we include all the states of the MSSM. From 1.5 TeV up to the Pati-Salam symmetry breaking scale we include all the remaining states which fill out three complete SUSY  $\mathbf{27}$  representations. The assumed threshold energies correspond

to those in [8], where a full discussion of MSSM and ESSM threshold effects is given. The only difference is that here we do not include the  $H', \overline{H}'$  states of the ESSM, so the gauge couplings do not converge at  $M_{GUT}$ . Instead  $M_{GUT}$  is taken to be the Pati-Salam symmetry breaking scale, which is determined as follows.

In the previous section we discussed the relation in Eq.4 between the hypercharge and Pati-Salam coupling constants at the Pati-Salam symmetry breaking scale. This can be turned into a boundary condition involving purely  $G_{321}$  couplings constants at the Pati-Salam breaking scale, since  $SU(3)_c$  comes from  $SU(4)$  so  $\alpha_3 = \alpha_4$  at this scale, and, as remarked,  $D_{LR}$  symmetry requires that  $\alpha_{2R} = \alpha_{2L}$  at the Pati-Salam symmetry breaking scale. Therefore Eq.(4) can be re-expressed as:

$$\frac{5}{\alpha_1} = \frac{3}{\alpha_{2L}} + \frac{2}{\alpha_3}. \quad (5)$$

Having specified the low energy matter content, and thresholds, Eq.5 allows a unique determination of the Pati-Salam breaking scale, by simply running up the gauge couplings until the condition is satisfied. In practice,  $\alpha_3$  runs quite slowly (its one loop beta-function is zero), while the inverse hypercharge coupling decreases most rapidly and the condition is satisfied for a Pati-Salam symmetry breaking scale about an order of magnitude higher energy scale than the crossing point of  $\alpha_1$  and  $\alpha_2$  assumed in [9]. Assuming the above matter content and threshold corrections, the Pati-Salam symmetry is found to be broken at  $M_{GUT} = 10^{16.44(4)}$  GeV as illustrated in Figure 1. This is close to the conventional GUT energy scale, and justifies our use of the notation  $M_{GUT}$  to denote the Pati-Salam breaking scale.

Above the scale  $M_{GUT}$  we run up the two Pati-Salam gauge couplings, namely  $\alpha_4$  and  $\alpha_{2L} = \alpha_{2R}$ , including, in addition to the three SUSY **27** matter representations, also a Pati-Salam SUSY Higgs breaking sector consisting of either  $\mathbf{16}_H + \overline{\mathbf{16}}_H$  or  $\mathbf{27}_H + \overline{\mathbf{27}}_H$ . The Pati-Salam couplings are found to converge at either  $10^{18.83(7)}$  GeV or  $10^{18.97(9)}$  GeV, respectively, as shown in Figure 1.<sup>6</sup> These values are close to the Planck scale  $M_p = 1.2 \times 10^{19}$  GeV, and suggests a Planck scale unification of all forces with gravity.

The value of the gauge coupling constant at the unification scales  $10^{18.83(7)}$  GeV or  $10^{18.97(9)}$  GeV is  $\alpha_P = 0.166(7)$  or  $\alpha_P = 0.321(46)$  for the  $\mathbf{16}_H + \overline{\mathbf{16}}_H$  or  $\mathbf{27}_H + \overline{\mathbf{27}}_H$  particle spectra, respectively. These values of the unified gauge coupling at the Planck scale are much larger than the conventional values of  $\alpha_{GUT}$ , and indeed are larger even than  $\alpha_3(M_Z)$ , however they are still in the perturbative regime. Of course there are expected to be large threshold corrections coming from Planck scale physics which are not included in our analysis. Indeed, we would expect that QFT breaks down as we approach the Planck scale, so that our RG analysis ceases to be valid as we approach the Planck scale, as remarked in the Introduction. The precise energy scale  $E_{new}$  at which quantum field theories of gravity are expected to break down and new physics

---

<sup>6</sup>If we were to drop the  $D_{LR}$  symmetry then, with a minimal Pati-Salam Higgs content consisting of just  $H_R$  and  $\overline{H}_R$ , the equation  $\frac{5}{\alpha_1} = \frac{3}{\alpha_{2R}} + \frac{2}{\alpha_3}$  at the Pati-Salam scale would predict that the Pati-Salam symmetry is broken at  $10^{14.4(1)}$  GeV and that unification would occur at  $10^{19.72(15)}$  GeV.

$Z_2$	$R$ -charges	Incomplete $E_6$ multiplets
+	$\frac{2}{3}$	$\phi_0 = S_3, h_3$
-	$\frac{2}{3}$	$\phi_i = F_i, F_i^c, \mathcal{D}_i, h_\alpha, S_\alpha$
+	$\frac{1}{3}$	$\mathbf{16}_H$ or $\mathbf{27}_H$
+	$\frac{1}{3}$	$\overline{\mathbf{16}}_H$ or $\overline{\mathbf{27}}_H$
-	0	$\Sigma$
+	$\frac{4}{3}$	$M$

Table 1: The  $Z_2$  and  $R$ -charge assignments of the Pati-Salam respecting incomplete  $E_6$  multiplets of the MESSM. The four  $\mathbf{27}$  multiplets are divided into  $\phi_0$  and  $\phi_i$  where  $i = 1 \dots 3$ .  $\phi_0$  contains the MSSM Higgs doublets  $h_3$  and a Pati-Salam symmetry singlet  $S_3$  which gives mass to the  $\mathcal{D}_i$  and  $h_\alpha$  particles in  $\phi_i$  where  $\alpha = 1, 2$ .  $\phi_i$  contains the quarks and leptons  $F_i$  and  $F_i^c$  as well as the exotic quarks  $\mathcal{D}_i$  and non-Higgs  $h_\alpha$ . The  $\mathbf{16}_H$  and  $\overline{\mathbf{16}}_H$  or  $\mathbf{27}_H$  and  $\overline{\mathbf{27}}_H$  multiplets of  $SO(10)$  and  $E_6$  respectively break the Pati-Salam symmetry and are given a GUT scale mass from the  $E_6$  singlet  $M$ .  $\Sigma$  is another  $E_6$  singlet that gets a VEV within the energy range  $10^{7-11}$  GeV to sufficiently suppress proton decay and to allow the exotic particles to decay with a rate that avoids cosmological problems.

takes over is discussed in [11] based on estimates of the scale of violation of (tree-level) unitarity. An upper bound for this new physics energy scale is given by  $E_{new}^2 = 20[G(\frac{2}{3}N_s + N_f + 4N_V)]^{-1}$  where  $N_s$ ,  $N_f$  and  $N_V$  are the number of scalars, fermions and vectors respectively that gravity couples to. Assuming three low-energy  $\mathbf{27}$  multiplets,  $E_{new}$  would be equal to  $10^{18.6}$  GeV which sets an upper bound for the scale at which our quantum field theory analysis (and with any corrections from effective quantum gravity theories included) can no longer be trusted. We have shown that the gauge coupling constants are predicted to be very close to one another at this scale and that, if extrapolated, unify just below  $M_p$ . We have naively extrapolated the RGEs up to  $M_p$ , even though new physics associated with quantum gravity must enter an order of magnitude below this. The fact that the two PS couplings are very close to each other at  $E_{new}$ , and are on a convergent trajectory must be regarded, at best, as a suggestive hint of a unification of the gauge fields with gravity in this approach.

## 4 Constructing a Realistic Model

We have shown that a low energy matter and Higgs content corresponding to three  $\mathbf{27}$  multiplets of the  $E_6$ , embedded in a left-right Pati-Salam symmetry at the GUT scale, can lead to Planck scale unification. However, assuming superpotential interactions corresponding to the  $E_6$  respecting operator  $\mathbf{27}_i \mathbf{27}_j \mathbf{27}_k$  where  $i, j$  and  $k$  are family indices, leads to trouble with phenomenology due to proton decay mediated by TeV scale colour triplets, and flavour changing neutral currents (FCNCs) due to multiple Higgs-like doublets. The solution proposed in [7, 8] is to consider incomplete  $E_6$  supermultiplets at low energies, which allows extra symmetries to appear at low energies that forbid proton decay and suppress FCNCs, while allowing the exotic colour triplets to decay with a lifetime less than about one second, to avoid conflict with nucleosynthesis. The same symmetries cannot be used here since they would not respect the Pati-Salam symmetry,



so we must therefore seek alternative symmetries that are consistent with the Pati-Salam symmetry.

We now propose a realistic supersymmetric model which contains the matter content of three  $\mathbf{27}$ 's of  $E_6$  at low energy, arising from four incomplete  $\mathbf{27}$ 's of  $E_6$  at high energy, and is embeddable in Pati-Salam (PS), without leading to conflict with phenomenology or cosmology. We shall call such a model the minimal exceptional supersymmetric standard model (MESSM) to distinguish it from the ESSM. In the MESSM we shall impose a  $Z_2 \otimes U(1)_R$  symmetry, under which incomplete multiplets of four  $E_6$   $\mathbf{27}$  states  $\phi_0$  and  $\phi_i$ , where  $i = 1 \dots 3$ , transform as shown in Table 1. As in the ESSM, we assume that two Higgs doublets which transform under the PS symmetry as  $h_3 = (1, 2, 2)$  and Higgs singlet  $S_3 = (1, 1, 1)$  arise from the  $\phi_0$  representation, which replace the Higgs in the  $\phi_3$ , leaving the corresponding states  $h_\alpha = (1, 2, 2)$  and  $S_\alpha = (1, 1, 1)$  in the  $\phi_{1,2}$  representations which do not develop VEVs or couple to quarks or leptons, and are “non-Higgs”. The remaining matter content of the  $\phi_i$ , consist of usual quarks and leptons which transform under PS as  $F_i = (4, 1, 2)$  and  $F_i^c = (\bar{4}, 1, 2)$ , the Pati-Salam singlets  $S_\alpha = (1, 1, 1)$  and coloured vector-like quarks  $\mathcal{D}_i = D_i + \bar{D}_i = (6, 1, 1)$ . Interactions between the non-Higgs and matter particles and exotic quarks and matter particles from the  $\phi_i \phi_j \phi_k$  operator would introduce flavour changing neutral currents and rapid proton decay respectively. We suppress these phenomenologically problematic terms by the  $Z_2$  symmetry. However, forbidding all the interactions between the exotic quarks and matter particles could cause serious cosmological problems since this would ensure that the lightest exotic quark is a stable particle whereas cosmology requires the lightest exotic to decay with a lifetime less than about 0.1s to avoid any problems with nucleosynthesis [18]. Therefore the  $Z_2$  symmetry must be broken by the VEV of a new singlet  $\Sigma$  which is sufficiently large to enable the exotic quarks to decay rapidly, but sufficiently small to be consistent with proton decay. This turns out to be possible as we discuss shortly. As well as the three incomplete  $\mathbf{27}$  multiplets of  $E_6$ , there are additional  $\mathbf{16}_H + \bar{\mathbf{16}}_H$  or  $\mathbf{27}_H + \bar{\mathbf{27}}_H$  Higgs multiplets that break the Pati-Salam symmetry and are assumed to reside at the GUT scale. To give a GUT scale mass to these Higgs multiplets we introduce another  $E_6$  singlet  $M$ , which gets a VEV at the GUT scale.

Using Table 1 the only superpotential terms that are allowed are the following:  $\phi_0 \phi_0 \phi_0$  which generates the term  $S_3 h_3 h_3$  which gives an effective  $\mu$ -term when  $S_3$  gets a VEV;  $\phi_0 \phi_i \phi_j$  which generates the rest of the MSSM superpotential terms and gives mass to the “non-MSSM” particles  $\mathcal{D}_i$ ,  $h_\alpha$  and  $S_\alpha$ ;  $\frac{1}{M_p} \Sigma \phi_i \phi_j \phi_k$  which allows the exotic quarks to decay once  $\Sigma$  gets a VEV at an energy scale as discussed below;  $\frac{1}{M_p} \Sigma \phi_i \phi_0 \phi_0$  which gives a mass mixing between Higgs and non-Higgs;  $\frac{1}{M_p} \phi_i \phi_j \bar{\mathbf{16}}_H \bar{\mathbf{16}}_H$  (or  $\frac{1}{M_p} \phi_i \phi_j \mathbf{27}_H \bar{\mathbf{27}}_H$ ) which generates right-handed neutrino masses;<sup>7</sup>  $\frac{1}{M_p} \phi_0 \phi_0 \bar{\mathbf{16}}_H \bar{\mathbf{16}}_H$  (or  $\frac{1}{M_p} \phi_0 \phi_0 \mathbf{27}_H \bar{\mathbf{27}}_H$ ) which are harmless; and  $M \mathbf{16}_H \bar{\mathbf{16}}_H$  (or  $M \mathbf{27}_H \bar{\mathbf{27}}_H$ ) which gives a GUT scale mass to the Higgs multiplets  $\mathbf{16}_H + \bar{\mathbf{16}}_H$  or  $\mathbf{27}_H + \bar{\mathbf{27}}_H$ .

<sup>7</sup>The superpotential term  $\frac{1}{M_p} \phi_i \phi_j \bar{\mathbf{16}}_H \bar{\mathbf{16}}_H$  is meant to represent all the Pati-Salam operators that are found once the  $SO(10)$  multiplet  $\bar{\mathbf{16}}_H$  is decomposed into its Pati-Salam representations. Similar meanings apply to the superpotential terms  $\frac{1}{M_p} \phi_i \phi_j \mathbf{27}_H \bar{\mathbf{27}}_H$ ,  $\frac{1}{M_p} \phi_0 \phi_0 \bar{\mathbf{16}}_H \bar{\mathbf{16}}_H$ ,  $\frac{1}{M_p} \phi_0 \phi_0 \mathbf{27}_H \bar{\mathbf{27}}_H$ ,  $M \mathbf{16}_H \bar{\mathbf{16}}_H$  and  $M \mathbf{27}_H \bar{\mathbf{27}}_H$ .

The  $D_i + \bar{D}_i$  components of the superpotential term  $\phi_i\phi_j\phi_k$  cause proton decay through the decay channels  $p \rightarrow K^+\bar{\nu}$  via  $d = 5$  operators (via the  $\phi_0\phi_j\phi_k$  term which is responsible for the triplet mass  $m_D$ ) and  $p \rightarrow \pi^0 e^+$  via  $d = 6$  operators with matrix elements proportional to  $1/m_D$  and  $1/m_D^2$  respectively. If the exotic quarks get a mass of order  $m_D = 1.5$  TeV from  $\phi_0\phi_j\phi_k$  then we estimate that the term  $\phi_i\phi_j\phi_k$  must be multiplied by an effective Yukawa coupling smaller than about  $10^{-8}$  for the proton's lifetime to be above  $1.6 \times 10^{33}$  years and  $5.0 \times 10^{33}$  years which are the present experimental limits for the  $p \rightarrow K^+\bar{\nu}$  and  $p \rightarrow \pi^0 e^+$  decay modes respectively [19]. In the MESSM the  $\phi_i\phi_j\phi_k$  terms are forbidden by a  $Z_2$  symmetry but are effectively generated from the non-renormalizable terms  $\frac{1}{M_p}\Sigma\phi_i\phi_j\phi_k$  when  $\Sigma$  gets a VEV which must therefore be less than  $10^{11}$  GeV to avoid experimentally observable proton decay. The effective operators  $\phi_i\phi_j\phi_k$  from  $\frac{1}{M_p}\Sigma\phi_i\phi_j\phi_k$  are also the source of exotic quark decay in this model and, for the exotic quarks to have a lifetime less than 0.1s, we estimate that the  $\phi_i\phi_j\phi_k$  operators must be multiplied by an effective Yukawa coupling no less than  $10^{-12}$ , in which case  $\Sigma$  must get a VEV greater than  $10^7$  GeV. Therefore, for the model to be phenomenologically acceptable, we require that the  $E_6$  singlet  $\Sigma$  should get a VEV between  $10^{7-11}$  GeV.

As well as the  $Z_2$  and  $U(1)_R$  symmetries, a  $U(1)_\psi$  symmetry from  $E_6$  may also be assumed. In our RG analysis we assumed for simplicity that  $U(1)_\psi$  is broken at  $M_p$ . However  $U(1)_\psi$  may be unbroken at  $M_p$ , and is broken instead at  $M_{GUT}$  by the Pati-Salam symmetry breaking Higgs  $H_R$  and  $\bar{H}_R$ . These Higgs also break the diagonal generator  $T_R^3$  of  $SU(2)_R$  and the  $B - L$  generator of  $SU(4)$  down to hypercharge  $Y$ , with  $Y = T_R^3 + (B - L)/2$  taking a zero value for the right-handed neutrino and anti-neutrino components. However this is not the only Abelian generator that is preserved by this Higgs sector. Under  $E_6 \rightarrow SO(10) \otimes U(1)_\psi$ ,  $\mathbf{27} \rightarrow \mathbf{16}_{1/2} + \mathbf{10}_{-1} + \mathbf{1}_2$ , and the  $U(1)_\psi$  generator may be written as  $T_\psi = \text{diag}(1/2, -1, 2)$ .<sup>8</sup> The right-handed neutrino component of the Higgs sector which develops the VEV will therefore also preserve the generators  $T_\psi - (B - L)/2$  and  $T_\psi + T_R^3$ , in addition to  $Y = T_R^3 + (B - L)/2$ . It is straightforward to show that precisely one additional Abelian generator orthogonal to  $U(1)_Y$  is preserved, namely:

$$X = (T_\psi + T_R^3) - c_{12}^2 Y \quad (6)$$

where  $c_{12} = \cos \theta_{12}$  and the mixing angle is given by

$$\tan \theta_{12} = \frac{g_{2R}}{g_{B-L}}, \quad g_{B-L} = \sqrt{\frac{3}{2}} g_4, \quad (7)$$

where the Pati-Salam coupling constants  $g_{2R}$  and  $g_4$  are evaluated at  $M_{GUT}$ .

The  $U(1)_X$  associated with the preserved generator in Eq.6 is an anomaly-free gauge group which plays the same role in solving the  $\mu$  problem as the  $U(1)_N$  of the ESSM, since it allows the coupling  $Sh_u h_d$  which generates an effective  $\mu$  term, while forbidding  $S_3$  and the  $\mu h_u h_d$ .  $U(1)_X$  is broken by the  $S$  singlet VEV near the TeV scale, yielding

<sup>8</sup>The respective  $E_6$  normalized generator is  $G^{78} = \frac{1}{\sqrt{6}} \text{diag}(\frac{1}{2}, -1, 2)$ .

a physical  $Z'$  which may be observed at the LHC. We emphasize that this  $Z'$  is distinct from those usually considered in the literature based on linear combinations of the  $E_6$  subgroups  $U(1)_\psi$  and  $U(1)_\chi$  since, in the MESSM,  $U(1)_\chi$  is necessarily broken at  $M_p$ . In particular the  $Z'$  of the MESSM based on  $U(1)_X$  and that of the ESSM based on  $U(1)_N$  will have different physical properties.

## 5 Summary

In this paper we have proposed and discussed a supersymmetric standard model, valid below the conventional GUT scale  $M_{GUT} \sim 10^{16}$  GeV. The particle content consists of three complete SUSY **27** representations of  $E_6$ . However the  $E_6$  gauge group is broken at  $M_p$  and the low energy gauge group is just that of the SM, supplemented by an additional  $U(1)_X$  gauge group. The Higgs doublets which break electroweak symmetry arise from a **27** representation of  $E_6$  as in the ESSM. The model is an example of a low energy Supersymmetric Standard Model (SSM) whose spectrum contains only complete GUT representations. In this respect it is quite unlike any of the conventional SSM's in the literature such as the MSSM, NMSSM or ESSM which all contain low energy Higgs (or non-Higgs) states which do not form complete GUT representations.

We have shown that while gauge coupling at  $M_{GUT} \sim 10^{16}$  GeV is lost, the new model suggests a gauge coupling unification near the Planck scale  $M_p \sim 10^{19}$  GeV. Although we have naively extrapolated the RGEs up to  $M_p$ , we have discussed the fact that in reality there will be new physics effects associated with quantum gravity that will enter about an order of magnitude below this scale, so the idea of Planck scale unification must be considered with caution. Therefore our results can only be regarded as suggestive of Planck scale unification. All we can say is that a supersymmetric model with the low energy matter content of three **27**'s of  $E_6$ , embedded in a Pati-Salam gauge group above  $M_{GUT}$ , gives rise to a high energy theory with two gauge couplings which are very close to each other and converging at the scale at which quantum gravity effects are expected to set in.

As remarked, above  $M_{GUT} \sim 10^{16}$  GeV the model is embedded into a left-right symmetric Supersymmetric Pati-Salam model, which may be regarded as a ‘‘surrogate SUSY GUT’’ with all the nice features of  $SO(10)$  but without proton decay or doublet-triplet splitting problems. The Pati-Salam gauge group is broken at  $M_{GUT} \sim 10^{16}$  GeV by a Higgs sector contained in either  $\mathbf{16}_H + \overline{\mathbf{16}}_H$  or  $\mathbf{27}_H + \overline{\mathbf{27}}_H$ , leaving only the desired spectrum below this scale. The three right-handed neutrinos and sneutrinos gain masses set by the scale  $M_{GUT}^2/M_p$ . At the TeV scale the extra exotic states of the model, which fill out three complete SUSY **27** representations (minus the three right-handed neutrinos and sneutrinos) may be discovered at the LHC, providing an observable footprint of an underlying  $E_6$  gauge group broken at the Planck scale. We have shown that it is possible to construct a realistic supersymmetric model of this kind, which is consistent with phenomenology and cosmology, by using incomplete  $E_6$  multiplets and assuming a symmetry  $Z_2 \otimes U(1)_R$ , and we called the resulting model the MESSM to distinguish it from the ESSM.

In our unification analysis performed in this paper we assumed for simplicity that the  $U(1)_\psi$  gauge group is broken at  $M_p$ . However, in order to solve the  $\mu$  problem, we have shown that it is necessary that the  $U(1)_\psi$  gauge group survives down to  $M_{GUT} \sim 10^{16}$  GeV, so that below this scale a low energy  $U(1)_X$  gauge group emerges as a linear combination of  $U(1)_\psi$  and diagonal Pati-Salam generators. Although we have not considered the effect of the  $U(1)_\psi$  and  $U(1)_X$  gauge groups in the RG analysis, we have checked that Planck scale unification is still possible if they are included. The phenomenology of a low energy  $Z'$ , corresponding to a  $U(1)_X$  which is not a simple linear combination of the  $E_6$  gauge group  $U(1)_\psi$  and  $U(1)_\chi$ , has not so far been considered in the literature and requires a dedicated study [20].

## Acknowledgments

We would like to thank Peter Zerwas for encouragement. RJH acknowledges a PPARC Studentship. SFK acknowledges partial support from the following grants: PPARC Rolling Grant PPA/G/S/2003/00096; EU Network MRTN-CT-2004-503369; EU ILIAS RII3-CT-2004-506222; NATO grant PST.CLG.980066.

## References

- [1] For a recent review see e.g. D. J. H. Chung, L. L. Everett, G. L. Kane, S. F. King, J. D. Lykken and L. T. Wang, Phys. Rept. **407** (2005) 1 [arXiv:hep-ph/0312378].
- [2] J. Ellis, S. Kelley, D.V. Nanopoulos, Phys. Lett. B **260** (1991) 131; P. Langacker, M. Luo, Phys. Rev. D **44** (1991) 817; U. Amaldi, W. de Boer, H. Furstenuau, Phys. Lett. B **260** (1991) 447; F. Anselmo, L. Cifarelli, A. Peterman, A. Zichichi, Nuovo Cimento **104A** (1991) 1817, **105A** (1992) 581.
- [3] S. Raby, Rept. Prog. Phys. **67** (2004) 755 [arXiv:hep-ph/0401155].
- [4] For a recent discussion see e.g. G. G. Ross, arXiv:hep-ph/0411057.
- [5] S. F. King, Phys. Lett. B **325** (1994) 129 [Erratum-ibid. B **325** (1994) 538].
- [6] S. F. King, S. Moretti and R. Nevzorov, Phys. Lett. B **634** (2006) 278 [arXiv:hep-ph/0511256].
- [7] S. F. King, S. Moretti and R. Nevzorov, Phys. Rev. D **73** (2006) 035009 [arXiv:hep-ph/0510419].
- [8] S. F. King, S. Moretti and R. Nevzorov, arXiv:hep-ph/0701064.
- [9] W. Kilian and J. Reuter, Phys. Lett. B **642** (2006) 81 [arXiv:hep-ph/0606277].
- [10] For other discussions of Planck scale unification see e.g. J. D. Lykken and S. Wilenbrock, Phys. Rev. D **49** (1994) 4902 [arXiv:hep-ph/9309258].

- [11] T. Han and S. Willenbrock, Phys. Lett. B **616** (2005) 215 [arXiv:hep-ph/0404182].
- [12] D. Chang, R. N. Mohapatra and M. K. Parida, Phys. Rev. D **30** (1984) 1052.
- [13] J. L. Hewett and T. G. Rizzo, Phys. Rept. **183** (1989) 193; M. Cvetič and P. Langacker, Phys. Rev. D **54** (1996) 3570 [arXiv:hep-ph/9511378];
- [14] S. F. King and Q. Shafi, Phys. Lett. B **422** (1998) 135 [arXiv:hep-ph/9711288].
- [15] J. C. Pati and A. Salam, Phys. Rev. D **10** (1974) 275.
- [16] W. M. Yao *et al.* [Particle Data Group], J. Phys. G **33** (2006) 1.
- [17] M. E. Machacek, M. T. Vaughn Nucl. Phys. B **222** (1983) 83;
- [18] M. Kawasaki, K. Kohri and T. Moroi, Phys. Rev. D **71** (2005) 083502 [arXiv:astro-ph/0408426].
- [19] S. Raby in [16].
- [20] R. Howl and S.F. King, in preparation.