

Non-Abelian Discrete Flavor Symmetries

Ernest Ma

*Department of Physics and Astronomy, University of California,
Riverside, California 92521, USA*

Abstract

This is an incomplete survey of some non-Abelian discrete symmetries which have been used recently in attempts to understand the flavor structure of leptons and quarks. To support such symmetries, new scalar particles are required. In some models, they are very massive, in which case there may not be much of a trace of their existence at the TeV scale. In other models, they are themselves at the TeV scale, in which case there is a reasonable chance for them to be revealed at the LHC (Large Hadron Collider) at CERN.

Introduction

Leptons and quarks come in three families. Their flavor structure, i.e. the specific values of their mass and mixing matrices, has long been a puzzle and a subject of study. In recent years, with the steady accumulation of data on solar and atmospheric neutrino oscillations, the lepton mixing matrix has been determined to a large extent and it came as a surprise to many that it does not resemble at all the known quark mixing matrix. Is there a way to understand this? One possible approach is the use of non-Abelian discrete symmetries, such as S_3 and A_4 among others. In this report, I will survey this topic, offering a basic recipe for constructing models, with an extensive (but nevertheless incomplete) list of references.

Finite Groups

To obtain a non-Abelian discrete symmetry, a simple heuristic way is to choose two specific noncommuting matrices and form all possible products. As a first example, consider the two 2×2 matrices:

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} \omega & 0 \\ 0 & \omega^{-1} \end{pmatrix}, \quad (1)$$

where $\omega^n = 1$, i.e. $\omega = \exp(2\pi i/n)$. Since $A^2 = 1$ and $B^n = 1$, this group contains Z_2 and Z_n . For $n = 1, 2$, we obtain Z_2 and $Z_2 \times Z_2$ respectively, which are Abelian. For $n = 3$, the group generated has 6 elements and is in fact the smallest non-Abelian finite group S_3 , the permutation group of 3 objects. This particular representation is not the one found in text books, but is related to it by a unitary transformation [1], and was first used in 1990 for a model of quark mass matrices [2, 3]. For $n = 4$, the group generated has 8 elements which are in fact $\pm 1, \pm\sigma_1, \pm\sigma_2, \pm i\sigma_3$, where $\sigma_{1,2,3}$ are the usual Pauli spin matrices. This forms the group D_4 , i.e. the symmetry group of the square, which was first used in 2003 [4, 5]. If the 8 elements are $\pm 1, \pm i\sigma_{1,2,3}$ instead, then they form the group of quaternions Q , which

has also been used [6] for quark and lepton mass matrices. In general, the groups generated by Eq. (1) have $2n$ elements and may be denoted as $\Delta(2n)$.

Consider next the two 3×3 matrices:

$$A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} \omega & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega^{-3} \end{pmatrix}. \quad (2)$$

Since $A^3 = 1$ and $B^n = 1$, this group contains Z_3 and Z_n . For $n = 1$, we obtain Z_3 . For $n = 2$, the group generated has 12 elements and is A_4 , the even permutation group of 4 objects, which was first used in 2001 in a model of lepton mass matrices [7, 8]. It is also the symmetry group of the tetrahedron, one of five perfect geometric solids, identified by Plato with the element “fire” [9]. In general, the groups generated by Eq. (2) have $3n^2$ elements and may be denoted as $\Delta(3n^2)$ [10]. They are in fact subgroups of $SU(3)$. In particular, $\Delta(27)$ has also been used [11, 12]. Generalizing to $k \times k$ matrices, we then have the series $\Delta(kn^{k-1})$. However, since there are presumably only 3 families, $k > 3$ is probably not of much interest.

Going back to $k = 2$, but using instead the following two matrices:

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} \omega & 0 \\ 0 & 1 \end{pmatrix}. \quad (3)$$

Now again $A^2 = 1$ and $B^n = 1$, but the group generated will have $2n^2$ elements. Call it $\Sigma(2n^2)$. For $n = 1$, it is just Z_2 . For $n = 2$, it is D_4 again. For $k = 3$, consider

$$A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} \omega & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (4)$$

then the groups generated have $3n^3$ elements and may be denoted as $\Sigma(3n^3)$. They are in fact subgroups of $U(3)$. For $n = 1$, it is just Z_3 . For $n = 2$, it is $A_4 \times Z_2$. For $n = 3$, the group $\Sigma(81)$ has been used [13] to understand the Koide formula [14] as well as lepton mass matrices [15]. In general, we have the series $\Sigma(kn^k)$.

Model Recipe

(I) Choose a group, e.g. S_3 or A_4 , and write down its possible representations. For example S_3 has $\underline{1}$, $\underline{1}'$, $\underline{2}$; A_4 has $\underline{1}$, $\underline{1}'$, $\underline{1}''$, $\underline{3}$. Work out all product decompositions. For example $\underline{2} \times \underline{2} = \underline{1} + \underline{1}' + \underline{2}$ in S_3 , and $\underline{3} \times \underline{3} = \underline{1} + \underline{1}' + \underline{1}'' + \underline{3} + \underline{3}$ in A_4 .

(II) Assign $(\nu, l)_{1,2,3}$ and $l_{1,2,3}^c$ to the representations of your choice. If you want to consider only renormalizable interactions, you will need to add Higgs doublets (and perhaps also triplets and singlets) and assign them. You may also want to consider adding neutrino singlets.

(III) Because of your choice of particle content and their representations, the Yukawa structure of your model is restricted. As the Higgs bosons acquire vacuum expectation values (which may be related because of some extra or residual symmetry), your lepton mass matrices will have certain particular forms. If the number of parameters involved are less than the number of observables, you then have one or more predictions. Of course, the forms themselves have to be consistent with the known values of m_e , m_μ , m_τ , etc.

(IV) Because there will be more than one Higgs doublet in such models, flavor nonconservation will appear at some level. You will need to work out its phenomenological consequences, making sure that your model is consistent with present experimental constraints. You can then proceed to explore its observability at the TeV scale.

(V) If you insist on using only the one Higgs doublet of the Standard Model, then you must consider effective nonrenormalizable interactions to support the discrete flavor symmetry. In such models, there are no predictions beyond the forms of the mass matrices themselves.

(VI) Quarks can be considered in the same way. The two quark mass matrices \mathcal{M}_u and \mathcal{M}_d must be nearly aligned so that their mixing matrix involves only small angles. In contrast, \mathcal{M}_ν and \mathcal{M}_l should have different structures so that large angles can be obtained.

S_3

Being the simplest, the non-Abelian discrete symmetry S_3 was used already [16] in the early days of strong interactions. There are many recent applications [17, 18, 19, 20, 21, 22, 23, 24, 25, 26], some of which are discussed in my talk at VI-Silafae [27]. Typically, such models often require extra symmetries beyond S_3 to reduce the number of parameters, or assumptions of how S_3 is spontaneously and softly broken. For illustration, consider the model of Kubo et al. [17] which has recently been updated by Felix et al. [28]. The symmetry used is actually $S_3 \times Z_2$, with the assignments

$$(\nu, l), \quad l^c, \quad N, \quad (\phi^+, \phi^0) \sim \underline{1} + \underline{2}, \quad (5)$$

and equal vacuum expectation values for the two Higgs doublets transforming as $\underline{2}$ under S_3 . The Z_2 symmetry serves to eliminate 4 Yukawa couplings otherwise allowed by S_3 , resulting in an inverted ordering of neutrino masses with

$$\theta_{23} \simeq \pi/4, \quad \theta_{13} \simeq 0.0034, \quad m_{ee} \simeq 0.05 \text{ eV}, \quad (6)$$

where m_{ee} is the effective Majorana neutrino mass measured in neutrinoless double beta decay. This model relates θ_{13} to the ratio m_e/m_μ .

A_4

To understand why quarks and leptons have very different mixing matrices, A_4 turns out to be very useful. It allows the two different quark mass matrices to be diagonalized by the same unitary transformations, implying thus no mixing as a first approximation, but because of the assumed Majorana nature of the neutrinos, a large mismatch may occur in the lepton sector, thus offering the possibility of obtaining the so-called tribimaximal mixing matrix

[29, 30], which is a good approximation to present data. One way of doing this is to consider the decomposition

$$U_{l\nu} = \begin{pmatrix} \sqrt{2/3} & 1/\sqrt{3} & 0 \\ -1/\sqrt{6} & 1/\sqrt{3} & -1/\sqrt{2} \\ -1/\sqrt{6} & 1/\sqrt{3} & 1/\sqrt{2} \end{pmatrix} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 \\ 1/\sqrt{2} & 0 & -i/\sqrt{2} \\ 1/\sqrt{2} & 0 & i/\sqrt{2} \end{pmatrix}, \quad (7)$$

where $\omega = \exp(2\pi i/3) = -1/2 + i\sqrt{3}/2$. The matrix involving ω has equal moduli for all its entries and was conjectured already in 1978 [31, 32] to be a possible candidate for the 3×3 neutrino mixing matrix.

Since $U_{l\nu} = U_l^\dagger U_\nu$, where U_l, U_ν diagonalize $\mathcal{M}_l \mathcal{M}_l^\dagger, \mathcal{M}_\nu \mathcal{M}_\nu^\dagger$ respectively, Eq. (7) may be obtained if we have

$$U_l^\dagger = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \quad (8)$$

and

$$\begin{aligned} \mathcal{M}_\nu &= \begin{pmatrix} a+2b & 0 & 0 \\ 0 & a-b & d \\ 0 & d & a-b \end{pmatrix} \\ &= \begin{pmatrix} 0 & 1 & 0 \\ 1/\sqrt{2} & 0 & -i/\sqrt{2} \\ 1/\sqrt{2} & 0 & i/\sqrt{2} \end{pmatrix} \begin{pmatrix} a-b+d & 0 & 0 \\ 0 & a+2b & 0 \\ 0 & 0 & -a+b+d \end{pmatrix} \begin{pmatrix} 0 & 1/\sqrt{2} & 1/\sqrt{2} \\ 1 & 0 & 0 \\ 0 & -i/\sqrt{2} & i/\sqrt{2} \end{pmatrix}. \end{aligned} \quad (9)$$

It was discovered in Ref. [7] that Eq. (8) is naturally obtained with A_4 if

$$(\nu, l)_{1,2,3} \sim \underline{3}, \quad l_{1,2,3}^c \sim \underline{1} + \underline{1}' + \underline{1}'', \quad (\phi^+, \phi^0)_{1,2,3} \sim \underline{3} \quad (11)$$

for $\langle \phi_1^0 \rangle = \langle \phi_2^0 \rangle = \langle \phi_3^0 \rangle$. This assignment also allows m_e, m_μ, m_τ to take on arbitrary values because there are here exactly three independent Yukawa couplings invariant under A_4 . If we use this also for quarks [8], then U_u^\dagger and U_d^\dagger are also given by Eq. (8), resulting in $V_{CKM} = 1$, i.e. no mixing. This should be considered as a good first approximation because the observed mixing angles are all small. In the general case without any symmetry, we would have expected U_u and U_d to be very different.

It was later discovered in Ref. [33] that Eq. (9) may also be obtained with A_4 , using two further assumptions. Consider the most general 3×3 Majorana mass matrix in the form

$$\mathcal{M}_\nu = \begin{pmatrix} a+b+c & f & e \\ f & a+b\omega+c\omega^2 & d \\ e & d & a+b\omega^2+c\omega \end{pmatrix}, \quad (12)$$

where a comes from $\underline{1}$, b from $\underline{1}'$, c from $\underline{1}''$, and (d, e, f) from $\underline{3}$ of A_4 . To get Eq. (9), we need $e = f = 0$, i.e. the effective scalar A_4 triplet responsible for neutrino masses should have its vacuum expectation value along the (1,0,0) direction, whereas that responsible for charged-lepton masses should be (1,1,1) as I remarked earlier. This misalignment is a technical challenge to all such models [34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45]. The other requirement is that $b = c$. Since they come from different representations of A_4 , this is rather *ad hoc*. A very clever solution [34, 39] is to eliminate both, i.e. $b = c = 0$. This results in a normal ordering of neutrino masses with the prediction [36]

$$|m_{\nu_e}|^2 \simeq |m_{ee}|^2 + \Delta m_{atm}^2/9. \quad (13)$$

Other applications [46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62] of A_4 have also been considered. A natural (spinorial) extension of A_4 is the binary tetrahedral group [63, 64] which is under active current discussion [65, 66, 67, 68].

Others

Other recent applications of non-Abelian discrete flavor symmetries include those of D_4 [4, 5, 69], Q_4 [6], D_5 [70, 71], D_6 [72], Q_6 [73, 74, 75], D_7 [76], S_4 [77, 78, 79, 80, 81], $\Delta(27)$ [11, 12], $\Delta(75)$ [82, 83], $\Sigma(81)$ [13, 15], and $B_3 \times Z_2^3$ [84, 85] which has 384 elements.

This work was supported in part by the U. S. Department of Energy under Grant No. DE-FG03-94ER40837.

References

- [1] E. Ma, arXiv:hep-ph/0409075.
- [2] E. Ma, Phys. Rev. D **43** (1991) 2761.
- [3] N. G. Deshpande, M. Gupta and P. B. Pal, Phys. Rev. D **45** (1992) 953.
- [4] W. Grimus and L. Lavoura, Phys. Lett. B **572** (2003) 189 [arXiv:hep-ph/0305046].
- [5] W. Grimus, A. S. Joshipura, S. Kaneko, L. Lavoura and M. Tanimoto, JHEP **0407** (2004) 078 [arXiv:hep-ph/0407112].
- [6] M. Frigerio, S. Kaneko, E. Ma and M. Tanimoto, Phys. Rev. D **71** (2005) 011901 [arXiv:hep-ph/0409187].
- [7] E. Ma and G. Rajasekaran, Phys. Rev. D **64** (2001) 113012 [arXiv:hep-ph/0106291].
- [8] K. S. Babu, E. Ma and J. W. F. Valle, Phys. Lett. B **552** (2003) 207 [arXiv:hep-ph/0206292].
- [9] E. Ma, Mod. Phys. Lett. A **17** (2002) 2361 [arXiv:hep-ph/0211393].
- [10] C. Luhn, S. Nasri and P. Ramond, arXiv:hep-th/0701188.
- [11] E. Ma, Mod. Phys. Lett. A **21** (2006) 1917 [arXiv:hep-ph/0607056].
- [12] I. de Medeiros Varzielas, S. F. King and G. G. Ross, Phys. Lett. B **648** (2007) 201 [arXiv:hep-ph/0607045].
- [13] E. Ma, arXiv:hep-ph/0612022.
- [14] Y. Koide, Lett. Nuovo Cim. **34** (1982) 201.
- [15] E. Ma, arXiv:hep-ph/0701016.
- [16] Y. Yamaguchi, Phys. Lett. **9** (1964) 281.
- [17] J. Kubo, A. Mondragon, M. Mondragon and E. Rodriguez-Jauregui, Prog. Theor. Phys. **109** (2003) 795 [Erratum-ibid. **114** (2005) 287] [arXiv:hep-ph/0302196].

- [18] J. Kubo, H. Okada and F. Sakamaki, Phys. Rev. D **70** (2004) 036007 [arXiv:hep-ph/0402089].
- [19] S. L. Chen, M. Frigerio and E. Ma, Phys. Rev. D **70** (2004) 073008 [Erratum-ibid. D **70** (2004) 079905] [arXiv:hep-ph/0404084].
- [20] W. Grimus and L. Lavoura, JHEP **0508** (2005) 013 [arXiv:hep-ph/0504153].
- [21] L. Lavoura and E. Ma, Mod. Phys. Lett. A **20** (2005) 1217 [arXiv:hep-ph/0502181].
- [22] T. Teshima, Phys. Rev. D **73** (2006) 045019 [arXiv:hep-ph/0509094].
- [23] Y. Koide, Phys. Rev. D **73** (2006) 057901 [arXiv:hep-ph/0509214].
- [24] R. N. Mohapatra, S. Nasri and H. B. Yu, Phys. Lett. B **639** (2006) 318 [arXiv:hep-ph/0605020].
- [25] S. Morisi and M. Picariello, Int. J. Theor. Phys. **45** (2006) 1267 [arXiv:hep-ph/0505113].
- [26] S. Kaneko, H. Sawanaka, T. Shingai, M. Tanimoto and K. Yoshioka, Prog. Theor. Phys. **117** (2007) 161 [arXiv:hep-ph/0609220].
- [27] E. Ma, arXiv:hep-ph/0612013.
- [28] O. Felix, A. Mondragon, M. Mondragon and E. Peinado, arXiv:hep-ph/0610061.
- [29] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. B **530** (2002) 167 [arXiv:hep-ph/0202074].
- [30] X. G. He and A. Zee, Phys. Lett. B **560** (2003) 87 [arXiv:hep-ph/0301092].
- [31] N. Cabibbo, Phys. Lett. B **72** (1978) 333.
- [32] L. Wolfenstein, Phys. Rev. D **18** (1978) 958.
- [33] E. Ma, Phys. Rev. D **70** (2004) 031901 [arXiv:hep-ph/0404199].
- [34] G. Altarelli and F. Feruglio, Nucl. Phys. B **720** (2005) 64 [arXiv:hep-ph/0504165].
- [35] K. S. Babu and X. G. He, arXiv:hep-ph/0507217.
- [36] E. Ma, Phys. Rev. D **72** (2005) 037301 [arXiv:hep-ph/0505209].

- [37] A. Zee, Phys. Lett. B **630** (2005) 58 [arXiv:hep-ph/0508278].
- [38] E. Ma, Phys. Rev. D **73** (2006) 057304 [arXiv:hep-ph/0511133].
- [39] G. Altarelli and F. Feruglio, Nucl. Phys. B **741** (2006) 215 [arXiv:hep-ph/0512103].
- [40] X. G. He, Y. Y. Keum, R. R. Volkas, JHEP **0604** (2006) 039 [arXiv:hep-ph/0601001].
- [41] B. Adhikary, B. Brahmachari, A. Ghosal, E. Ma and M. K. Parida, Phys. Lett. B **638** (2006) 345 [arXiv:hep-ph/0603059].
- [42] B. Adhikary and A. Ghosal, Phys. Rev. D **75** (2007) 073020 [arXiv:hep-ph/0609193].
- [43] F. Yin, Phys. Rev. D **75** (2007) 073010 [arXiv:0704.3827 [hep-ph]].
- [44] G. Altarelli, F. Feruglio and Y. Lin, Nucl. Phys. B **775** (2007) 31 [arXiv:hep-ph/0610165].
- [45] X. G. He, Nucl. Phys. Proc. Suppl. **168** (2007) 350 [arXiv:hep-ph/0612080].
- [46] E. Ma, Mod. Phys. Lett. A **17** (2002) 289 [arXiv:hep-ph/0201225].
- [47] E. Ma, Mod. Phys. Lett. A **17** (2002) 627 [arXiv:hep-ph/0203238].
- [48] K. S. Babu, T. Kobayashi and J. Kubo, Phys. Rev. D **67** (2003) 075018 [arXiv:hep-ph/0212350].
- [49] S. L. Chen, M. Frigerio and E. Ma, Nucl. Phys. B **724** (2005) 423 [arXiv:hep-ph/0504181].
- [50] M. Hirsch, A. Villanova del Moral, J. W. F. Valle and E. Ma, Phys. Rev. D **72** (2005) 091301 [Erratum-ibid. D **72** (2005) 119904] [arXiv:hep-ph/0507148].
- [51] E. Ma, Mod. Phys. Lett. A **20** (2005) 1953 [arXiv:hep-ph/0502024].
- [52] E. Ma, Mod. Phys. Lett. A **20** (2005) 2601 [arXiv:hep-ph/0508099].
- [53] E. Ma, Mod. Phys. Lett. A **20** (2005) 2767 [arXiv:hep-ph/0506036].
- [54] E. Ma, H. Sawanaka and M. Tanimoto, Phys. Lett. B **641** (2006) 301 [arXiv:hep-ph/0606103].

- [55] E. Ma, Mod. Phys. Lett. A **21** (2006) 2931 [arXiv:hep-ph/0607190].
- [56] E. Ma, Mod. Phys. Lett. A **22** (2007) 101 [arXiv:hep-ph/0610342].
- [57] L. Lavoura and H. Kuhbock, Mod. Phys. Lett. A **22** (2007) 181 (2007) [arXiv:hep-ph/0610050].
- [58] I. de Medeiros Varzielas, S. F. King and G. G. Ross, Phys. Lett. B **644** (2007) 153 [arXiv:hep-ph/0512313].
- [59] S. F. King and M. Malinsky, Phys. Lett. B **645** (2007) 351 [arXiv:hep-ph/0610250].
- [60] S. Morisi, M. Picariello and E. Torrente-Lujan, Phys. Rev. D **75** (2007) 075015 [arXiv:hep-ph/0702034].
- [61] Y. Koide, arXiv:hep-ph/0701018.
- [62] M. Hirsch, A. S. Joshipura, S. Kaneko and J. W. F. Valle, arXiv:hep-ph/0703046.
- [63] A. Aranda, C. D. Carone and R. F. Lebed, Phys. Lett. B **474** (2000) 170 [arXiv:hep-ph/9910392].
- [64] A. Aranda, C. D. Carone and R. F. Lebed, Phys. Rev. D **62** (2000) 016009 [arXiv:hep-ph/0002044].
- [65] P. D. Carr and P. H. Frampton, arXiv:hep-ph/0701034.
- [66] F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, Nucl. Phys. B **775** (2007) 120 [arXiv:hep-ph/0702194].
- [67] M. C. Chen and K. T. Mahanthappa, arXiv:0705.0714 [hep-ph].
- [68] P. H. Frampton and T. W. Kephart, arXiv:0706.1186 [hep-ph].
- [69] T. Kobayashi, H. P. Nilles, F. Ploger, S. Raby and M. Ratz, Nucl. Phys. B **768** (2007) 135 [arXiv:hep-ph/0611020].
- [70] E. Ma, Fizika B **14** (2005) 35 [arXiv:hep-ph/0409288].
- [71] C. Hagedorn, M. Lindner and F. Plentinger, Phys. Rev. D **74** (2006) 025007 [arXiv:hep-ph/0604265].

- [72] Y. Kajiyama, J. Kubo and H. Okada, Phys. Rev. D **75** (2007) 033001 [arXiv:hep-ph/0610072].
- [73] P. H. Frampton and T. W. Kephart, Phys. Rev. D **51** (1995) 1 [arXiv:hep-ph/9409324].
- [74] K. S. Babu and J. Kubo, Phys. Rev. D **71** (2005) 056006 [arXiv:hep-ph/0411226].
- [75] J. Kubo, Phys. Lett. B **622** (2005) 303 [arXiv:hep-ph/0506043].
- [76] S. L. Chen and E. Ma, Phys. Lett. B **620** (2005) 151 [arXiv:hep-ph/0505064].
- [77] E. Ma, Phys. Lett. B **632** (2006) 352 [arXiv:hep-ph/0508231].
- [78] C. Hagedorn, M. Lindner and R. N. Mohapatra, JHEP **0606** (2006) 042 [arXiv:hep-ph/0602244].
- [79] Y. Cai and H. B. Yu, Phys. Rev. D **74** (2006) 115005 [arXiv:hep-ph/0608022].
- [80] H. Zhang, arXiv:hep-ph/0612214.
- [81] Y. Koide, arXiv:0705.2275 [hep-ph].
- [82] D. B. Kaplan and M. Schmaltz, Phys. Rev. D **49** (1994) 3741 [arXiv:hep-ph/9311281].
- [83] M. Schmaltz, Phys. Rev. D **52** (1995) 1643 [arXiv:hep-ph/9411383].
- [84] W. Grimus and L. Lavoura, JHEP **0601** (2006) 018 [arXiv:hep-ph/0509239].
- [85] W. Grimus and L. Lavoura, arXiv:hep-ph/0611149.