

Charged hadron R_{AA} as a function of p_T at LHC

T. Renk and K. J. Eskola

Department of Physics, P.O. Box 35 FI-40014 University of Jyväskylä, Finland and
Helsinki Institute of Physics, P.O. Box 64 FI-00014, University of Helsinki, Finland

E-mail: trenk@phys.jyu.fi

Abstract. We compute the nuclear suppression factor R_{AA} for charged hadrons within a radiative energy loss picture using a hydrodynamical evolution to describe the soft medium inducing energy loss. A minijet + saturation picture provides initial conditions for LHC energies and leading order perturbative QCD (LO pQCD) is used to compute the parton spectrum before distortion by energy loss.

We calculate the suppression of hard hadrons induced by the presence of a soft medium produced in central Pb-Pb collisions at $\sqrt{s} = 5.5$ TeV at the LHC. Note that this prediction depends on knowledge of the medium. In the present calculation, the medium evolution is likewise predicted and has to be confirmed before the suppression can be tested. Note further that the calculation is only valid where hadron production is dominated by fragmentation and that it cannot be generalized to the suppression of jets since the requirement of observing a hard hadron leads to showers in which the momentum flow is predominantly through a single parton. This is not so for jets in which the momentum flow is shared on average among several partons (which requires a different framework).

We describe the soft medium evolution by the boost-invariant hydrodynamical model discussed in [1] where the initial conditions for LHC are computed from perturbative QCD+saturation [2]. Our calculation for the propagation of partons through the medium follows the BDMPS formalism for radiative energy loss using quenching weights [3]. Details of the implementation can be found in [4].

The probability density $P(x_0, y_0)$ for finding a hard vertex at the transverse position $\mathbf{r}_0 = (x_0, y_0)$ and impact parameter \mathbf{b} is given by the normalized product of the nuclear profile functions. We compute the energy loss probability $P(\Delta E)_{path}$ for any given path from a vertex through the medium by evaluating the line integrals

$$\omega_c(\mathbf{r}_0, \phi) = \int_0^\infty d\xi \xi \hat{q}(\xi) \quad \text{and} \quad \langle \hat{q}L \rangle(\mathbf{r}_0, \phi) = \int_0^\infty d\xi \hat{q}(\xi). \quad (1)$$

Along the path where we assume the relation

$$\hat{q}(\xi) = K \cdot 2 \cdot \epsilon^{3/4}(\xi) (\cosh \rho - \sinh \rho \cos \alpha) \quad (2)$$

between the local transport coefficient $\hat{q}(\xi)$, the energy density ϵ and the local flow rapidity ρ as given in the hydrodynamical model. The angle α is between flow and

parton trajectory. We view the constant K as a tool to account for the uncertainty in the selection of α_s and possible non-perturbative effects increasing the quenching power of the medium (see [4]) and adjust it such that pionic R_{AA} for central Au-Au collisions at RHIC is described. The result for LHC is then an extrapolation with K fixed.

Using the numerical results of [3], we obtain $P(\Delta E; \omega_c, R)_{path}$ for ω_c and $R = 2\omega_c^2 / \langle \hat{q}L \rangle$. From this distribution given a single path, we can define the averaged energy loss probability distribution $P(\Delta E)_{TAA}$ by averaging over all possible paths, weighted with the probability density $P(x_0, y_0)$ for finding a hard vertex in the transverse plane.

We consider all partons as absorbed whose energy loss is formally larger than their initial energy. The momentum spectrum of produced partons is calculated in LO pQCD. The medium-modified perturbative production of hadrons is obtained from the convolution

$$d\sigma_{med}^{AA \rightarrow h+X} = \sum_f d\sigma_{vac}^{AA \rightarrow f+X} \otimes \langle P(\Delta E) \rangle_{TAA} \otimes D_{f \rightarrow h}^{vac}(z, \mu_F^2) \quad (3)$$

with $D_{f \rightarrow h}^{vac}(z, \mu_F^2)$ the fragmentation function. From this we compute the nuclear modification factor R_{AA} as

$$R_{AA}(p_T, y) = \frac{dN_{AA}^h / dp_T dy}{T_{AA}(\mathbf{b}) d\sigma^{pp} / dp_T dy}. \quad (4)$$




Figure 1. Expectation for the p_T dependence of the nuclear suppression factor R_{AA} for charged hadrons in central Pb-Pb collisions at midrapidity at the LHC.

Fig. 1 shows the expected behaviour of R_{AA} with hadronic transverse momentum p_T at midrapidity. On quite general grounds, we expect a rise of R_{AA} with p_T for any energy loss model in which the energy loss probability does not strongly depend on the initial parton energy as more of the shift in energy becomes accessible (see [4]). The detailed form of the rise is then sensitive to the form of $P(\Delta E)_{TAA}$.

References

- [1] K. J. Eskola, H. Honkanen, H. Niemi, P. V. Ruuskanen and S. S. Räsänen, Phys. Rev. C **72** (2005) 044904.
- [2] K. J. Eskola, K. Kajantie, P. V. Ruuskanen and K. Tuominen, Nucl. Phys. B **570** (2000) 379.
- [3] C. A. Salgado and U. A. Wiedemann, Phys. Rev. D **68**, (2003) 014008.
- [4] T. Renk and K. J. Eskola, hep-ph/0610059.