

The infrared behavior of Landau gauge Yang-Mills theory in $d = 2, 3$ and 4 dimensions

Markus Huber¹, Reinhard Alkofer¹, Christian S. Fischer² and Kai Schwenzer¹

¹Institut für Physik, Universität Graz, Universitätsplatz 5, 8010 Graz, Austria

²Institut für Kernphysik, Darmstadt University of Technology,
Schlossgartenstraße 9, 64289 Darmstadt, Germany

Abstract

We develop a general power counting scheme for the infrared limit of Landau gauge $SU(N)$ Yang-Mills theory in arbitrary dimensions. Employing a skeleton expansion, we find that the infrared behavior is qualitatively independent of the spacetime dimension d . In the cases $d = 2, 3$ and 4 even the quantitative results for the infrared exponents of the vertices differ only slightly. Therefore, corresponding lattice simulations provide interesting qualitative information for the physical case. We furthermore find that the loop integrals depend only weakly on the numerical values of the IR exponents.

Introduction: The infrared (IR) regime of Yang-Mills theory determines the fundamental non-perturbative properties of the non-Abelian gauge dynamics. In particular, the IR limit of Greens functions of the colored fields are likely connected to the problem of confinement as demonstrated within the scenarios of Kugo-Ojima [1] and Gribov-Zwanziger [2, 3]. Due to the intricate nature of the problem an investigation via all available methods is desirable. Such combined efforts via Dyson-Schwinger equations (DSE) [4, 5, 6, 7], renormalization group (RG) techniques [8] and lattice gauge theory studies [9] led during the last years to a coherent picture of the infrared regime in Landau gauge. Hereby, in the continuum approaches, the IR scaling limit of general Greens functions in four spacetime dimensions has been determined [10]. It exhibits, as argued for in [11], a dominance of the gauge fixing part of the action, i.e. the Faddeev-Popov ghost. A simultaneous analysis of DSE and RG methods allowed furthermore to show that this IR fixpoint is unique [12].

The Greens functions of Yang-Mills theory in Landau gauge are studied extensively on the lattice, see e.g. [9]. These analyses show the IR increase of the ghost dressing function required in both the Kugo-Ojima and the Gribov-Zwanziger scenario, as well as the IR scaling of the gluon dressing function in accordance with DSE and RG analyses. The influence of finite volume corrections has been studied within DSE analyses on compact manifolds [13] which agree qualitatively with the finite IR limit of current lattice results. However, a confirmation that even the gluon propagator vanishes weakly in the IR, which is necessary in the Gribov-Zwanziger scenario, has not yet been possible in four-dimensional lattice simulations due to the large lattice sizes required to probe the IR regime. The situation is similar for the important but even harder determination of the scaling behavior of vertex functions or the quantitative extraction of IR exponent. Lower dimensional lattices could allow to test qualitative aspects of these Greens functions in a substantially simpler setting. Naturally, such a surrogate study requires that the generic features are identical to those in four-dimensional spacetime. Recent lattice studies in two [14] and three spacetime dimensions [15] indeed suggest a power-law behavior similar to the four dimensional case.

$SU(N)$ Yang-Mills theory in three spacetime dimensions is also relevant from two other points of view. In the high temperature limit only the lowest Matsubara mode contributes, leading to a dimensional reduction and thus to an effective field theory in three dimensions. The qualitative aspects of this effective theory, in particular with regard to its confining nature, may be covered by ordinary three-dimensional gauge theory [16]. Similarly, Yang-Mills theory in the canonical quantization approach in Coulomb gauge

[17] with an appropriate choice for the vacuum wave functional is structurally similar to the corresponding three-dimensional theory in Landau gauge.

These results motivate to study the dependence of the infrared limit on the parameters of the theory in more detail. It will turn out that the infrared solution of Yang-Mills theory is determined by the spacetime dimension d and one parameter κ that determines the IR exponents of propagators and vertices. In this work we establish a manifest power counting scheme for general vertex functions of Landau gauge Yang-Mills theory in arbitrary dimensions. We find qualitatively similar results as in the four-dimensional analysis given in [10]. Furthermore we discuss the dependence of the Dyson-Schwinger equations on the infrared scaling parameter κ . In four dimensions it takes a value $\kappa \approx 0.595$ within an approximation based on the propagator DSEs and using a bare ghost-gluon vertex. A possible dressing of this vertex could change this value slightly [6]. Therefore we study to what extent these dynamical building blocks in the DSEs depend on κ , employing an analysis of the IR-dominant integrals.

IR exponents for arbitrary d : In the following we will perform a scaling analysis for the IR regime of Yang-Mills theory in arbitrary dimensions. In contrast to the generic IR limit we study the limit where the coupling g is kept fixed and has no inherent scaling dependence. Instead in dimensions other than four it fixes the fundamental scale of the theory. This limit is more directly accessible in lattice simulations and has been analyzed in [14, 15]. A similar IR analysis has previously been performed for the physically important case $d=4$ in refs. [10, 12]. Although the scaling in arbitrary dimension may be abstracted from these results via the consideration of the appropriate canonical scaling dimensions, we chose to give a concise and nevertheless self-contained derivation of the general scaling relations and refer the reader to the corresponding work for details. We note, that our analysis is immediately applicable to the gauge sector of QCD since closed quark loops are IR-suppressed due to the finite current quark masses [18] and therefore there are *no* quark contributions to purely bosonic Greens functions in the IR regime. This has been verified in ref. [19].

Analog to the corresponding analysis in four dimensions [10], the starting point for the IR analysis is the non-renormalization of the ghost-gluon vertex in Landau gauge [20], which implies a finite vertex in the infrared [10]. This property depends purely on the transversality of the gluon propagator and is, as long as the ghost-gluon scattering kernel is not strongly IR-divergent, therefore valid in arbitrary dimensions. We come back to this point below.

$$\text{---}\bullet\text{---}^{-1} = \text{---}^{-1} - \text{---}\bullet\text{---} \text{---} \text{---}\bullet\text{---}$$

Figure 1: The DSE for the ghost propagator.

The infrared behavior of the propagators and vertices well below its inherent scale $g^{2/(4-d)}$ (respectively Λ_{QCD} in $d=4$) is determined via renormalization group arguments by scaling relations. The propagators of the gluons and ghosts

$$D_{\mu\nu}(p^2) = \left(\delta_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \right) \frac{Z(p^2)}{p^2}, \quad D^G(p^2) = -\frac{G(p^2)}{p^2}, \quad (1)$$

are given in terms of dressing functions whose IR behavior is described by a power law ansatz

$$Z(p^2) = c_{0,2} \cdot (p^2)^{\delta_{0,2}}, \quad G(p^2) = c_{2,0} \cdot (p^2)^{\delta_{2,0}}, \quad (2)$$

and similar for the vertices. Here we denote the IR exponent of a vertex with $2n$ ghost and m gluon legs by $\delta_{2n,m}$ and the corresponding coefficient by $c_{2n,m}$. Whereas this coefficient is a constant for the propagators it is generally a function of $2n-m-1$ momentum ratios.

For the integral on the right hand side of the ghost propagator Dyson-Schwinger equation, cf. fig. 1, one can use the standard expression [6, 21]

$$\int \frac{d^d q}{(2\pi)^d} (q^2)^{\nu_1} ((q-p)^2)^{\nu_2} = (4\pi)^{-\frac{d}{2}} \frac{\Gamma(\frac{d}{2} + \nu_1) \Gamma(\frac{d}{2} + \nu_2) \Gamma(-\nu_1 - \nu_2 - \frac{d}{2})}{\Gamma(-\nu_1) \Gamma(-\nu_2) \Gamma(d + \nu_1 + \nu_2)} (p^2)^{\frac{d}{2} + \nu_1 + \nu_2} \quad (3)$$




Figure 2: 1-loop part of the skeleton expansion of the DSE for the three-gluon vertex.

which shows that it scales proportional to the external momentum. The left hand side of the ghost DSE, which consists only of the inverse dressed ghost propagator, is proportional to $(p^2)^{-\delta_{2,0}+1}$. The 1 comes from the canonical dimension of the ghost propagator. The scaling dimensions on the right hand side are $d/2$ from the integral, $\delta_{0,2} - 1$ from the gluon propagator, $\delta_{2,0} - 1$ from the ghost propagator, $1/2$ from the bare ghost-gluon vertex, and $1/2$ from the dressed ghost-gluon vertex which features no anomalous scaling. This yields as condition for the IR exponents

$$1 - \delta_{2,0} = \frac{d}{2} + \delta_{0,2} - 1 + \delta_{2,0} - 1 + \frac{1}{2} + \frac{1}{2}. \quad (4)$$

Defining the parameter κ as $\kappa := -\delta_{2,0}$, we recover in four dimensions the well-known result $\delta_{0,2} = 2\kappa$. In d dimensions we have [5, 6]

$$\delta_{2,0} = -\kappa, \quad \delta_{0,2} = 2\kappa + 2 - \frac{d}{2}. \quad (5)$$

A comparison between different dimensions via κ is not possible directly, because κ has different values for different d [5, 6, 16]. We discuss this point further below.

In order to transform the infinite hierarchy of Dyson-Schwinger equations into a closed system we perform a skeleton expansion. This yields an infinite tower of graphs involving only primitively divergent vertices which we will analyze in a first step. The first order of the skeleton expansion of the DSE for the three-gluon vertex is depicted in fig. 2. We start with the ghost triangle which turns out to be one of the infrared leading diagrams. We have $d/2$ from the integral, $3(-\kappa - 1)$ from the three ghost propagators and $3/2$ from the three ghost-gluon vertices. We subtract the canonical dimension $1/2$ to get the anomalous IR exponent of the ghost triangle of the three-gluon vertex:

$$\delta_{0,3}^{gh\Delta} = \frac{d}{2} + 3(-\kappa - 1) + 3\frac{1}{2} - \frac{1}{2} = -3\kappa + \frac{d}{2} - 2. \quad (6)$$

For the four-gluon vertex we can apply the same procedure. In fig. 3 we show the first order of its skeleton expansion. Here, also the ghost rectangle gives one of the IR dominant contributions. Simple counting of the powers yields for the IR exponent of this diagram

$$\delta_{0,4}^{gh\Box} = \frac{d}{2} + 4(-\kappa - 1) + 4\frac{1}{2} = -4\kappa + \frac{d}{2} - 2. \quad (7)$$

Now we are in the position to derive the IR exponents of the other diagrams of the three-gluon vertex DSE in fig. 2. The gluon triangle contains three propagators as well as two dressed and one bare three-gluon vertex, so including the loop integral and subtracting the canonical dimension the IR exponent is $d/2 + 3(2\kappa + 1 - d/2) + 2(-3\kappa + d/2 - 3/2) + 1/2 - 1/2 = 0$. Correspondingly, the other corrections, including the two-loop diagrams that are not shown, are likewise subleading compared to the dominant




Figure 3: 1-loop part of the skeleton expansion of the DSE for the four-gluon vertex.

ghost loop. The same holds for the four-gluon vertex. In general, the IR exponents of these purely gluonic diagrams are *independent* of the dimension and therefore identical to eqs. (13) and (15) in [10].

With the scaling information of all primitively divergent Greens functions at hand we have all parts to calculate the IR exponent of an arbitrary n -point function. The different objects that can appear in corresponding general graphs are given in table 1. In terms of these building blocks the IR exponent of an arbitrary vertex v is given by

$$\begin{aligned} \delta_v = & (l - m_i + v_{0,3} + v_{0,4}) \frac{d}{2} + (2m_i - n_i - 3v_{0,3} - 4v_{0,4})\kappa + \\ & + \frac{1}{2}(2m_i - 2n_i + v_{2,1} + v_{0,3}^b - 3v_{0,3} - 4v_{0,4} - 2v), \end{aligned} \quad (8)$$

where v is the canonical scaling dimension of the vertex. This formula can be simplified using standard relations between the number of loops, the number of vertices and the number of propagators. The number of loops is given in terms of the internal lines and vertices via the topological relation

$$l = m_i + n_i + 1 - (v_{2,1} + v_{0,3}^b + v_{0,3} + v_{0,4}^b + v_{0,4}). \quad (9)$$

Further topological relations that connect the number of lines of a given species with the corresponding vertices allow to rewrite the expression in terms of the number of external lines which are denoted by m and n for gluons and ghost-antighost-pairs, respectively:

$$m + 2m_i = v_{2,1} + 3(v_{0,3}^b + v_{0,3}) + 4(v_{0,4}^b + v_{0,4}), \quad (10)$$

$$n + n_i = v_{2,1}. \quad (11)$$

object	number of objects	scaling dimension
loop	1	$d/2$
internal ghost line	n_i	$\delta_{2,0} - 1 = -\kappa - 1$
internal gluon line	m_i	$\delta_{0,2} - 1 = 2\kappa + 1 - d/2$
ghost-gluon vertex	$v_{2,1}$	$1/2$
bare 3-gluon vertex	$v_{0,3}^b$	$1/2$
dressed 3-gluon vertex	$v_{0,3}$	$\delta_{0,3} + 1/2 = -3\kappa + d/2 - 3/2$
bare 4-gluon vertex	$v_{0,4}^b$	0
dressed 4-gluon vertex	$v_{0,4}$	$\delta_{0,4} = -4\kappa + d/2 - 2$

Table 1: Infrared behavior of the building blocks of general loop diagrams within the skeleton expansion.

The canonical dimension of the vertex is finally given in terms of its external lines

$$v = \frac{4 - (2n + m)}{2} = 2 - n - \frac{m}{2}. \quad (12)$$

Inserting eqs. (9), (10), (11) and (12) into eq. (8) we get

$$\begin{aligned} \delta_{2n,m} = & (-n + 1 - v_{0,4}^b - v_{0,3}^b) \frac{d}{2} + (4v_{0,4}^b + 3v_{0,3}^b - m + n)\kappa + \\ & + (2v_{0,4}^b + 2v_{0,3}^b + 2n - 2). \end{aligned} \quad (13)$$

This formula is still dependent on bare vertices. However, when considering different terms of the skeleton expansion of an n -point function, one easily sees that terms containing bare three- and four-gluon vertices are not the IR-dominant ones because contributions from these terms are

$$v_{0,3}^b(3\kappa - \frac{d}{2} + 2), \quad (14)$$





$$v_{0,4}^b(4\kappa - \frac{d}{2} + 2), \quad (15)$$

and thereby always add a positive power of momenta. Consequently, we can neglect these diagrams, when counting only IR leading ones, i.e. we can set $v_{0,3}^b = v_{0,4}^b = 0$ in (13) and obtain as a final result for the IR exponent of an n -point function the simple expression

$$\delta_{2n,m} = (n - m)\kappa + (1 - n) \left(\frac{d}{2} - 2 \right). \quad (16)$$

This relation verifies a posteriori the assumption discussed above fig. 1: the ghost-gluon scattering kernel is exactly as IR-divergent as to make the ghost-gluon vertex finite in the IR, and this is independent of the value of the dimension d .

It remains to discuss higher order corrections in the skeleton expansion. From eq. (16) we can see immediately that there are infinitely many higher orders terms which have the same IR behavior as the leading term: Insertions which lead to higher orders only contribute with dressed vertices and propagators, and these were all considered in the derivation of eq. (16). These insertions with their corresponding IR exponents are given by:

	$(2\kappa + 1 - \frac{d}{2}) + 2(-\kappa - 1) + 2\frac{1}{2} + \frac{d}{2} = 0$
<p>A</p> 	$2(2\kappa + 1 - \frac{d}{2}) + (-\kappa - 1) + (-3\kappa + \frac{d}{2} - \frac{3}{2}) + \frac{1}{2} + \frac{d}{2} = 0$
<p>B</p> 	$3(2\kappa + 1 - \frac{d}{2}) + 2(-3\kappa + \frac{d}{2} - \frac{3}{2}) + \frac{d}{2} = 0$
<p>C</p> 	$2(2\kappa + 1 - \frac{d}{2}) + (-4\kappa + \frac{d}{2} - 2) + \frac{d}{2} = 0$
<p>D</p>	




Figure 4: The (mostly two) possible solutions for the IR exponent κ as a function of dimension d , cf. also refs. [11] and [16]. The linear fits are exact in some ranges of dimension d and approximate the real solutions to good precision in others. The first branch contains the solutions $\kappa = 0.2$, $\kappa = 0.3976\dots$ and $\kappa = 0.5953\dots$ for $d = 2, 3$ and 4 , respectively, whereas the second one includes the solutions $\kappa = 0$, $\kappa = 0.5$ and $\kappa = 1$ [5].

It should be noted that in contrast to a uniform scaling limit we have kept the (dimensionful) coupling g fixed. The IR power laws, eq. (16), are then valid for momenta $p^2 \ll g^{4/(4-d)}$. The resulting momentum dependent couplings from the ghost-gluon vertex, α^{gh} , the three-gluon vertex, α^{3g} , and the four gluon vertex, α^{4g} , (see [10] for a definition) are then given by

$$\alpha^{gh}(p^2) = \frac{g^2}{4\pi} [Z_{2,1}(p^2)]^2 G^2(p^2) Z(p^2) \xrightarrow{p^2 \rightarrow 0} (p^2)^{(4-d)/2}, \quad (17)$$

$$\alpha^{3g}(p^2) = \frac{g^2}{4\pi} [Z_{0,3}(p^2)]^2 Z^3(p^2) \xrightarrow{p^2 \rightarrow 0} (p^2)^{(4-d)/2}, \quad (18)$$

$$\alpha^{4g}(p^2) = \frac{g^2}{4\pi} [Z_{0,4}(p^2)] Z^2(p^2) \xrightarrow{p^2 \rightarrow 0} (p^2)^{(4-d)/2}. \quad (19)$$

Here $Z_{2,1}$ denotes the dressing function of the tree-level structure of the ghost-gluon vertex and $Z_{0,3}$ and $Z_{0,4}$ the corresponding dressing functions of the three- and four-gluon vertices, respectively. To obtain these results the scaling laws, eq. (16), have been used. From this one may better understand the usefulness of the skeleton expansion: apart from B all diagrammatic pieces given above contain dressing factors of the same multiplicity as the couplings, cf. eqs. (17-19). Thus all anomalous dimensions cancel and the canonical dimension $(4-d)/2$ of these couplings is cancelled by the canonical dimensions and the extra loop. As a result the skeleton expansion indeed works independent of the value of the spacetime dimension d as could have been expected from the previously studied four-dimensional case [4, 5, 6, 10, 12].

Finally, let us compare the IR behavior of the Yang-Mills Greens functions in different dimensions. The corresponding values for the IR exponent κ (using a bare ghost-gluon vertex) are displayed in fig. 4. Apart from the region $d < 1$ we always have two possible values, which arise from an IR analysis of the ghost and gluon propagator DSEs. We show the IR exponents of the ghost and gluon propagators and the three- and four-gluon vertices in table 2. Whereas the situation in four dimensions is not conclusive, yet, the solutions belonging to the first branch in fig. 4 have been found in lattice studies in two and three dimensions. These results for the IR exponents of the propagators, the ghost-gluon vertex [22] and the three-gluon vertex in two dimensions [14] agree within errors with the corresponding values given in table 2. One can see that for this branch the qualitative behavior does not change in different dimensions. The other branch in fig. 4 that starts at $\kappa = -1$ for $d = 0$ and ends at $\kappa = 2$ for $d = 6$ has not been seen in lattice simulations. We will give some additional arguments below that this branch may be unphysical.

Dependence of the loop integrals on the IR-exponent: The Dyson-Schwinger equations involve loop integrals over bare and dressed vertex functions. As just discussed the corresponding dressing functions exhibit a power law with appropriate infrared exponents. The corresponding coefficients depend on the IR exponent κ . For the respective coefficients of the propagators, eq. (2), which can be computed from the 2-point integral, eq. (3), this κ -dependence is shown in fig. 5, where κ -independent prefactors have been dropped. The physical requirement that the dressing functions have to be positive restricts the possible values of κ to $\kappa \leq 0.5$ in two, $0.25 < \kappa \leq 0.75$ in three and $\kappa > 0.5$ in four dimensions [23]. We note that all known solutions fulfill these requirements. The decisive point is, however, that although the curves in fig. 5 differ considerable, the physical values obtained from the Dyson-Schwinger solution are far away from these poles and thereby the κ -dependence in their vicinity is rather mild and qualitatively similar in each case.




Figure 5: The κ -dependence of the unrenormalized dressing integrals appearing in the DSEs for the ghost (left) and gluon propagator (right). The dotted, dashed and solid lines show the curves for $d = 2, 3$ and 4 respectively, see also ref. [6]. The full and open dots represent the solutions of the first and second branch in fig. 4.

In general the vertex integrals feature more complicated tensor structures with an increasing number of independent tensor components. However, there are general methods to decompose such tensor integrals to standard scalar integrals [24]. The tensor integrals in the DSE for the ghost-gluon and three-gluon vertex reduce to 3-point integrals of the form

$$I_3(p, q) \equiv \int \frac{d^d k}{(2\pi)^d} \frac{1}{((k+p)^2)^{\nu_1}} \frac{1}{((k-q)^2)^{\nu_2}} \frac{1}{(k^2)^{\nu_3}} \quad (20)$$

where any additional momenta due to vertex functions in the numerator are included in the denominators for appropriate values ν_i . A general expression for such scalar 1-loop 3-point integrals in arbitrary dimensions and with arbitrary powers of the propagators has been obtained in [21, 25]. An explicit evaluation of this result requires an expression for the resulting Appell function F_4 whose defining series does not converge in the considered Euclidean regime. A converging expression can be obtained by analytic continuation and has been given in [26]. Unfortunately, the result given there is slightly incorrect. The correct and lengthy expression which agrees with a direct numerical integration of these integrals

Dimension		4	3	2	4	3	2
κ		0.5953... ≈ 0.6	0.3976... ≈ 0.4	0.2	1	0.5	0
Ghost	$-\kappa - 1$	-1.6	-1.4	-1.2	-2	-1.5	-1
Gluon	$2\kappa + 1 - d/2$	0.2	0.3	0.4	1	0.5	0
3-gluon	$-3\kappa + d/2 - 3/2$	-1.3	-1.2	-1.1	-1.5	-1.5	-0.5
4-gluon	$-4\kappa + d/2 - 2$	-2.4	-2.1	-1.8	-4	-2.5	-1

Table 2: The dimension dependence of the infrared behavior of Yang-Mills Green functions. For each d the two solutions for κ corresponding to the two alternative branches in fig. 4 are displayed.

will be given in a forthcoming publication [27] where we will also give the detailed results for the different tensor structures and their kinematic dependence. Here we will merely discuss the dependence of the overlap of the tree-level tensor with the IR-dominant ghost triangle. According to the discussed power counting, the ghost loop presents the IR-leading contribution to the Dyson Schwinger equation for the three-gluon vertex in fig. 2. Since this loop involves only ghost-gluon vertices that remain bare to leading order, its IR behavior can be analyzed semi-perturbatively using the scaling form of the dressed ghost propagators, see eq. (2). We consider the special kinematic configuration given by the symmetric point $p^2 = q^2 = r^2$. At this point the dressing function of the leading ghost loop correction to the three-gluon integral scales according to eq. (6) as $Z_{0,3} = c_{0,3}(\kappa) \cdot (p^2)^{-3\kappa+d/2-2}$ where the dependence of the coefficient on the scaling parameter κ has been made explicit. The coefficient is shown as a function of κ in fig. 6, where the full and open points represent the values for the first and second branch in fig. 4 and κ -independent factors have been dropped again.




Figure 6: The κ -dependence of the overlap of the unrenormalized ghost triangle with the tree-level tensor at the symmetric point. The dotted, dashed and solid lines show the curves for $d = 2, 3$ and 4 respectively. The full and open dots represent the solutions of the first and second branch in fig. 4.

The vertex integral has zeros at $\kappa = 0$ in two, $\kappa = 0.5$ in three and $\kappa = 1$ in four dimensions which coincide precisely with the solutions of the second branch. These zeros depend on the kinematic configuration and appear at higher values of κ away from the symmetric point. (The corresponding scalar integrals, cf. eq. (20), feature even poles at these values of κ which precisely cancel for the tree-level tensor.) According to eq. (18) such zeros in the vertex dressing function would lead to an IR-vanishing coupling. Since Yang-Mills theory is apparently a strongly interacting theory, this indicates that these solutions might not be physically relevant. As in the case of the propagators the κ dependence in the vicinity of the solutions of the first branch is similar in all dimensions. The negative value of the three-gluon vertex in two dimensions nevertheless leads to a positive IR limit of the coupling α^{3g} in eq. (18). However, it would cause problems in its renormalization group flow and might be an artifact of the skeleton expansion and cured by higher orders.

Conclusions: We have studied generic features of the IR limit of $SU(N)$ Yang-Mills theory established in [4, 6, 7, 10] in more detail. As in the four-dimensional case, the IR behavior of Greens functions can be extracted via a skeleton expansion in arbitrary dimensions. We find that the IR limit of Greens functions is surprisingly insensitive on the spacetime dimension. This already constitutes the first important result of this study. As a consequence Yang-Mills theory in lower dimensions has a qualitatively similar IR limit as the four dimensional theory. Thus the confinement mechanism might possess identical features in different

dimensions. Corresponding lattice simulations should provide interesting qualitative information for the physical case. Indeed a recent study on large two dimensional lattices [14] finds infrared exponents in agreement with the value $\kappa = 0.2$, thus confirming the Gribov-Zwanziger scenario also in two dimensions.

The results on the mild κ -dependence of the remaining DSE solutions suggest that ghost dominance is a rather robust mechanism and should not depend on the details of the employed truncation scheme. This is further substantiated by the fact that the main non-linearities in the DSE system, which enable the non-trivial fixpoint, arise in the propagator equations whereas there is no non-linear feedback to leading order in the vertex equations.

Acknowledgements: We are grateful to A. Maas and J. Pawłowski for valuable discussions and to A. Maas for a critical reading of the manuscript. This work has been supported in part by the DFG under contract AL279/5-1, by the FWF under contract M979-N16 and by the Helmholtz-University Young Investigator Grant VH-NG-332.

References

- [1] T. Kugo and I. Ojima, Prog. Theor. Phys. Suppl. **66** (1979) 1.
- [2] V. N. Gribov, Nucl. Phys. B **139** (1978) 1.
- [3] D. Zwanziger, Nucl. Phys. B **364** (1991) 127.
- [4] L. von Smekal, R. Alkofer and A. Hauck, Phys. Rev. Lett. **79** (1997) 3591, [arXiv:hep-ph/9705242].
- [5] D. Zwanziger, Phys. Rev. D **65** (2002) 094039, [arXiv:hep-th/0109224].
- [6] C. Lerche and L. von Smekal, Phys. Rev. D **65** (2002) 125006, [arXiv:hep-ph/0202194].
- [7] R. Alkofer and L. von Smekal, Phys. Rept. **353** (2001) 281; C. S. Fischer, J. Phys. G **32** (2006) R253, [arXiv:hep-ph/0007355].
- [8] J. M. Pawłowski, D. F. Litim, S. Nedelko and L. von Smekal, Phys. Rev. Lett. **93** (2004) 152002, [arXiv:hep-th/0312324].
- [9] A. Sternbeck, E. M. Ilgenfritz, M. Mueller-Preussker and A. Schiller, Phys. Rev. D **72** (2005) 014507, [arXiv:hep-lat/0506007]; A. Cucchieri and T. Mendes, Phys. Rev. D **73** (2006) 071502, [arXiv:hep-lat/0602012]; P. O. Bowman *et al.*, [arXiv:hep-lat/0703022]; O. Oliveira and P. J. Silva, [arXiv:0705.0964 [hep-lat]].
- [10] R. Alkofer, C. S. Fischer and F. J. Llanes-Estrada, Phys. Lett. B **611** (2005) 279, [arXiv:hep-th/0412330].
- [11] D. Zwanziger, Phys. Rev. D **69** (2004) 016002, [arXiv:hep-ph/0303028].
- [12] C. S. Fischer and J. M. Pawłowski, Phys. Rev. D **75** (2007) 025012, [arXiv:hep-th/0609009].
- [13] C. S. Fischer, B. Gruter and R. Alkofer, Annals Phys. **321** (2006) 1918, [arXiv:hep-ph/0506053]; C. S. Fischer, A. Maas, J. M. Pawłowski and L. von Smekal, to be published in Annals Phys., [arXiv:hep-ph/0701050].
- [14] A. Maas, to be published in Phys. Rev. D, [arXiv:0704.0722 [hep-lat]].
- [15] A. Cucchieri, A. Maas and T. Mendes, Phys. Rev. D **74** (2006) 014503, [arXiv:hep-lat/0605011]; A. Cucchieri, T. Mendes and A. R. Taurines, Phys. Rev. D **67** (2003) 091502, [arXiv:hep-lat/0302022].
- [16] A. Maas, J. Wambach, B. Gruter and R. Alkofer, Eur. Phys. J. C **37** (2004) 335, [arXiv:hep-ph/0408074].
- [17] W. Schleifenbaum, M. Leder and H. Reinhardt, Phys. Rev. D **73** (2006) 125019, [arXiv:hep-th/0605115].

- [18] T. Appelquist and J. Carazzone, Phys. Rev. D **11** (1975) 2856.
- [19] C. S. Fischer and R. Alkofer, Phys. Rev. D **67** (2003) 094020, [arXiv:hep-ph/0301094].
- [20] J. C. Taylor, Nucl. Phys. B **33** (1971) 436.
- [21] C. Anastasiou, E. W. N. Glover and C. Oleari, Nucl. Phys. B **572** (2000) 307, [arXiv:hep-ph/9907494].
- [22] A. Cucchieri, T. Mendes and A. Mihara, JHEP **0412** (2004) 012, [arXiv:hep-lat/0408034].
- [23] A. Maas, PhD thesis, TU Darmstadt, 2004 [arXiv:hep-ph/0501150].
- [24] A. I. Davydychev, Phys. Lett. B **263** (1991) 107.
- [25] A. I. Davydychev, Vestn. Mosk. Univ. (Ser. 3) 28, No. 3 (1987) 8 [Mosc. Univ. Phys. Bull. 42, No. 3, (1987) 6]; E. E. Boos and A. I. Davydychev, Theor. Math. Phys. **89** (1991) 1052 [Teor. Mat. Fiz. **89** (1991) 56].
- [26] H. Exton, J. Phys. A: Math. Gen. 28 (1995) 631.
- [27] R. Alkofer, M. Huber and K. Schwenzer, in preparation.