

# Nuclear Suppression of Jets and $R_{AA}$ at the LHC

G.Y. Qin, J. Ruppert, S. Turbide, C. Gale and S. Jeon

Department of Physics, McGill University, Montreal, QC, H3A 2T8, Canada

**Abstract.** The nuclear modification factor  $R_{AA}$  for charged hadron production at the LHC is predicted from jet energy loss induced by gluon bremsstrahlung. The Arnold, Moore, and Yaffe [1] formalism is used, together with an ideal hydrodynamical model [2].

We present a calculation of the nuclear modification factor  $R_{AA}$  for charged hadron production as a function of  $p_T$  in Pb+Pb collisions at  $\sqrt{s} = 5.5$  ATeV in central collisions at mid-rapidity at the LHC. The net-energy loss of the partonic jets is calculated by applying the Arnold, Moore, and Yaffe (AMY) formalism to calculate gluon bremsstrahlung [1]. The details of jet suppression relies on an understanding of the nuclear medium, namely the temperatures and flow profiles that are experienced by partonic jets while they interact with partonic matter at  $T \geq T_c$ . Our predictions use a boost-invariant ideal hydrodynamic model with initial conditions calculated from perturbative QCD + saturation [2, 3]. It is emphasized that the reliability of this work hinges on the validity of hydrodynamics at the LHC. It has been verified that  $R_{AA}$  for  $\pi_0$  production as a function of  $p_T$  as obtained in the same boost-invariant ideal hydrodynamical model adjusted to Au+Au collisions at  $\sqrt{s} = 0.2$  ATeV [2] is in agreement with preliminary data from PHENIX in central collisions at RHIC (and the result is very close to the one obtained in 3D hydrodynamics presented in [4]). In AMY the strong coupling constant  $\alpha_s$  is a direct measure of the interaction strength between the jet and the thermalized soft medium and is the only quantity not uniquely determined in the model, once the temperature and flow evolution is fixed by the initial conditions and subsequent hydrodynamical expansion. We found that assuming a constant  $\alpha_s = 0.33$  describes the experimental data in most central collisions at RHIC. It is conjectured that  $\alpha_s$  should not be changed very much at the LHC since the initial temperature is about twice larger than the one at RHIC whereas  $\alpha_s$  is only logarithmically dependent on temperature. We present results for  $\alpha_s = 0.33$  and 0.25.


For details of the calculation of nuclear suppression, we refer the reader to [4]. The extension to the LHC once the medium evolution and  $\alpha_s$  are fixed is straightforward. The initial jets are produced with an initial momentum distribution of jets computed from pQCD in the factorization formalism including nuclear shadowing effects. The probability density  $\mathcal{P}_{AB}(\vec{r}_\perp)$  of finding a hard jet at the transverse position  $\vec{r}_\perp$  in central A+B collisions is given by the normalized product of the nuclear thickness functions,  $\mathcal{P}_{AB}(\vec{r}_\perp) = T_A(\vec{r}_\perp)T_B(\vec{r}_\perp)/T_{AB}$  and is calculated for Pb+Pb collisions. The evolution

of the jet momentum distribution  $P_j(p, t) = dN_j(p, t)/dpdy$  in the medium is calculated by solving a set of coupled rate equations with the following generic form,

$$\frac{dP_j(p, t)}{dt} = \sum_{ab} \int dk \left[ P_a(p+k, t) \frac{d\Gamma_{jb}^a(p+k, p)}{dkdt} - P_j(p, t) \frac{d\Gamma_{ab}^j(p, k)}{dkdt} \right], \quad (1)$$

where  $d\Gamma_{ab}^j(p, k)/dkdt$  is the transition rate for the partonic process  $j \rightarrow a + b$  which depends on the temperature and flow profiles experienced by the jets traversing the medium. The hadron spectrum  $dN_{AB}^h/d^2p_T dy$  is obtained by the fragmentation of jets after their passing through the medium. The nuclear modification factor  $R_{AA}$  is computed as

$$R_{AA}^h(\vec{p}_T, y) = \frac{1}{N_{coll}} \frac{dN_{AA}^h/d^2p_T dy}{dN_{pp}^h/d^2p_T dy}. \quad (2)$$



**Figure 1.** The  $p_T$  dependence of the nuclear modification factor  $R_{AA}$  for charged hadrons in central Pb+Pb collisions at mid-rapidity at the LHC.

In Fig. 1 we present a prediction for charged hadron  $R_{AA}$  as a function of  $p_T$  at mid-rapidity for central collisions at the LHC. We consider that these two values of  $\alpha_s$  define a sensible band of physical parameters.

**Acknowledgments:** We thank the authors of [2] for providing their hydrodynamical evolution calculation at RHIC and LHC energies, T. Renk for discussions, and the Natural Sciences and Engineering Research Council of Canada for support.

## References

- [1] P. Arnold, G. D. Moore and L. G. Yaffe, JHEP **0111**, 057 (2001); JHEP **0112**, 009 (2001); JHEP **0206**, 030 (2002).
- [2] K. J. Eskola, H. Honkanen, H. Niemi, P. V. Ruuskanen and S. S. Räsänen, Phys. Rev. C **72**, 044904 (2005).
- [3] K. J. Eskola, K. Kajantie, P. V. Ruuskanen and K. Tuominen, Nucl. Phys. B **570** (2000) 379.
- [4] G. Y. Qin, J. Ruppert, S. Turbide, C. Gale, C. Nonaka and S. A. Bass, arXiv:0705.2575 [hep-ph].