

# Forward-Backward (F-B) rapidity correlations in a two step scenario

**J. Dias de Deus and J.G. Milhano**

Instituto Superior Técnico/CENTRA, Av. Rovisco Pais, P-1049-001 Lisboa, Portugal

Two-step scenario models for particle production are based on:

- (i) creation of extended objects in rapidity (glasma longitudinal colour fields or coloured strings); followed by
- (ii) local emission of particles.

The first step guarantees the presence of F-B correlations due to fluctuations in the colour/number of sources, while the second step accounts for local effects such as resonances.

The F-B correlation parameter  $b$  is defined via

$$\langle n_B \rangle_F = a + b n_F, \quad b \equiv D_{FB}^2 / D_{FF}^2, \quad (1)$$

where  $D^2$  is the variance. In general, correlations are measured in two rapidity windows separated by a rapidity gap so that F-B short range correlations are eliminated. In the two-step scenario models we write [1, 2, 3, 4],

$$D_{FB}^2 \equiv \langle n_F n_B \rangle - \langle n_F \rangle \langle n_B \rangle = \frac{\langle n_F \rangle \langle n_B \rangle}{K}, \quad (2)$$

$$D_{FF}^2 \equiv \langle n_F^2 \rangle - \langle n_F \rangle^2 = \frac{\langle n_F \rangle^2}{K} + \langle n_F \rangle, \quad (3)$$

where  $1/K$  is the normalized — e.g., in the number of elementary collisions — long range fluctuation and depends on centrality, energy and rapidity length of the windows. We have assumed, for simplicity, that local emission is of Poisson type.

From (1, 2, 3) we obtain

$$b = \frac{\langle n_B \rangle / \langle n_F \rangle}{1 + K / \langle n_F \rangle}. \quad (4)$$

It should be noticed that  $b$  may be larger than 1, and that a Colour Glass Condensate (CGC) model calculation [5] shows a structure similar to (4):  $b = A[1 + B]^{-1}$  (for a discussion on general properties of (4) and on the CGC model, see [4]).


A simple way of testing (4) is by fixing the backward rapidity window, or  $\langle n_B \rangle$ , in the region of high particle density and move the forward window along the rapidity axis. We can rewrite (4) in the form

$$b = \frac{x}{1 + K'x}, \quad (5)$$

where  $K' \equiv K/\langle n_B \rangle$  is a constant and  $x \equiv \langle n_B \rangle / \langle n_F \rangle$ . In (5), one has  $1 < x < \infty$  with the limiting behaviour:

$$x \rightarrow 1, \quad b \rightarrow \frac{1}{1+K'}; \quad x \rightarrow \infty, \quad b \rightarrow \frac{1}{K'}. \quad (6)$$

The behaviour of (5) is shown in Figure 1 (drawn for  $K' = 1$ ).



**Figure 1.** F-B correlation parameter  $b$  (5) with  $K' = 1$ .

A similar curve is obtained for B-F correlations in the backward region of rapidity. Note that in  $aA$  collisions,  $a \leq A$ , the centrality and energy dependence of  $K'$  is given by [3, 6],

$$K' \sim a^{1/2} A^{-1/6} e^{\lambda Y}, \quad (7)$$

where  $Y$  is the beam rapidity and  $\lambda$  a positive parameter. In the symmetric situation,  $a = A$  and  $K'$  **increases** with centrality (and the curve of the figure moves down) while in the asymmetric situation,  $a = 1, 2 \ll A$  and  $K'$  **decreases** with centrality (and the curve in the figure moves up). As the energy increases  $K'$  increases (and the curve moves down).

J. G. M. acknowledges the financial support of the Fundação para a Ciência e a Tecnologia of Portugal (contract SFRH/BPD/12112/2003).

## References

- [1] J. Dias de Deus, C. Pajares and C. A. Salgado, Phys. Lett. B **407** (1997) 335 [arXiv:hep-ph/9702398].
- [2] M. A. Braun, C. Pajares and V. V. Vechernin, Phys. Lett. B **493**, 54 (2000) [arXiv:hep-ph/0007241].
- [3] P. Brogueira and J. Dias de Deus, arXiv:hep-ph/0611329.
- [4] J. Dias de Deus and J. G. Milhano, *in preparation*.
- [5] N. Armesto, M. A. Braun and C. Pajares, Phys. Rev. C **75**, 054902 (2007) [arXiv:hep-ph/0702216].
- [6] J. Dias de Deus and J. G. Milhano, arXiv:hep-ph/0701215.