

R_{pA} ratio: total shadowing due to running coupling

E. Iancu¹ and D.N. Triantafyllopoulos²

¹ Service de Physique Théorique de Saclay, F-91191 Gif-sur-Yvette, France

² ECT*, Villa Tambosi, Strada delle Tabarelle 286, I-38050 Villazzano (TN), Italy

Abstract. We predict that the R_{pA} ratio at the most forward rapidities to be measured at LHC should be strongly suppressed, close to “total shadowing” ($R_{pA} \simeq A^{-1/3}$), as a consequence of running coupling effects in the nonlinear QCD evolution.

We present predictions for the nuclear modification factor, or “ R_{pA} ratio”, at forward pseudorapidities ($\eta > 0$) and relatively large transverse momenta (p_\perp) for the produced particles, in the kinematical range to be accessible at LHC. These predictions are based on a previous, systematic, study of the R_{pA} ratio within the Color Glass Condensate formalism with running coupling [1]. The ratio can be approximated by

$$R_{pA} \simeq \frac{1}{A^{1/3}} \frac{\Phi_A(Y, p_\perp)}{\Phi_p(Y, p_\perp)}, \quad (1)$$

where $Y = \eta + \ln \sqrt{s}/p_\perp$ and $\Phi(Y, p_\perp)$ is the unintegrated gluon distribution of the corresponding target hadron at fixed impact parameter. When the energy increases one expects more and more momentum modes of this distribution to saturate to a value of order $1/\alpha_s$, and the corresponding saturation momentum reads

$$Q_s^2(Y) = \Lambda^2 \exp \sqrt{B(Y - Y_0) + \ln^2 \frac{Q_s^2(Y_0)}{\Lambda^2}}, \quad (2)$$

with $\Lambda = 0.2\text{GeV}$, $B = 2.25$ and $Y_0 = 4$. The initial condition for the nucleus and the proton are taken as $Q_s^2(A, Y_0) = 1.5\text{GeV}^2$ and $Q_s^2(p, Y_0) = 0.25\text{GeV}^2$ respectively, so that $Q_s^2(A, Y_0) = A^{1/3} Q_s^2(p, Y_0)$ for $A = 208$. The functional form of this expression is motivated by the solution to the nonlinear QCD evolution equations with running coupling [2, 3], while the actual values of the numbers B and Y_0 have been chosen in such a way to agree with the HERA/RHIC phenomenology. As shown in Fig. 1, with increasing Y the two saturation momenta approach to each other and clearly for sufficiently large Y , a nucleus will not be more dense than a proton [3].

For momenta p_\perp larger than Q_s , the gluon distribution satisfies geometrical scaling [2, 4], i.e. it is a function of only the combined variable $p_\perp/Q_s(Y)$:

$$\Phi(p_\perp, Y) \propto \left[\frac{Q_s^2(Y)}{p_\perp^2} \right]^\gamma \left(\ln \frac{p_\perp^2}{Q_s^2(Y)} + c \right), \quad (3)$$

with $\gamma = 0.63$ and $c = \mathcal{O}(1)$. This holds within the scaling window $Q_s \lesssim p_\perp \lesssim Q_g$, where $\ln Q_g^2(Y)/Q_s^2(Y) \sim [\ln Q_s^2(Y)/\Lambda^2]^{1/3}$ and for large Y this is proportional to $Y^{1/6}$. The geometrical scaling lines for a proton and a nucleus are shown in Fig. 1. Note




Figure 1. Left: The ratio of the saturation momenta. ($Y = 12$ corresponds to a pseudorapidity $\eta = 6$ for the produced particles). Right: Geometric scaling windows.

that, since Q_g is increasing much faster than Q_s , a *common scaling window* exists, at $Q_s(A, Y) \lesssim p_\perp \lesssim Q_g(p, Y)$ (and for sufficiently large Y), where the gluon distributions for both the nucleus and the proton are described by Eq. (3).

Within this window, it is straightforward to calculate the R_{pA} ratio. This is shown in Fig. 2 for two values of pseudorapidity. The upper, dotted, line is the asymptotic prediction of a fixed-coupling scenario, in which the ratio $Q_s^2(A, Y)/Q_s^2(p, Y) = \text{const.} = A^{1/3}$, while the lowest, straight, curve is the line of total shadowing $R_{pA} = 1/A^{1/3}$. Our prediction with running coupling is the line in between and it is very close to total shadowing. This is clearly a consequence of the fact that the proton and the nuclear saturation momenta approach each other with increasing energy.




Figure 2. The ratio R_{pA} as a function of p_\perp^2 at $\sqrt{s} = 8.8$ TeV.

Note finally that in the present analysis we have neglected the effects of particle number fluctuations (or “Pomeron loops”). This is appropriate since Pomeron loops effects are suppressed by the running of the coupling [5], and thus can be indeed ignored at all energies of phenomenological interest (in particular, at LHC).

- [1] E. Iancu, K. Itakura, and D.N. Triantafyllopoulos, Nucl. Phys. A **742** (2004) 182.
- [2] A.H. Mueller and D.N. Triantafyllopoulos, Nucl. Phys. B **640** (2002) 331.
- [3] A.H. Mueller, Nucl. Phys. A **724** (2003) 223.
- [4] E. Iancu, K. Itakura, and L. McLerran, Nucl. Phys. A **708** (2002) 327.
- [5] A. Dumitru, E. Iancu, L. Portugal, G. Soyez, D.N. Triantafyllopoulos, arXiv:0706.2540.