

Transverse momentum broadening of vector bosons in heavy ion collisions at the LHC

Zhong-Bo Kang¹ and Jian-Wei Qiu^{1,2}

¹Department of Physics and Astronomy, Iowa State University, Ames IA 50011, USA

²Physics Department, Brookhaven National Laboratory, Upton, NY 11793, USA

E-mail: kangzb@iastate.edu and jwq@iastate.edu

Abstract. We calculate in perturbative QCD the transverse momentum broadening of vector bosons in heavy ion collisions at the Large Hadron Collider (LHC). We predict transverse momentum broadening of W/Z bosons constructed from their leptonic decay channels, which should be a clean probe of initial-state medium effect. We also predict the upper limit of transverse momentum broadening of J/ψ and Υ production as a function of N_{part} at the LHC energy.

Nuclear transverse momentum broadening of heavy vector bosons (γ^* , W/Z , and heavy quarkonia) is defined as a difference between the averaged transverse momentum square measured in nuclear collisions and that measured in collisions of free nucleons,

$$\Delta\langle q_T^2 \rangle_{AB} \equiv \langle q_T^2 \rangle_{AB} - \langle q_T^2 \rangle_{NN} \approx \int dq_T^2 q_T^2 \frac{d\sigma_{AB}^{(D)}}{dq_T^2} \bigg/ \int dq_T^2 \frac{d\sigma_{NN}}{dq_T^2} . \quad (1)$$

Since single scattering is localized in space, the broadening is a result of multiple parton scattering, and is a good probe for nuclear medium properties. Because the mass scale of the vector bosons is much larger than the characteristic momentum scale of the hot medium, the broadening is likely dominated by double partonic scattering as indicated in Eq. (1). The broadening caused by the double scattering can be systematically calculated in terms of high twist formalism in QCD factorization [1, 2].

At the LHC energies, a lot W and Z , and J/ψ and Υ will be produced. Most reconstructed W/Z bosons will come from their leptonic decays. Their transverse momentum broadening is a result of purely initial-state multiple scattering. By calculating the double scattering effect, we obtain [2, 3]

$$\Delta\langle q_T^2 \rangle_{pA}^W = \frac{4\pi^2\alpha_s(M_W)}{3} \lambda_W^2 A^{1/3} , \quad \Delta\langle q_T^2 \rangle_{pA}^Z = \frac{4\pi^2\alpha_s(M_Z)}{3} \lambda_Z^2 A^{1/3} \quad (2)$$

for hadron-nucleus collisions. The $\lambda^2 A^{1/3}$ in Eq. (2) was introduced in [1] as a ratio of nuclear four parton correlation function over normal parton distribution. The λ is proportional to the virtuality or transverse momentum of soft gluons participating in the coherent double scattering. For collisions with a large momentum transfer, Q , the λ^2 should be proportional to $\ln(Q^2)$ [3] and the saturation scale Q_s^2 if the active parton x is small. By fitting Fermilab E772 Drell-Yan data, it was found that $\lambda_{\text{DY}}^2 \approx 0.01 \text{GeV}^2$

at $\sqrt{s} = 38.8\text{GeV}$ [2]. From the λ_{DY}^2 , we estimate the value of λ^2 for production of a vector boson of mass M_V at the LHC energy as

$$\lambda_V^2(\text{LHC}) \approx \lambda_{\text{DY}}^2 \frac{\ln(M_V^2)}{\ln(Q_{\text{DY}}^2)} \left(\frac{M_V/5500}{Q_{\text{DY}}/38.8} \right)^{-0.3}, \quad (3)$$

where we used $Q_s^2 \propto 1/x^\delta$ with $\delta \approx 0.3$ [4] and $\sqrt{s} = 5500$ GeV for the LHC heavy ion collisions. For an averaged $Q_{\text{DY}} \sim 6$ GeV, we obtain $\lambda_{W/Z}^2 \approx 0.05$ at the LHC energy. We can also apply our formula in Eq. (2) to the broadening in nucleus-nucleus collisions by replacing $A^{1/3}$ by an effective medium length L_{eff} . We calculate L_{eff} in Glauber model with inelastic nucleon-nucleon cross section $\sigma_{NN}^{\text{in}} = 70\text{mb}$ at the LHC energy. We plot our predictions (lower set curves) for the broadening of W/Z bosons in Fig.1.




Figure 1. Predicted broadening (maximum broadening) for W and Z (J/ψ and Y) production in p-A (left) and Pb-Pb (right) collisions at $\sqrt{s} = 5500$ GeV.

Heavy quark pairs are produced at a distance scale much less than the physical size of heavy quarkonia in high energy collisions. The pairs produced in heavy ion collisions can have final-state interactions before bound quarkonia could be formed. We found [3] that with both initial- and final-state double scattering, the broadening of heavy quarkonia is close to $2C_A/C_F$ times the Drell-Yan broadening in proton-nucleus collision, which is consistent with existing data [5]. If all soft gluons of heavy ion beams are stopped to form the hot dense medium in nucleus-nucleus collisions, final-state interaction between the almost stationary medium and the fast moving heavy quarks (or quarkonia) of transverse momentum q_T is unlikely to broaden the q_T spectrum, instead, it is likely to slow down the heavy quarks (or quarkonia) [3]. From Eq. (3). we obtain $\lambda_{J/\psi}^2 \approx 0.035$, and $\lambda_Y^2 \approx 0.049$ at the LHC energy; and we predict the maximum broadening for J/ψ and Y production (upper set curves) in Fig. 1.

This work is supported in part by the US Department of Energy under Grant No. DE-FG02-87ER40371 and contract number DE-AC02-98CH10886.

- [1] M. Luo, J. W. Qiu and G. Sterman, Phys. Rev. D **49**, 4493 (1994); D **50**, 1951 (1994).
- [2] X. f. Guo, Phys. Rev. D **58**, 114033 (1998) [arXiv:hep-ph/9804234].
- [3] Z. B. Kang and J. W. Qiu, in preparation.
- [4] K. Golec-Biernat and M. Wusthoff, Phys. Rev. D **59**, 014017 (1999).
- [5] J. C. Peng, AIP Conf. Proc. **494**, 503 (1999) [arXiv:hep-ph/9912371].