

Data-driven goodness-of-fit tests

Mikhail Langovoy*

*Machine Learning and Optimization Laboratory,
EPFL IC, INJ 339, Station 14,
CH-1015 Lausanne, Switzerland
e-mail: mikhail.langovoy@epfl.ch*

Abstract: We propose and study a general method for construction of consistent statistical tests on the basis of possibly indirect, corrupted, or partially available observations. The class of tests devised in the paper contains Neyman's smooth tests, data-driven score tests, and some types of multi-sample tests as basic examples. Our tests are data-driven and are additionally incorporated with model selection rules. The method allows to use a wide class of model selection rules that are based on the penalization idea. In particular, many of the optimal penalties, derived in statistical literature, can be used in our tests. We establish the behavior of model selection rules and data-driven tests under both the null hypothesis and the alternative hypothesis, derive an explicit detectability rule for alternative hypotheses, and prove a master consistency theorem for the tests from the class. The paper shows that the tests are applicable to a wide range of problems, including hypothesis testing in statistical inverse problems, multi-sample problems, and nonparametric hypothesis testing.

MSC subject classification: Primary 62G10; secondary 60F10, 62H10, 62E20.

Keywords and phrases: Data-driven test, model selection, large deviations, asymptotic consistency, generalized likelihood, random quadratic form, statistical inverse problems, stochastic processes.

1. Introduction

Constructing good tests for statistical hypotheses is an essential problem of statistics. There are two main approaches to constructing test statistics. In the first approach, roughly speaking, some measure of distance between the theoretical and the corresponding empirical distributions is proposed as the test statistic. Classical examples of this approach are the Cramer-von Mises and the Kolmogorov-Smirnov statistics (Kolmogorov [1933]). More generally, L^p -distance based tests, as well as graphical tests based on confidence bands, usually belong to this type, and so do kernel tests (see Dette [2002]) and RKHS-based kernel tests that recently became especially popular in machine learning (see, e.g., Sejdinovic et al. [2013], Zaremba et al. [2013], Chwialkowski et al. [2016]). Although these tests are capable of giving very good results, each of these tests is asymptotically optimal only in a finite number of directions of alternatives to a null hypothesis (see Nikitin [1995]).

Nowadays, there is an increasing interest to the second approach of constructing test statistics. The idea of this approach is to construct tests in such a way that the tests would be asymptotically optimal in some sense, or most powerful, at least in a reach enough set of directions. Test statistics constructed following

*Financial support of the Deutsche Forschungsgemeinschaft GK 1023 "Identifikation in Mathematischen Modellen" is gratefully acknowledged.

this approach are often called *score test* statistics. The pioneer of this approach was Neyman [1937]. See also, for example, Wilks [1938], Le Cam [1956], Neyman [1959], Bickel and Ritov [1992], Inglot and Ledwina [1996] for subsequent developments and improvements, and Fan et al. [2001a], Bickel et al. [2006] and Li and Liang [2008] for recent results in the field. This approach is also closely related to the theory of efficient (adaptive) estimation - Bickel et al. [1993], Ibragimov and Has'minskiĭ [1981]. Additionally, it was shown, at least in some basic situations, that data-driven score tests are asymptotically optimal in the sense of intermediate efficiency in an infinite number of directions of alternatives (see Inglot and Ledwina [1996]) and show good overall performance in practice (see Kallenberg and Ledwina [1995]).

Classical score tests have been extensively developed in recent literature: see, for example, the generalized likelihood ratio statistics for nonparametric models in Fan et al. [2001a], tailor-made tests in Bickel et al. [2006] and the semiparametric generalized likelihood ratio statistics in Li and Liang [2008]. See Bickel et al. [2006] for a recent general overview of this and other existing theories of statistical testing, and a discussion of some advantages and disadvantages of different classes of testing methods.

An important line of development in the area of optimal testing concerns with minimax testing, see Ingster [1993], and adaptive minimax testing, see Spokoiny [1996]. Those tests are optimal in a certain minimax sense against wide classes of nonparametric alternatives. In Fan et al. [2001a] it was shown that, for many types of statistical problems, generalized likelihood ratio statistics achieve optimal minimax and adaptive minimax rates. Within the method of the present paper, one can construct and use different variations of data-driven generalized likelihood ratio statistics, too. It is expected that these new tests would share the same optimality properties as the optimal tests from Fan et al. [2001a].

Historically, most of the systematic developments in statistical hypothesis testing were concerned with tests based on directly available observations, rather than with tests for those cases where the variables of interest are only available indirectly, and where the test has to be based on noisy, corrupted, or auxiliary third-party variables. Examples of this type of problems arise in many modern applications, such as testing hypothesis about noisy signals in statistical inverse problems Langovoy [2008], Holzmänn et al. [2007], Bissantz et al. [2007], or hypothesis testing for stochastic processes. However, most of the existing solutions were proposed either on a case-by-case basis, or for relatively specific types of problems. This paper aims to develop a methodological approach that is applicable for both direct and indirect types of hypothesis testing problems.

1.1. Main contributions

In this paper we propose and study a general method for construction of consistent statistical tests on the basis of possibly indirect, corrupted, or partially available observations.

We introduce a class of *data-driven* statistical *NT-tests*. These tests are concerned with testing hypotheses about variables of interest X_1, X_2, \dots, X_m on the basis of auxiliary third-party variables Y_1, Y_2, \dots, Y_n , and their corresponding test statistics are, in general, random quadratic forms of the type

$$T_S = \left\{ \frac{1}{\sqrt{n}} \sum_{j=1}^n l(Y_j) \right\} L \left\{ \frac{1}{\sqrt{n}} \sum_{j=1}^n l(Y_j) \right\}^T,$$

where $l = (l_1, \dots, l_S)$ is an appropriate vector-function with a random dimension chosen via a penalization-based selection rule S , and L serves as a normalization matrix. The choice of l , S and L is problem-dependent and will be treated in this paper.

Neyman's smooth tests, data-driven score tests, some types of multi-sample tests, and efficient deconvolution density tests serve as examples for this class of tests.

The main goal of this paper is to derive a unified approach for construction of consistent NT-tests on the basis of possibly indirect or noisy observations, and in proposing a master theorem establishing consistency of NT-tests. We establish the behavior of model selection rules and data-driven tests under both the null and the alternative hypothesis, derive an explicit detectability rule for alternative hypotheses, and prove a master consistency theorem for data-driven NT-tests.

The paper shows that the tests are applicable to a wide range of problems, including hypothesis testing in statistical inverse problems, multi-sample problems, dependent data problems, and nonparametric hypothesis testing. Semi- and nonparametric generalized likelihood ratio statistics from Fan et al. [2001a] and Li and Liang [2008], score processes from Bickel et al. [2006], and empirical likelihood from Owen [1988], could be used to build consistent data-driven NT-tests.

NT-tests are data-driven and are incorporated with model selection rules. Our method allows to use a wide class of model selection rules that are based on the penalization idea. In particular, many of the optimal penalties, derived in statistical literature, can be used in our tests. The method of this paper allows for big freedom in choice of penalties and dimension growth rates in data-driven model selection, and for flexibility in choosing models with convenient and realistic regularity assumptions. This allows to widen applicability of already existing data-driven tests.

It becomes possible to construct and properly tune data-driven tests for new statistical problems by combining recent developments in model selection, efficient testing, generalized likelihood, and other areas. We describe some of the possibilities in this paper. A statistician can take into account specific features of his particular problem and choose a test with desired properties among a family of consistent data-driven tests.

In many modern problems, e.g. in semiparametric hypothesis testing, the model is so complicated that the calculations necessary for a proof of consistency are hardly possible to carry out. Consistency theorems of this paper allow to bypass this step, and deliver consistency results in problems where direct computations are too cumbersome. In certain cases, our consistency theorems lead to generalization or improvement of already existing results (see, for example, Proposition 2 below).

For any data-driven NT-test we give an explicit rule to determine, for every particular alternative, whether the test will be consistent against this alterna-

tive. This rule allows us to describe, in a closed form, the set of detectable alternatives for every NT-test.

1.2. Paper outline

In Section 2, we describe the framework and introduce a class of NT-statistics, for the case of deterministic model dimension. This is one of the main building blocks for our tests. In Section 3, we define penalization-based model selection rules that will be used in our tests, and introduce a concept of data-driven tests. In Section 4, we establish behavior of model selection rules and data-driven NT-tests for the case when the alternative hypothesis is true. In this Section, we also derive an explicit consistency condition that allows to check whether any particular alternative could be detected by the data-driven NT-test. In Section 5 we describe what happens with a data-driven NT-test under the null hypothesis. Finally, in Section 6 a master consistency theorem for data-driven NT-tests is given. Additionally, each section contains examples illustrating the applicability of new concepts by connecting them to a variety of well-established special cases.

2. NT-statistics

2.1. Basic definition: NT-statistics with deterministic dimensions

Let X_1, X_2, \dots be a sequence of random variables with values in an arbitrary measurable space \mathbb{X} . Suppose that for every m the random variables X_1, \dots, X_m have the joint distribution P_m from the family of distributions \mathbb{P}_m . Suppose there is a given functional \mathcal{F} acting from the direct product of the families $\otimes_{m=1}^{\infty} \mathbb{P}_m = (\mathbb{P}_1, \mathbb{P}_2, \dots)$ to a known set Θ , and that $\mathcal{F}(P_1, P_2, \dots) = \theta$. We consider the following generic

Problem: test the hypothesis $H_0 : \theta \in \Theta_0 \subset \Theta$ against the alternative $H_A : \theta \in \Theta_1 = \Theta \setminus \Theta_0$, on the basis of *indirect observations* Y_1, \dots, Y_n having their values in an arbitrary measurable space \mathbb{Y} (i.e. *not necessarily on the basis of variables of interest* X_1, \dots, X_m).

Here Θ can be any set, for example, a functional space; correspondingly, parameter θ can be infinite dimensional. It is not assumed that Y_1, \dots, Y_n are independent or identically distributed. The measurable space \mathbb{Y} can be infinite dimensional. This allows to apply the results of this paper in statistics for stochastic processes. Additional assumptions on Y_i 's will be imposed below, when it would be necessary.

An important feature of our approach is that we are able to consider the case when the null hypothesis H_0 is not about observable Y_i 's, but about some other random variables X_1, \dots, X_m . This makes it possible to use our method in the case of statistical inverse problems (see Langovoy [2008], and Examples 2 and 3 below). Under conditions of Theorem 3 of the present paper, it would be still possible to extract from Y_i 's enough information in order to build a consistent test about X_i 's.

Now we introduce one of the main concepts of this paper.

Definition 1. (NT-statistics with deterministic dimensions). Suppose we have n random observations Y_1, \dots, Y_n with values in a measurable space \mathbb{Y} . Let k be a fixed number and $l = (l_1, \dots, l_k)$ be a vector-function, where $l_i : \mathbb{Y} \rightarrow \mathbb{R}$ for $i = 1, \dots, k$ are some known Lebesgue measurable functions. Set

$$L = \{E_0[l(Y)]^T l(Y)\}^{-1}, \quad (1)$$

where the mathematical expectation E_0 is taken with respect to P_0 , and P_0 is the (fixed and known in advance) distribution function of some auxiliary random variable Y , where Y is assuming its values in the space \mathbb{Y} . Assume that $E_0 l(Y) = 0$ and L is well defined in the sense that all its elements are finite. Put

$$T_k = \left\{ \frac{1}{\sqrt{n}} \sum_{j=1}^n l(Y_j) \right\}^T L \left\{ \frac{1}{\sqrt{n}} \sum_{j=1}^n l(Y_j) \right\}. \quad (2)$$

We call T_k a *statistic of Neyman's type* (or NT-statistic).

Here l_1, \dots, l_k can be some score functions, as was the case for the classical Neyman's test Neyman [1937], but it is possible to use any other functions, depending on the problem under consideration. We prove below that under additional assumptions it is possible to construct consistent tests of such form without using scores in (2). We will discuss different possible sets of meaningful additional assumptions on l_1, \dots, l_k in Sections 4 - 6.

Scores are based on the notion of maximum likelihood. In our constructions it is possible to use, for example, truncated, penalized or generalized likelihood to build a test. It is even possible to use functions l_1, \dots, l_k such that they are unrelated to any kind of likelihood.

If, for example, Y_i 's are equally distributed, then the natural choice for P_0 is their distribution function under the null hypothesis. Thus, L will be the inverse to the covariance matrix of the vector $l(Y)$. Such construction is often used in score tests for simple hypothesis. But our definition allows to use a reasonable substitution instead of the covariance matrix. This is helpful for testing in certain semi- or nonparametric models.

2.2. Examples of NT-tests

Example 1. Neyman's smooth tests. The basic example of an NT-statistic is the Neyman's smooth test statistic for simple hypotheses (see Neyman [1937]). Let X_1, \dots, X_n be i.i.d. random variables. Consider the problem of testing the simple null hypothesis H_0 that the X_i 's have the uniform distribution on $[0, 1]$. Let $\{\phi_j\}$ denote the family of orthonormal Legendre polynomials on $[0, 1]$. Then for every k one has the test statistic

$$T_k = \sum_{j=1}^k \left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n \phi_j(X_i) \right\}^2.$$

We see that Neyman's classical test statistic is an NT-statistic, with $Y_i = X_i$ for all i in this very simplest example.

Hypothesis testing in statistical inverse problems is an important but still a rather unexplored area in modern statistics. Applications to statistical inverse problems served as the main motivation for studying NT-tests.

Example 2. Statistical inverse problems. The most well-known example here is the deconvolution problem. This problem appears when one has noisy signals or measurements: in physics, seismology, optics and imaging, engineering. It is a building block for many complicated statistical inverse problems. The book Carroll et al. [2006] provides many examples related to deconvolution problems. In Langovoy [2008], data-driven score tests for the problem were constructed, thus solving the deconvolution density testing problem. The solution of Langovoy [2008] is a special case of the results of the present paper.

The basic deconvolution density testing problem is formulated as follows. Suppose that instead of X_i one observes Y_i , where

$$Y_i = X_i + \varepsilon_i,$$

and ε_i 's are i.i.d. with a known density h with respect to the Lebesgue measure λ ; also X_i and ε_i are independent for each i and $E\varepsilon_i = 0$, $0 < E\varepsilon_i^2 < \infty$. Assume that X has a density with respect to λ . Our null hypothesis H_0 is the simple hypothesis that X has a known density f_0 with respect to λ . Let us choose for every $k \leq d(n)$ an auxiliary parametric family $\{f_\theta\}$, $\theta \in \Theta \subseteq \mathbb{R}^k$ such that f_0 from this family coincides with f_0 from the null hypothesis H_0 . The true F possibly has no relation to the chosen $\{f_\theta\}$. Set

$$l(y) = \frac{\frac{\partial}{\partial \theta} \left(\int_{\mathbb{R}} f_\theta(s) h(y-s) ds \right) \Big|_{\theta=0}}{\int_{\mathbb{R}} f_0(s) h(y-s) ds} \quad (3)$$

and define the corresponding test statistic U_k by the formula (2). Under appropriate regularity assumptions, U_k is an NT-statistic (see Langovoy [2007]). As was shown in Langovoy [2008], the data-driven test U_S , based on U_k , is asymptotically consistent against a wide class of nonparametric alternatives.

Subsequently, Butucea et al. [2009] proposed an adaptive goodness-of-fit test for this model with indirect observations, which appears to be the first nonparametric adaptive test in statistical inverse problems. The test from Butucea et al. [2009] is restricted to polynomially smooth error density and to signal densities coming from even narrower smoothness classes. The data-driven test U_S from Langovoy [2007] was intended to cover wider scope of applications and to have a flexible penalization procedure, so optimality properties were not attainable over the intended consistency range. Adaptive modifications of U_S can be constructed via using special penalties, tailored to deliver minimax optimality for convolution densities from properly restricted classes (analogously to the method of Fan et al. [2001b]).

The following example serves as an illustration of applicability of NT-tests to multisample problems.

Example 3. Rank Tests for Independence. Let $X_1 = (V_1, W_1), \dots, X_n = (V_n, W_n)$ be i.i.d. random variables with the distribution function D and the marginal distribution functions F and G for V_1 and W_1 . Assume that F and G are continuous, but unknown. It is the aim to test the null hypothesis of independence

$$H_0 : D(v, w) = F(v)G(w), \quad v, w \in \mathbb{R}, \quad (4)$$

against a wide class of alternatives. The following construction was proposed in Kallenberg and Ledwina [1999].

Let ϕ_j denote the j -th orthonormal Legendre polynomial (i.e., $\phi_1(x) = \sqrt{3}(2x-1)$, $\phi_2(x) = \sqrt{5}(6x^2-6x+1)$, etc.). The score test statistic from Kallenberg and Ledwina [1999] is

$$T_k = \sum_{j=1}^k \left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n \phi_j \left(\frac{R_i - 1/2}{n} \right) \phi_j \left(\frac{S_i - 1/2}{n} \right) \right\}^2, \quad (5)$$

where R_i stands for the rank of V_i among V_1, \dots, V_n and S_i for the rank of W_i among W_1, \dots, W_n . Thus defined T_k satisfies Definition 1 of NT-statistics: put

$$Y_i = (Y_i^{(1)}, Y_i^{(2)}) := \left(\frac{R_i - 1/2}{n}, \frac{S_i - 1/2}{n} \right)$$

and $l_j(Y_i) := \phi_j(Y_i^{(1)}) \phi_j(Y_i^{(2)})$. Under the null hypothesis $L_k = E_{k \times k}$, and $E_0 l(Z) = 0$. Thus, T_k is an NT-statistic. New Y_i depends on the original X_i 's ($= (V_i, W_i)$'s) in a nontrivial way, but still contains some information about the pair of interest. \square

Many other examples, including data-driven tests based on partial likelihood (see Cox [1975]), can be found in Langovoy [2007] and Langovoy [2009].

3. Data-driven tests and model selection

3.1. Selection rules and data-driven tests: definitions

It was shown that for achieving optimality of efficient score tests it is important to select the right number of components in the test statistic (see Bickel and Ritov [1992], Eubank et al. [1993], Fan [1996], Kallenberg [2002]). Therefore, we provide a corresponding refinement for NT-statistics as well. Using the idea of a penalized likelihood, we propose a general mathematical framework for constructing rules to find reasonable model dimensions. We make our tests *data-driven*, i.e., the tests choose a reasonable number of components in the test statistics automatically and accordingly to the data. Our construction offers substantial freedom in the choice of penalties and possible basis models. This way, it is easy to take into account specific features of particular problems and to choose suitable statistical models and reasonable penalties in order to build tests with desired properties for each particular problem.

We will not restrict a possible number of components in test statistics by some fixed number, but instead we allow the number of components to grow

unlimitedly as the number of observations grows. This is important because the more observations Y_1, \dots, Y_n we have, the more information is available about the problem. This makes it possible to give a more detailed description of the phenomena under investigation. In our case this means that the complexity of the model and the possible number of components in the corresponding test statistic grow with n at a controlled rate.

Denote by M_k a statistical model designed for a specific statistical problem satisfying assumptions of Section 2. Assume that the true parameter value θ belongs to the parameter set of M_k , call it Θ_k . We say that the family of models M_k for $k = 1, 2, \dots$ is nested if for their parameter sets it holds that $\Theta_1 \subseteq \Theta_2 \subseteq \dots$. Here Θ'_k 's can be infinite-dimensional. We also do not require that all Θ'_k 's are different (analogously to Birgé and Massart [2001], p. 221). Let T_k be an *arbitrary* statistic for testing validity of the model M_k on the basis of observations Y_1, \dots, Y_n . The following definition applies for the sequence of statistics $\{T_k\}$.

Definition 2. (Penalty, selection rule, and data-driven tests) Consider a nested family of models M_k for $k = 1, \dots, d(n)$, where $d(n)$ is a control sequence, giving the largest possible model dimension for the case of n observations. Choose a function $\pi(\cdot, \cdot) : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{R}$, where \mathbb{N} is the set of natural numbers. Assume that $\pi(1, n) < \pi(2, n) < \dots < \pi(d(n), n)$ for all n and $\pi(j, n) - \pi(1, n) \rightarrow \infty$ as $n \rightarrow \infty$ for every $j = 2, \dots, d(n)$. Call $\pi(j, n)$ a *penalty attributed to the j -th model M_j and the sample size n* .

Then a *selection rule* S for the sequence of statistics $\{T_k\}$ is an integer-valued random variable satisfying the condition

$$S = \min\{k : 1 \leq k \leq d(n); T_k - \pi(k, n) \geq T_j - \pi(j, n), j = 1, \dots, d(n)\}. \quad (6)$$

We call T_S a *data-driven test statistic* for testing validity of the initial model.

The definition is expected to be applied, of course, only if the sequence $\{T_k\}$ is increasing unboundedly, in the sense that $T_k(Y_1, \dots, Y_n) \rightarrow \infty$ in probability as $k \rightarrow \infty$.

In statistical literature, one often tries to choose penalties such that they possess some sort of minimax or Bayesian optimality. Classical examples of the penalties constructed via this approach are Schwarz's penalty $\pi(j, n) = j \log n$ (see Schwarz [1978]), and minimum description length penalties, see Rissanen [1983]. For more examples of optimal penalties and recent developments, see Abramovich et al. [2007], Birgé and Massart [2001] or Bunea et al. [2007]. In this paper, we do not aim for optimality of the penalization; our goal is to be able to build consistent data-driven tests based on different choices of penalties. The penalization technique that we use in this paper allows for many possible choices of penalties. In our framework it is possible to use most of the penalties from the abovementioned papers. As an illustration, see Example 4 below.

3.2. Model selection rules: examples

Example 2, continued. Data-driven tests for inverse problems. One can incorporate U_k with the following selection rule widely used in statistical

literature:

$$S = \min\{k : 1 \leq k \leq d(n); T_k - k \log n \geq T_j - j \log n, j = 1, 2, \dots, d(n)\}. \quad (7)$$

This selection rule satisfies Definition 2, so thus defined U_S extends the test statistic from Langovoy [2008] and makes it a data-driven NT-statistic. \square

Example 4. Gaussian model selection. Birgé and Massart in Birgé and Massart [2001] proposed a method of model selection in a framework of Gaussian linear processes. This framework is quite general and includes as special cases a Gaussian regression with fixed design, Gaussian sequences and the model of Ibragimov and Has'minskii. In this example we briefly describe the construction (for more details see the original paper) and then discuss the relation with our results.

Given a linear subspace \mathbb{S} of some Hilbert space \mathbb{H} , we call Gaussian linear process on \mathbb{S} , with mean $s \in \mathbb{H}$ and variance ε^2 , any process Y indexed by \mathbb{S} of the form

$$Y(t) = \langle s, t \rangle + \varepsilon Z(t),$$

for all $t \in \mathbb{S}$, and where Z denotes a linear isonormal process indexed by \mathbb{S} (i.e. Z is a centered and linear Gaussian process with covariance structure $E[Z(t)Z(u)] = \langle t, u \rangle$). Birgé and Massart considered estimation of s in this model.

Let S be a finite dimensional subspace of \mathbb{S} and set $\gamma(t) = \|t\|^2 - 2Y(t)$. One defines the projection estimator on S to be the minimizer of $\gamma(t)$ with respect to $t \in S$. Given a finite or countable family $\{S_m\}_{m \in \mathcal{M}}$ of finite dimensional linear subspaces of S , the corresponding family of projection estimators \hat{s}_m , built for the same realization of the process Y , and given a nonnegative function pen defined on \mathcal{M} , Birgé and Massart estimated s by a penalized projection estimator $\tilde{s} = \hat{s}_{\hat{m}}$, where \hat{m} is any minimizer with respect to $m \in \mathcal{M}$ of the penalized criterion

$$crit(m) = -\|\hat{s}_m\|^2 + pen(m) = \gamma(\hat{s}_m) + pen(m).$$

They proposed some specific penalties pen such that the penalized projection estimator has the optimal order risk with respect to a wide class of loss functions. The method of model selection of this paper has a relation with the one of Birgé and Massart [2001].

4. Data-driven NT-tests: alternative hypothesis case

In this section we study the behavior of NT-statistics and model selection rules under alternatives.

4.1. Explicit consistency condition

Let P denote the alternative distribution of Y_i 's. Suppose that $E_P l(Y)$ exists. Now we formulate the following crucial explicit *consistency condition*:

Consistency condition

$\langle \mathbf{C} \rangle$ there exists integer $K = K(P) \geq 1$ such that
 $E_P l_1(Y) = 0, \dots, E_P l_{K-1}(Y) = 0, E_P l_K(Y) = C_P \neq 0,$

where l_1, \dots, l_k are as in Definition 1.

Example 1 (continued). Consistent smooth tests. In this basic problem, condition $\langle C \rangle$ is simply equivalent to the following one: there exists $K = K_{\mathbb{P}}$ such that $E_{\mathbb{P}} \phi_K(X) \neq 0$. \square

Example 2 (continued). Consistent density testing in inverse problems. Let F be a true distribution of X . Here F is not supposed to be parametric and possibly does not have a density. Let l be the score vector given by (3) and h be the known error density. As was shown in Langovoy [2008] and Langovoy [2007], if there exists an integer $K = K(F) \geq 1$ such that

$$E_{F*h} l_1(Y) = 0, \dots, E_{F*h} l_{K-1}(Y) = 0, E_{F*h} l_K(Y) \neq 0, \quad (8)$$

then U_S is asymptotically consistent against a wide class of nonparametric alternatives. \square

We see that conditions equivalent to $\langle C \rangle$ naturally appear in studies of consistency of data-driven score tests. This condition explicitly describes which directions of alternatives to the null hypothesis can be captured by an NT-test.

4.2. Model selection under alternatives

We impose now two additional assumptions on the abstract model of Section 2. First, we assume that Y_1, Y_2, \dots are identically distributed. We do *not* assume that Y_1, Y_2, \dots are independent. It is possible that the sequence of interest X_1, X_2, \dots consists of dependent and nonidentically distributed random variables.

It is only important that the new (possibly obtained by a complicated transformation) sequence Y_1, Y_2, \dots obeys the consistency conditions. Then it is possible to build consistent tests of hypotheses about X_i 's. The intuition behind this is that, even after a complicated transformation, the transformed sequence still can contain some part of the information about the sequence of interest. However, if the transformed sequence Y_1, Y_2, \dots is not chosen reasonably, then test can be meaningless: it can be formally consistent, but against an empty or almost empty set of alternatives.

Another assumption we impose is that the first K components of $l(Y_i)$'s satisfy both the law of large numbers and the multivariate central limit theorem, i.e. that for the vectors $l(Y_1) = (l_1(Y_1), \dots, l_K(Y_1)), \dots, l(Y_n) = (l_1(Y_n), \dots, l_K(Y_n))$ it holds that

$$\frac{1}{n} \sum_{j=1}^n l(Y_j) \rightarrow E_P l(Y) \quad \text{in } P\text{-probability as } n \rightarrow \infty,$$

$$n^{-1/2} \sum_{j=1}^n (l(Y_j) - E_P l(Y)) \rightarrow_d \mathcal{N} \quad (9)$$

where \mathcal{N} stands for a K -dimensional normal distribution with zero mean vector.

These assumptions restrict the choice of the function l and leave us with a uniquely determined P_0 . Nonetheless, random variables of interest X_1, \dots, X_n are still allowed to be nonidentically distributed, and their transformed counterparts Y_1, \dots, Y_n are still allowed to be dependent.

Without loss of generality, below we always assume that

$$\lim_{n \rightarrow \infty} d(n) = \infty. \quad (10)$$

The (relatively degenerate) case when $d(n)$ is nondecreasing and bounded from above by some constant D can be handled analogously to the method of this paper, see Langovoy [2007].

To clarify the statement of the next theorem, we remind that in Definition 1 L is a $k \times k$ -matrix. Below we will sometimes denote it L_k in order to stress the model dimension. Accordingly, ordered eigenvalues of L_k will be denoted $\lambda_1^{(k)} \geq \lambda_2^{(k)} \geq \dots \geq \lambda_k^{(k)}$. We have the sequence of matrices $\{L_k\}_{k=1}^\infty$ and each matrix has its own eigenvalues. Whenever possible, we will use the simplified notation from Definition 1.

Proposition 1. *Let $\langle C \rangle$ and (10) holds and*

$$\lim_{n \rightarrow \infty} \sup_{k \leq d(n)} \frac{\pi(k, n)}{n \lambda_k^{(k)}} = 0. \quad (11)$$

Then

$$\lim_{n \rightarrow \infty} P(S \geq K) = 1.$$

Remark 1. Condition (11) means that not only n tends to infinity, but that it is also possible for k to grow infinitely, but at the controlled rate. This condition is needed to ensure that the penalty π is not so large that it trivializes the data-driven model choice. It is easy to check (11) for any specific function π since the eigenvalues of L_k are known in advance, and it is easy to satisfy (11) because we are free to choose π from the whole range of possible penalty functions.

4.3. Data-driven NT-tests under detectable alternatives

Now suppose that the alternative distribution P is such that $\langle C \rangle$ is satisfied and that there exists a sequence $\{r_n\}_{n=1}^\infty$ such that $\lim_{n \rightarrow \infty} r_n = \infty$ and

$$\langle \mathbf{A} \rangle \quad P \left(\frac{1}{n} \left| \sum_{i=1}^n [l_K(Y_i) - E_P l_K(Y_i)] \right| \geq y \right) = O \left(\frac{1}{r_n} \right).$$

Note that in $\langle A \rangle$ we do not require uniformity in y , i.e. r_n gives us the rate, but the exact bound can depend on y . In some sense condition $\langle A \rangle$ is a way to make the weak law of large numbers for $l_K(Y_i)$'s more precise. The following lemma shows that condition $\langle A \rangle$ is often easily satisfied.

Lemma 1. *Let $l_K(Y_i)$'s be bounded i.i.d. random variables with finite expectation and variance σ^2 . Then condition $\langle A \rangle$ is satisfied with $r_n = \exp(ny^2/2\sigma)$.*

Therefore, one can often expect exponential rates in $\langle A \rangle$, but even a much slower rate is not a problem. The proof of this standard lemma is omitted.

The main result of this section is the following theorem that describes asymptotic behavior of NT-tests under alternative hypotheses.

Theorem 1. *Let $\langle A \rangle$, $\langle C \rangle$, (10) and (11) holds and*

$$d(n) = o(r_n) \quad \text{as } n \rightarrow \infty. \quad (12)$$

Then $T_S \rightarrow_P \infty$ as $n \rightarrow \infty$.

Remark 2. Sometimes it is possible to find asymptotic distributions of NT-statistics, both under the null hypothesis and under alternatives. In order to do this, one can approximate the quadratic form T_k by certain quadratic form in N_1, \dots, N_k , where N_i 's are Gaussian random variable with suitably chosen mean and covariance structure. When this approximation is possible, the error of approximation is often of order $n^{-1/2}$ and depends on the smallest eigenvalue of the covariance of $l(Y_1)$. See Götze and Tikhomirov [1999], p. 1078 for more details.

Note that for proving consistency of data-driven NT-tests one doesn't need to know asymptotic distributions of underlying NT-statistics. Theorem 1 is strong enough for our purposes.

5. Data-driven NT-tests under the null hypothesis

Now we study the asymptotic behavior of data-driven NT-statistics under the null hypothesis.

5.1. Proper penalties and majorants for data-driven tests

The following two technical definitions are useful for data-driven model selection in general.

Definition 3. (Penalties of proper weight). Let $\{T_k\}$ be a sequence of NT-statistics and S be a selection rule for it. Suppose that $\lambda_1 \geq \lambda_2 \geq \dots$ are ordered eigenvalues of L , where L is defined by (1). We say that the penalty $\pi(k, n)$ in S is of *proper weight*, if the following conditions holds:

1. there exists sequences of real numbers $\{s(k, n)\}_{k,n=1}^\infty$, $\{t(k, n)\}_{k,n=1}^\infty$, such that

(a)

$$\lim_{n \rightarrow \infty} \sup_{k \leq u_n} \frac{s(k, n)}{n\lambda_k^{(k)}} = 0,$$

where $\{u_n\}_{n=1}^\infty$ is some real sequence such that $\lim_{n \rightarrow \infty} u_n = \infty$.

- (b) $\lim_{n \rightarrow \infty} t(k, n) = \infty$ for every $k \geq 2$,
 $\lim_{k \rightarrow \infty} t(k, n) = \infty$ for every fixed n .
- 2. $s(k, n) \leq \pi(k, n) - \pi(1, n) \leq t(k, n)$ for all k, n .
- 3.

$$\lim_{n \rightarrow \infty} \sup_{k \leq m_n} \frac{\pi(k, n)}{n \lambda_k^{(k)}} = 0,$$

where $\{m_n\}_{n=1}^\infty$ is some real sequence such that $\lim_{n \rightarrow \infty} m_n = \infty$.

Definition 3 is more general than required for our present goals. As can be seen from Theorem 2 below, in our research on consistency of data-driven tests we can assume $u_n = m_n = d(n)$ for all n , where $d(n)$ is the model complexity growth rate from Definition 2. We have chosen to state Definition 3 in its full generality because of its general usefulness in model selection.

Condition 2 of Definition 3 is needed to ensure that the penalty π actually penalizes models with large dimension k . The choice of $s(k, n)$ and $t(k, n)$ will be explained below.

For notational convenience, for $l = (l_1, \dots, l_k)$ from Definition 1 we denote

$$\bar{l}_j := \frac{1}{n} \sum_{i=1}^n l_j(Y_i), \quad \bar{l} := (\bar{l}_1, \bar{l}_2, \dots, \bar{l}_k) \quad (13)$$

and define a quadratic form

$$Q_k(\bar{l}) = (\bar{l}_1, \bar{l}_2, \dots, \bar{l}_k) L(\bar{l}_1, \bar{l}_2, \dots, \bar{l}_k)^T. \quad (14)$$

In the new notation, $T_k = Q_k(\bar{l})$, where T_k is the statistic from Definition 1. Below we also use the notation of Definition 3.

Definition 4. (Proper majorant). Let S be with a penalty of proper weight. Assume that there exists a Lebesgue measurable function $\varphi(\cdot, \cdot) : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, such that φ is monotonically decreasing in the second argument and monotonically nondecreasing in the first one, and assume that

- 1. for every $\varepsilon > 0$ there exists $K = K_\varepsilon$ such that for every $n > n(\varepsilon)$

$$\langle \mathbf{B1} \rangle \quad \sum_{k=K_\varepsilon}^{u_n} \varphi(k; s(k, n)) < \varepsilon,$$

where $\{u_n\}_{n=1}^\infty$ is as in Definition 3.

- 2.

$$\langle \mathbf{B2} \rangle \quad P_0(n Q_k(\bar{l}) \geq y) \leq \varphi(k; y)$$

for all $k \geq 1$ and $y \in [s(k, n); t(k, n)]$, where P_0 is as in Definition 1.

We call φ a *proper majorant* for (large deviations of) the statistic T_S . Equivalently, we say that (large deviations of) the statistic T_S are *properly majorated* by φ .

Again, one is free to assume in this Definition that $u_n = d(n)$ for all n . In Definitions 3 and 4 we need some lower bound $s(k, n)$ to be sure that the penalty π is not "too light", i.e. that the penalty somehow affects the choice of the model dimension and protects us from choosing a "too complicated" model. In nontrivial cases, it follows from $\langle B1 \rangle$ that $s(k, n) \rightarrow \infty$ as $k \rightarrow \infty$. The exact choice of functions s and t for each specific problem is dictated by the variant of inequalities $\langle B1 \rangle$ and $\langle B2 \rangle$ that holds in this particular statistical problem.

Example 1 (continued). Model selection for data-driven smooth tests. Put $\pi(k, n) = k \log n$. Then Prohorov's inequality (see Appendix) suggests the following possible φ in Definition 4:

$$\varphi(k; y) = \frac{150210}{\Gamma(k/2)} \left(\frac{y^2}{2} \right)^{\frac{k-1}{2}} \exp \left\{ -\frac{y^2}{4} \right\}, \quad (15)$$

with, for example, $s(k, n) := 3k$ and $t(k, n) := 1/4\sqrt{n/k}$. All conditions of Definitions 3 and 4 can be easily checked. It follows that one can afford any model dimension growth rate such that $d(n) = o(\{n/\log n\}^{1/3})$. \square

For many test statistics Prohorov's inequality can't be used to estimate the large deviations. This is usually the case for more complicated models where the matrix L is not diagonal. This is typical for statistical inverse problems as well as for problems with dependent data. If one deals with likelihood-based NT-statistics, then the easiest way is to use large deviations results that were established in studies on usual likelihood tests and their efficiencies. Otherwise, one has to find a direct way to estimate the test statistic.

Remark 3. There is a general approach that makes it possible to check conditions of Definition 4, and to study distributions of NT-statistics for moderate sample sizes, in many particular situations. This method consists of two steps. On the first step, one approximates the quadratic form $Q(\tilde{l}(Y))$ by the simpler quadratic form $Q(N)$, where N is the Gaussian random variable with the same mean and covariance structure as $l(Y)$. This approximation is possible, for example, under conditions given in Götze and Tikhomirov [1999] or Bentkus and Götze [1996]. These authors gave the rate of convergence for such approximation. The second step is to establish a large deviation result for the quadratic form $Q(N)$; this form has a more predictable distribution. Even for strongly dependent random variables, one can hope to use results from Horváth and Shao [1999].

5.2. Data-driven NT-statistics under null hypotheses

The following is the key result describing the behaviour of a data-driven NT-statistic under the null hypothesis.

Theorem 2. *Let $\{T_k\}$ be a sequence of NT-statistics and S be a selection rule for the sequence. Assume that the penalty in S is of proper weight and that large deviations of statistics T_S are properly majorated. Suppose that*

$$d(n) \leq \min\{u_n, m_n\}. \quad (16)$$

Then $S = O_{P_0}(1)$ and $T_S = O_{P_0}(1)$.

Remark 4. Usually it happens that if the penalty does not satisfy all the conditions of Definitions 3 and 4, then T_S has the same distribution under both alternative and null hypotheses and the test is inconsistent.

As the first corollary, we have the following generalization of previously known null-hypothesis results for Neyman's smooth test. As was shown in Langovoy [2007] and Langovoy [2009], this corollary allows to build statistical tests for new problems that go far beyond the classic case of testing simple hypothesis of uniformity.

Proposition 2. *Let $\{T_k\}$ be a family of NT-statistics and S a selection rule for the family. Assume that Y_1, \dots, Y_n are i.i.d. Let $El(Y_1) = 0$ and assume that for every k the vector $(l_1(Y_i), \dots, l_k(Y_i))$ has the unit covariance matrix. Suppose that $\|(l_1(Y_1), \dots, l_k(Y_1))\|_k \leq M(k)$ a.e., where $\|\cdot\|_k$ is the norm of the k -dimensional Euclidean space. Assume $\pi(k, n) - \pi(1, n) \geq 2k$ for all $k \geq 2$ and*

$$\lim_{n \rightarrow \infty} \frac{M(d(n))\pi(d(n), n)}{\sqrt{n}} = 0. \quad (17)$$

Then $S = O_{P_0}(1)$ and $T_S = O_{P_0}(1)$.

Example 1 (continued). Data-driven smooth tests under the null hypothesis. As a simple corollary, we derive the following slight improvement of Theorem 3.2 from Kallenberg [2002].

Corollary 3. *Let T_S be the Neyman's smooth data-driven test statistic for the case of simple hypothesis of uniformity. Assume that $\pi(k, n) - \pi(1, n) \geq 2k$ for all $k \geq 2$ and that for all $k \leq d(n)$*

$$\lim_{n \rightarrow \infty} \frac{d(n)\pi(d(n), n)}{\sqrt{n}} = 0.$$

Then $S = O_{P_0}(1)$ and $T_S = O_{P_0}(1)$.

Proof. It is enough to note that in this case $M(k) = \sqrt{(k-1)(k+3)}$ and apply Proposition 2. \square

To prove consistency of a test based on some test statistic, usually it is required to use inequalities for large deviations of this test statistic. NT-statistics are no exception: in order to prove consistency of an NT-test, one first has to establish large deviations inequalities for NT-statistics under consideration. The model regularity assumptions and the model dimension growth rate $d(n)$ will be strongly connected with this inequality.

In certain cases it can be desirable to have $d(n)$ growing at a fast rate, even by the cost of restrictive regularity assumptions that can arise from using a sharp large deviations inequality. Sometimes, it is better to use a simple crude inequality that requires only little additional assumptions, but allows only for a slow growth of $d(n)$. One useful point of Definitions 3 and 4 and Theorem 3 below is that one can be sure in advance that whatever large deviations inequality he is able to prove, he will succeed in proving a consistency theorem, provided that the chosen inequality satisfies conditions $\langle B1 \rangle$ and $\langle B2 \rangle$. Moreover, once an inequality is chosen, the rate $d(n)$ is obtained from Theorem 3.

6. Consistency theorem

Now we formulate the general consistency theorem for data-driven NT-tests. We understand consistency of the test based on T_S in the sense that under the null hypothesis T_S is bounded in probability, while under fixed alternatives $T_S \rightarrow \infty$ in probability.

Theorem 3. (*Consistency of data-driven NT-tests.*) *Let $\{T_k\}$ be a sequence of NT-statistics and S be a selection rule for it. Assume that the penalty in S is of proper weight. Assume that conditions (A), (10) and (11) are satisfied and that $d(n) = o(r_n)$, $d(n) \leq \min\{u_n, m_n\}$. Then the test based on T_S is consistent against any alternative distribution P satisfying condition (C).*

Without a general consistency theorem, one has to perform the whole proof of consistency of a data-driven test anew for every particular problem. This becomes especially difficult in many semi- and nonparametric problems. Using the general consistency Theorem 3, some type of consistency result can be obtained for any data-driven NT-statistics, and types of detectable alternatives can be characterized explicitly.

Example 4 (continued). Data-driven tests for Gaussian linear processes. In the model of Birgé and Massart $\gamma(t)$ is the least squares criterion and \hat{s}_m is the least squares estimator of s , which is in this case the maximum likelihood estimator. Therefore $\|\hat{s}_m\|^2$ is the Neyman score for testing the hypothesis $s = 0$ within this model. Risk-optimizing penalties pen proposed in Birgé and Massart [2001] satisfy the conditions of Definition 2 (after the change of notations $pen(m) = \pi(m, n)$; for the explicit expressions of pen 's see the original paper). Therefore, $\|\hat{s}_{\hat{m}}\|^2$ is, in our terminology, the data-driven NT-statistic. As follows from the consistency Theorem 3, $\|\hat{s}_{\hat{m}}\|^2$ can be used for testing $s = 0$ and has a good range of consistency, even though this particular penalty probably does not lead to adaptively optimal testing, see Baraud et al. [2003]. \square

Many other examples of applications of the consistency theorem can be found in Langovoy [2007] and Langovoy [2009].

Remark 5. In general, it seems to be possible to use the idea of a score process and some other technics from Bickel et al. [2006] in order to construct and analyze NT-statistics. This can be seen by the fact that such applications as in Example 6 naturally appear in both papers. The difference with the above paper would be that we prefer to use test statistics of the form (2) rather than integrals or suprema of score processes.

In semi- and nonparametric models, generalized likelihood ratios from Fan et al. [2001a] and Li and Liang [2008], as well as different modifications of empirical likelihood, could also be a powerful tool for constructing NT-statistics.

Acknowledgments. Author would like to thank Fadoua Balabdaoui, Shota Gugushvili and Axel Munk for helpful discussions. Most of this research was done at Georg-August-University of Göttingen, Germany.

References

- A. N. Kolmogorov. Sulla Determinazione Empirica di una Legge di Distribuzione. *Giornale dell'Istituto Italiano degli Attuari*, 4:83–91, 1933.
- Holger Dette. A consistent test for heteroscedasticity in nonparametric regression based on the kernel method. *Journal of statistical planning and inference*, 103(1):311–329, 2002.
- Dino Sejdinovic, Bharath Sriperumbudur, Arthur Gretton, and Kenji Fukumizu. Equivalence of distance-based and rkhs-based statistics in hypothesis testing. *The Annals of Statistics*, pages 2263–2291, 2013.
- Wojciech Zaremba, Arthur Gretton, and Matthew Blaschko. B-test: A non-parametric, low variance kernel two-sample test. In *Advances in neural information processing systems*, pages 755–763, 2013.
- Kacper Chwialkowski, Heiko Strathmann, and Arthur Gretton. A kernel test of goodness of fit. In *International Conference on Machine Learning*, pages 2606–2615, 2016.
- Ya. Nikitin. *Asymptotic efficiency of nonparametric tests*. Cambridge University Press, Cambridge, 1995. ISBN 0-521-47029-3.
- J. Neyman. Smooth test for goodness of fit. *Skand. Aktuarietidskr.*, 20:150–199, 1937.
- S.S. Wilks. The large-sample distribution of the likelihood ratio for testing composite hypotheses. *Ann. Math. Stat.*, 9:60–62, 1938.
- L. Le Cam. On the asymptotic theory of estimation and testing hypotheses. In *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, 1954–1955, vol. I*, pages 129–156, Berkeley and Los Angeles, 1956. University of California Press.
- J. Neyman. Optimal asymptotic tests of composite statistical hypotheses. In *Probability and statistics: The Harald Cramér volume (edited by Ulf Grenander)*, pages 213–234. Almqvist & Wiksell, Stockholm, 1959.
- P. J. Bickel and Y. Ritov. Testing for goodness of fit: a new approach. In *Nonparametric statistics and related topics (Ottawa, ON, 1991)*, pages 51–57. North-Holland, Amsterdam, 1992.
- T. Inglot and T. Ledwina. Asymptotic optimality of data-driven Neyman’s tests for uniformity. *Ann. Statist.*, 24(5):1982–2019, 1996. ISSN 0090-5364.
- J. Fan, C. Zhang, and J. Zhang. Generalized likelihood ratio statistics and Wilks phenomenon. *Ann. Stat.*, 29(1):153–193, 2001a.
- P. J. Bickel, Y. Ritov, and T. M. Stoker. Tailor-made tests for goodness of fit to semiparametric hypotheses. *Ann. Statist.*, 34(2):721–741, 2006. ISSN 0090-5364.
- Runze Li and Hua Liang. Variable selection in semiparametric regression modeling. *Ann. Statist.*, 36(1):261–286, 2008. ISSN 0090-5364. . URL <http://dx.doi.org/10.1214/009053607000000604>.
- P. J. Bickel, C. A. J. Klaassen, Y. Ritov, and J. A. Wellner. *Efficient and adaptive estimation for semiparametric models*. Johns Hopkins Series in the Mathematical Sciences. Johns Hopkins University Press, Baltimore, MD, 1993. ISBN 0-8018-4541-6.
- I. A. Ibragimov and R. Z. Has’minskii. *Statistical estimation*, volume 16 of *Applications of Mathematics*. Springer-Verlag, New York, 1981. ISBN 0-387-90523-5. Asymptotic theory, Translated from the Russian by Samuel Kotz.
- W. C. M. Kallenberg and T. Ledwina. Consistency and Monte Carlo simulation

- of a data driven version of smooth goodness-of-fit tests. *Ann. Statist.*, 23(5): 1594–1608, 1995. ISSN 0090-5364.
- Yu. I. Ingster. Asymptotically minimax hypothesis testing for nonparametric alternatives. I, II, III. *Math. Methods Statist.*, 2:85–114, 171–189, 249–268, 1993. ISSN 1066-5307.
- V. G. Spokoiny. Adaptive hypothesis testing using wavelets. *Ann. Statist.*, 24(6):2477–2498, 1996. ISSN 0090-5364.
- M. Langovoy. Data-driven efficient score tests for deconvolution hypotheses. *Inverse Problems*, 24(2):025028 (17pp), 2008. URL <http://stacks.iop.org/0266-5611/24/025028>.
- Hajo Holzmann, Nicolai Bissantz, and Axel Munk. Density testing in a contaminated sample. *Journal of multivariate analysis*, 98(1):57–75, 2007.
- Nicolai Bissantz, Thorsten Hohage, Axel Munk, and Frits Ruymgaart. Convergence rates of general regularization methods for statistical inverse problems and applications. *SIAM Journal on Numerical Analysis*, 45(6):2610–2636, 2007.
- A. B. Owen. Empirical likelihood ratio confidence intervals for a single functional. *Biometrika*, 75(2):237–249, 1988. ISSN 0006-3444.
- R. J. Carroll, D. Ruppert, L. A. Stefanski, and C. M. Crainiceanu. *Measurement error in nonlinear models*, volume 105 of *Monographs on Statistics and Applied Probability*. Chapman & Hall/CRC, Boca Raton, FL, second edition, 2006. ISBN 978-1-58488-633-4; 1-58488-633-1. A modern perspective.
- M. Langovoy. *Data-driven goodness-of-fit tests*. University of Göttingen, Göttingen, 2007. Ph.D. thesis.
- Cristina Butucea, Catherine Matias, Christophe Pouet, et al. Adaptive goodness-of-fit testing from indirect observations. *Ann. Inst. Henri Poincaré Probab. Stat.*, 45(2):352–372, 2009.
- J. Fan, C. Zhang, and J. Zhang. Generalized likelihood ratio statistics and Wilks phenomenon. *Ann. Statist.*, 29(1):153–193, 2001b. ISSN 0090-5364.
- W. C. M. Kallenberg and T. Ledwina. Data-driven rank tests for independence. *J. Amer. Statist. Assoc.*, 94(445):285–301, 1999. ISSN 0162-1459.
- D. R. Cox. Partial likelihood. *Biometrika*, 62(2):269–276, 1975. ISSN 0006-3444.
- M. Langovoy. *Model selection, large deviations and consistency of data-driven tests*. EURANDOM Report No. 2009-007. EURANDOM, Eindhoven, 2009.
- R. L. Eubank, J. D. Hart, and V. N. LaRiccia. Testing goodness of fit via nonparametric function estimation techniques. *Comm. Statist. Theory Methods*, 22(12):3327–3354, 1993. ISSN 0361-0926.
- J. Fan. Test of significance based on wavelet thresholding and Neyman’s truncation. *J. Amer. Statist. Assoc.*, 91(434):674–688, 1996. ISSN 0162-1459.
- W. C. M. Kallenberg. The penalty in data driven Neyman’s tests. *Math. Methods Statist.*, 11(3):323–340 (2003), 2002. ISSN 1066-5307.
- L. Birgé and P. Massart. Gaussian model selection. *J. Eur. Math. Soc. (JEMS)*, 3(3):203–268, 2001. ISSN 1435-9855.
- G. Schwarz. Estimating the dimension of a model. *Ann. Statist.*, 6(2):461–464, 1978. ISSN 0090-5364.
- J. Rissanen. A universal prior for integers and estimation by minimum description length. *Ann. Statist.*, 11(2):416–431, 1983. ISSN 0090-5364.
- Felix Abramovich, Vadim Grinshtein, and Marianna Pensky. On optimality of Bayesian testimation in the normal means problem. *Ann. Statist.*, 35(5):2261–2286, 2007. ISSN 0090-5364. URL

<http://dx.doi.org/10.1214/009053607000000226>.

- Florentina Bunea, Alexandre B. Tsybakov, and Marten H. Wegkamp. Aggregation for Gaussian regression. *Ann. Statist.*, 35(4):1674–1697, 2007. ISSN 0090-5364. . URL <http://dx.doi.org/10.1214/009053606000001587>.
- F. Götze and A. N. Tikhomirov. Asymptotic distribution of quadratic forms. *Ann. Probab.*, 27(2):1072–1098, 1999. ISSN 0091-1798.
- V. Bentkus and F. Götze. Optimal rates of convergence in the CLT for quadratic forms. *Ann. Probab.*, 24(1):466–490, 1996. ISSN 0091-1798.
- L. Horváth and Q.-M. Shao. Limit theorems for quadratic forms with applications to Whittle’s estimate. *Ann. Appl. Probab.*, 9(1):146–187, 1999. ISSN 1050-5164.
- Yannick Baraud, Sylvie Huet, and Béatrice Laurent. Adaptive tests of linear hypotheses by model selection. *The Annals of Statistics*, 31(1):225–251, 2003.
- A. V. Prohorov. Sums of random vectors. *Teor. Veroyatnost. i Primenen.*, 18: 193–195, 1973. ISSN 0040-361x.

Appendix.

Proof. (Proposition 1). By the law of large numbers, as $n \rightarrow \infty$,

$$\frac{1}{n} \sum_{i=1}^n l_K(Y_i) \rightarrow_P C_P \neq 0. \quad (18)$$

We get

$$\begin{aligned} T_K &= \left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n \vec{l}(Y_i) \right\} L_k \left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n \vec{l}(Y_i) \right\}^T \\ &\geq \lambda_K^{(k)} \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \vec{l}(Y_i) \right\|^2 \\ &\geq \lambda_K^{(k)} \cdot \frac{1}{n} \left(\sum_{i=1}^n l_K(Y_i) \right)^2. \end{aligned} \quad (19)$$

By (18)

$$\begin{aligned} T_K - \pi(K, n) &\geq n \lambda_K^{(k)} \cdot \left(\frac{1}{n} \sum_{i=1}^n l_K(Y_i) \right)^2 - \pi(K, n) \\ &= n \lambda_K^{(k)} (C_K^2 + o_P(1) C_K) - \pi(K, n) \\ &= n \lambda_K^{(k)} C_K^2 + o_P(n \lambda_K^{(k)}) - \pi(K, n), \end{aligned}$$

and, because K and C_K are constants determined by fixed P , condition (11) yields

$$T_K - \pi(K, n) \rightarrow_P \infty \quad \text{as } n \rightarrow \infty. \quad (20)$$

On the other hand, by (9)

$$\left(\frac{1}{\sqrt{n}} \sum_{i=1}^n l_1(Y_i), \dots, \frac{1}{\sqrt{n}} \sum_{i=1}^n l_{K-1}(Y_i) \right) \rightarrow_P \mathcal{N},$$

where \mathcal{N} is a $(K-1)$ -dimensional multivariate normal distribution with the expectation vector equal to zero. This implies that $T_k = O_P(1)$ for all $k = 1, 2, \dots, K-1$, because

$$T_k \leq \lambda_1^{(k)} \left\| \frac{1}{n} \sum_{i=1}^n l(Y_i) \right\|^2 = \lambda_1^{(k)} O_P(1) = O_P(1)$$

and $\lambda_1^{(1)}, \lambda_1^{(2)}, \dots, \lambda_1^{(K-1)}$ are constants and $K < \infty$. Now by (20)

$$\lim_{n \rightarrow \infty} \sum_{k=1}^{K-1} P(T_k - \pi(k, n) \geq T_K - \pi(K, n)) = 0.$$

But for $d(n) \geq K$

$$P(S < K) \leq \sum_{k=1}^{K-1} P(T_k - \pi(k, n) \geq T_K - \pi(K, n)),$$

and the theorem follows. \square

Because of assumption $\langle A \rangle$ we can prove the following lemma.

Lemma 2.

$$P\left(\left|\frac{1}{n} \sum_{i=1}^n l_K(Y_i)\right| \leq \sqrt{\frac{x}{\lambda_K n}}\right) = O\left(\frac{1}{r_n}\right).$$

Proof. Denote $x_n := \sqrt{\frac{x}{\lambda_K n}}$, and remember that by $\langle C \rangle$ we have $E_P l_K(Y_i) = C_K$. Obviously, $x_n \rightarrow 0$ as $n \rightarrow \infty$. We have

$$\begin{aligned} P\left(\left|\frac{1}{n} \sum_{i=1}^n l_K(Y_i)\right| \leq x_n\right) &= P\left(-x_n \leq \frac{1}{n} \sum_{i=1}^n l_K(Y_i) \leq x_n\right) \\ &= P\left(-x_n - C_K \leq \frac{1}{n} \sum_{i=1}^n (l_K(Y_i) - E_P l_K(Y_i)) \leq x_n - C_K\right). \end{aligned}$$

Here we get two cases. First, suppose $C_K > 0$. Then we continue as follows:

$$\begin{aligned} &P\left(-x_n - C_K \leq \frac{1}{n} \sum_{i=1}^n (l_K(Y_i) - E_P l_K(Y_i)) \leq x_n - C_K\right) \\ &\leq P\left(\frac{1}{n} \sum_{i=1}^n (l_K(Y_i) - E_P l_K(Y_i)) \leq x_n - C_K\right) \\ &\leq P\left(\left|\frac{1}{n} \sum_{i=1}^n (l_K(Y_i) - E_P l_K(Y_i))\right| \geq |x_n - C_K|\right) \end{aligned}$$

(for all $n \geq$ some n_K)

$$\leq P\left(\left|\frac{1}{n} \sum_{i=1}^n (l_K(Y_i) - E_P l_K(Y_i))\right| \geq \frac{C_K}{2}\right) = O\left(\frac{1}{r_n}\right)$$

by $\langle A \rangle$, and so we proved the lemma for the case $C_K > 0$. In case if $C_K < 0$, we write

$$\begin{aligned} P\left(-x_n - C_K \leq \frac{1}{n} \sum_{i=1}^n (l_K(Y_i) - E_P l_K(Y_i)) \leq x_n - C_K\right) \\ \leq P\left(\frac{1}{n} \sum_{i=1}^n (l_K(Y_i) - E_P l_K(Y_i)) \geq -x_n - C_K\right) \end{aligned}$$

and then we proceed analogously to the previous case. \square

Proof. (Theorem 1). Let $x > 0$. Since $T_j > T_K$ if $j > K$ and (10) holds, we get by Proposition 1 that

$$\begin{aligned} P(T_S \leq x) &= \sum_{j=K}^{d(n)} P(T_j \leq x, S = j) + o(1) \\ &\leq d(n) P(T_K \leq x) + o(1) \\ &\leq d(n) P\left(\lambda_K \frac{1}{n} \left(\sum_{i=1}^n l_K(Y_i)\right)^2 \leq x\right) + o(1) \\ &= d(n) P\left(\left|\frac{1}{n} \sum_{i=1}^n l_K(Y_i)\right| \leq \sqrt{\frac{x}{\lambda_K n}}\right) + o(1). \end{aligned}$$

Now by Lemma 2 and (12) we get

$$P(T_S \leq x) = O\left(\frac{d(n)}{r_n}\right) + o(1) = o(1).$$

\square

Proof. (Theorem 2). If $S \geq K$, then $T_k - T_1 \geq \pi(k, n) - \pi(1, n)$ for some $K \leq k \leq d(n)$ and so, equivalently,

$$\begin{aligned} \left\{\frac{1}{\sqrt{n}} \sum_{i=1}^n l(Y_i)\right\}^T L \left\{\frac{1}{\sqrt{n}} \sum_{i=1}^n l(Y_i)\right\} \\ - \left\{\frac{1}{\sqrt{n}} \sum_{i=1}^n l_1(Y_i)\right\}^2 \{E_0[l_1(Y)]^T l_1(Y)\}^{-1} \geq \pi(k, n) - \pi(1, n) \end{aligned} \quad (21)$$

for some $K \leq k \leq d(n)$, where $l = (l_1, l_2, \dots, l_k)$. We can rewrite (21) in terms of the notation (13)-(14) as follows:

$$(\sqrt{n} \bar{l}_1, \dots, \sqrt{n} \bar{l}_k) L (\sqrt{n} \bar{l}_1, \dots, \sqrt{n} \bar{l}_k)^T \quad (22)$$

$$= n(\bar{l}_1, \dots, \bar{l}_k) L(\bar{l}_1, \dots, \bar{l}_k)^T \geq \frac{n \bar{l}_1^2}{E_0 l_1^2} + (\pi(k, n) - \pi(1, n)),$$

for some $K \leq k \leq d(n)$. Denote $\Delta(k, n) := \pi(k, n) - \pi(1, n)$; then with the help of (14) we rewrite (22) as

$$n Q_k(\bar{l}) \geq \Delta(k, n) + \frac{n \bar{l}_1^2}{E_0 l_1^2}, \quad (23)$$

for some $K \leq k \leq d(n)$. Clearly,

$$\begin{aligned} P_0(S \geq K) &\leq P_0((21) \text{ holds for some } K \leq k \leq d(n)) \\ &= P_0((23) \text{ holds for some } K \leq k \leq d(n)) \\ &\leq P_0(n Q_k(\bar{l}) \geq \Delta(k, n) \text{ for some } K \leq k \leq d(n)). \end{aligned}$$

But now by condition $\langle B2 \rangle$ we have

$$\begin{aligned} P_0(S \geq K) &\leq P_0(n Q_k(\bar{l}) \geq \Delta(k, n) \text{ for some } K \leq k \leq d(n)) \\ &\leq \sum_{k=K}^{d(n)} P_0(n Q_k(\bar{l}) \geq \Delta(k, n)) \\ &\leq \sum_{k=K}^{d(n)} \varphi(k; \Delta(k, n)), \end{aligned} \quad (24)$$

if only $d(n) \leq \min\{u_n, m_n\}$ (see Definition 3). Thus, because of the condition $\langle B2 \rangle$, for each $\varepsilon > 0$ there exists $K = K_\varepsilon$ such that for all $n > n(\varepsilon)$ we have $P_0(S \geq K) \leq \varepsilon$, i.e. $S = O_{P_0}(1)$.

Now, by standard inequalities, it is possible to show that $T_S = O_{P_0}(1)$. Let us write for an arbitrary real $t > 0$

$$\begin{aligned} P_0(|T_S| \geq t) &= \sum_{m=1}^{K_\varepsilon} P_0(|T_m| \geq t; S = m) \\ &\quad + \sum_{m=K_\varepsilon+1}^{d(n)} P_0(|T_m| \geq t; S = m) \\ &\leq \sum_{m=1}^{K_\varepsilon} P_0(|T_m| \geq t) + \sum_{m=K_\varepsilon+1}^{d(n)} P_0(S = m) \\ &= \sum_{m=1}^{K_\varepsilon} P_0(|T_m| \geq t) + P_0(S \geq K_\varepsilon + 1) \\ &\leq \sum_{m=1}^{K_\varepsilon} P_0(|T_m| \geq t) + \varepsilon \\ &=: R(t) + \varepsilon. \end{aligned}$$

For $t \rightarrow \infty$ we have $P_0(|T_m| \geq t) \rightarrow 0$ for every fixed m , so $R(t) \rightarrow 0$ as $t \rightarrow \infty$. Now it follows that for arbitrary $\varepsilon > 0$

$$\overline{\lim}_{t \rightarrow \infty} P_0(|T_S| \geq t) \leq \varepsilon,$$

therefore

$$\overline{\lim}_{t \rightarrow \infty} P_0(|T_S| \geq t) = 0$$

and

$$\lim_{t \rightarrow \infty} P_0(|T_S| \geq t) = 0.$$

This completes the proof. \square

Proof. (Theorem 3). Follows from Theorems 1 and 2 and our definition of consistency. \square

In the next proof we will need the following theorem from Prohorov [1973].

Theorem 4. *Let Z_1, \dots, Z_n be i.i.d. random vectors with values in \mathbb{R}^k . Let $EZ_i = 0$ and let the covariance matrix of Z_i be equal to the identity matrix. Assume $\|Z_1\|_k \leq L$ a.e. Then, for $2k \leq y^2 \leq nL^{-2}$, we have*

$$Pr\left(\|n^{-1/2} \sum_{i=1}^n Z_i\|_k \geq y\right) \leq \frac{150210}{\Gamma(k/2)} \left(\frac{y^2}{2}\right)^{\frac{k-1}{2}} \exp\left\{-\frac{y^2}{2}(1 - \eta_n)\right\},$$

where $0 \leq \eta_n \leq L y n^{-1/2}$.

Proof. (Proposition 2) Here T_S is an NT-statistic with $L_k = E_{k \times k}$ and $\lambda_1^{(k)} = \dots = \lambda_k^{(k)} = 1$. Therefore Theorem 2 is applicable. Put $s(k, n) = \sqrt{2k}$, $t(k, n) = 0.5\sqrt{n} M(k)^{-1}$. The Prohorov inequality is applicable if $M(k) \pi(k, n) \leq \sqrt{n}$ and $M^2(k) \pi(k, n) \leq n$ for all $k \leq d(n)$; therefore assumption (17) guarantees that the Prohorov inequality is applicable and, moreover, that $\langle B2 \rangle$ holds with

$$\varphi(k; y) = \frac{150210}{\Gamma(k/2)} \left(\frac{y^2}{2}\right)^{\frac{k-1}{2}} \exp\left\{-\frac{y^2}{4}\right\}. \quad (25)$$

Since φ is exponentially decreasing in y under (17), it is a matter of simple calculations to prove that $\langle B1 \rangle$ is satisfied with $u_n = d(n)$ for any sequence $\{d(n)\}$ such that (17) holds. \square