

International Journal of Modern Physics A
© World Scientific Publishing Company

CP violation in Charged Higgs Bosons decays $H^\pm \rightarrow W^\pm(\gamma, Z)$ in the Minimal Supersymmetric Standard Model (MSSM)

ABDESSLAM ARHRIB

*Département de Mathématiques, Faculté des Sciences et Techniques, B.P 416 Tangier, Morocco
and National Central University, Department of Physics, Chung-li, Taiwan 32054, R.O.C.
aarhrib@ictp.it*

RACHID BENBRIK

*Chung Yuan Christian University, Department of Physics, Chung-Li, Taiwan 320, R.O.C.
LPHEA, Département de Physique, Faculté des Sciences-Semlalia B.P 2390 Marrakech, Morocco
rbenbrik@phys.cycu.edu.tw*

MOHAMED CHABAB

*LPHEA, Département de Physique, Faculté des Sciences-Semlalia
B.P 2390 Marrakech, Morocco
mchabab@ucam.ac.ma*

WEI TING CHANG AND TZU-CHIANG YUAN

Department of Physics, National Tsing Hua University, Hsinchu, Taiwan 300, R.O.C.

One loop mediated charged Higgs bosons decays $H^\pm \rightarrow W^\pm V$, $V = Z, \gamma$ are studied in the Minimal Supersymmetric Standard Model (MSSM) with and without CP violating phases. We evaluate the MSSM contributions to these processes taking into account $B \rightarrow X_s \gamma$ constraint as well as experimental constraints on the MSSM parameters. In the MSSM, we found that in the intermediate range of $\tan \beta \lesssim 10$ and for large A_t and large μ , where the lightest top squark becomes very light and hence non-decoupled, the branching ratio of $H^\pm \rightarrow W^\pm Z$ can be of the order 10^{-3} while the branching ratio of $H^\pm \rightarrow W^\pm \gamma$ is of the order 10^{-5} . We found also that the CP violating phases of soft SUSY parameters can modify the branching ratio by about one order of magnitude. We also show that MSSM with CP violating phases lead to CP-violating asymmetry in the decays $H^+ \rightarrow W^+ V$ and $H^- \rightarrow W^- V$. Such CP asymmetry can be rather large and can reach 80% in some region of parameter space.

1. Introduction

Supersymmetric (SUSY) theories, in particular the Minimal Supersymmetric Standard Model (MSSM), are currently considered as the most theoretically well motivated extensions of the Standard Model. Recently, phenomenology of the MSSM with complex SUSY parameters has received growing attention¹. Such complex phases provide new sources of CP violation which may explain electroweak baryogenesis scenarios², and CP violating phenomena in K and B decays³. It has

been shown in ⁴ that by assuming universality of the gaugino masses at a high energy scale, the effects of complex soft SUSY parameters in the MSSM can be parameterized by two independent CP violating phases: the phase of the Higgsino mass term μ ($\text{Arg}(\mu)$) and the phase of the trilinear scalar coupling parameters $A = A_f$ ($\text{Arg}(A_f)$) of the sfermions \tilde{f} . The presence of large SUSY phases can give contributions to electric dipole moments of the electron and neutron (EDM) which exceed the experimental upper bounds. In a variety of SUSY models such phases turn out to be severely restricted by such constraints, *i.e.* $\text{Arg}(\mu) < (10^{-2})$ for a SUSY mass scale of the order of few hundred GeV ⁵. However, the possibility of having large CP violating phases can still be consistent with experimental data in any of the following three scenarios: i) Effective SUSY models ⁵, ii) Cancellation mechanism ⁶ and iii) Non-universality of trilinear couplings A_f ⁷. In the MSSM, after electroweak symmetry breaking we are left with 5 physical Higgs particles (2 charged Higgs H^\pm , 2 CP-even H^0, h^0 and one CP-odd A^0). In this study, our concerns is the charged Higgs decays $H^\pm \rightarrow W^\pm V$, $V = Z, \gamma$ with and without CP violating phases. Those channels have a very clear signature and might emerge easily at future colliders. For instance, if $H^\pm \rightarrow W^\pm Z$ is enhanced enough, this decay may lead to three leptons final state if both W and Z decay leptonically and that would be the corresponding golden mode for charged Higgs boson. Charged Higgs phenomenology has been extensively studied in the literature. It has been shown that SUSY one-loop contributions can lead to decay rates asymmetry of $H^\pm \rightarrow t\bar{b}, b\bar{t}$, $H^\pm \rightarrow \tilde{u}_i \tilde{d}_j^*, \tilde{u}_i^* \tilde{d}_j$ and $H^\pm \rightarrow \tilde{\chi}_i^+ \tilde{\chi}_j^0, \tilde{\chi}_i^- \tilde{\chi}_j^0$ ⁸. Similar study has been done for the single charged Higgs production at hadron collider ⁹ and shown that SUSY CP phases can lead to CP asymmetry.

In this paper, we will discuss both the branching ratios of $H^\pm \rightarrow W^\pm V$ with $V = Z, \gamma$ ¹⁰ as well as the CP-violating asymmetry in the decay rates $H^+ \rightarrow W^+ V$ and $H^- \rightarrow W^- V$ that emerge from the presence of CP violating phases in MSSM.

The paper is organized as follows. In section II, we describe our calculations and the one-loop renormalization scheme we will use for $H^\pm \rightarrow W^\pm V$ and present the source of CP violation in $H^\pm \rightarrow W^\pm V$. In Section III, we present our numerical results and discussions, and section IV contains our conclusions.

2. CP violation in Charged Higgs decay: $H^\pm \rightarrow W^\pm V$

2.1. One loop amplitude $H^\pm \rightarrow W^\pm V$

In MSSM, at tree level, the coupling $H^\pm W^\pm \gamma$ is absent because of electromagnetic gauge invariance $U(1)_{\text{em}}$, while the absence of $H^\pm W^\pm Z$ is due to the isospin symmetry of the kinetic Lagrangian of the Higgs doublet fields. We emphasize here that it is possible to construct models with an even larger scalar sector than 2 Higgs doublets. One of the most popular being the Higgs Triplet Model (HTM) ¹¹. A noteworthy difference between MSSM and HTM is that the HTM contains a tree level $ZW^\pm H^\mp$ coupling. Therefore, in MSSM, decays modes like $H^\pm \rightarrow W^\pm \gamma$,

$H^\pm \rightarrow W^\pm Z$ are mediated at one loop level ^{12,13,14,15,16}. Although these decays are rare processes, loop or/and threshold effects can give rise to substantial enhancement. Moreover, once worked out, any experimental deviation from the results within such a model should bring some fruitful information on the new physics and also allow to distinguish between models. Let us give the general structure of the one loop amplitude for $H^\pm \rightarrow W^\pm V$ and discuss the renormalization scheme introduced to deal with tadpoles and vector boson-scalar boson mixing.

The amplitude \mathcal{M} for a scalar decaying to two gauge bosons V_1 and V_2 can be written as

$$\mathcal{M} = \frac{g^3 \epsilon_{V_1}^{\mu*} \epsilon_{V_2}^{\nu*}}{16\pi^2 m_W} \mathcal{M}_{\mu\nu} \quad (1)$$

where ϵ_{V_i} are the polarization vectors of the V_i . According to Lorentz invariance, the general structure of the one loop amplitude $\mathcal{M}_{\mu\nu}$ of $S \rightarrow V_1^\mu V_2^\nu$ decay is

$$\mathcal{M}_{\mu\nu}(S \rightarrow W^\mu V^\nu) = \mathcal{F}_1 g_{\mu\nu} + \mathcal{F}_2 p_{1\mu} p_{2\nu} + \mathcal{F}_3 i \epsilon_{\mu\nu\rho\sigma} p_1^\rho p_2^\sigma \quad (2)$$


where $p_{1,2}$ are the momentum of V_1, V_2 vector bosons, $\mathcal{F}_{1,2,3}$ are form factors, and $\epsilon_{\mu\nu\rho\sigma}$ is the totally antisymmetric tensor. The form factor \mathcal{F}_1 has dimension 2 while the other are dimensionless. Therefore, it is expected that in case of $H^\pm \rightarrow W^\pm Z$ decay, \mathcal{F}_1 will grow quadratically with internal top quark mass while $\mathcal{F}_{2,3}$ will have only logarithmic dependence ^{13,a}. Quite contrary, for $H^\pm \rightarrow W^\pm \gamma$ decay, the electromagnetic gauge invariance implies that \mathcal{F}_1 and \mathcal{F}_2 are related by $\mathcal{F}_1 = 1/2(m_W^2 - m_{H^\pm}^2)\mathcal{F}_2$ ¹³. This means that only \mathcal{F}_2 and \mathcal{F}_3 will contribute to the decay $H^\pm \rightarrow W^\pm \gamma$ and then the amplitude of $H^\pm \rightarrow W^\pm \gamma$ will not grow quadratically with internal fermionic masses. In case of $H^\pm \rightarrow W^\pm Z$, there is no such constraint on the form factors. In fact the CP asymmetry in this process $H^\pm \rightarrow W^\pm Z$ is very large compared to the CP asymmetry in $H^\pm \rightarrow t\bar{b}$ ⁸. The reason is that $H^\pm \rightarrow W^\pm Z$ is loop dominated, while $H^\pm \rightarrow t\bar{b}$ is dominated by the tree level contribution which is CP conserving.

The one-loop amplitude for $H^\pm \rightarrow W^\pm V$ is evaluated in the 't Hooft-Feynman gauge using dimensional regularization. The typical Feynman diagrams that contribute to $H^\pm \rightarrow W^\pm V$ are depicted in Fig. 1. Those diagrams contain vertex diagrams (Fig. 1.1 \rightarrow 1.11), W^\pm - H^\pm mixing (Fig. 1.12 \rightarrow 1.14), H^\pm - G^\pm mixing (Fig. 1.15 \rightarrow 1.17) and H^\pm - W^\pm mixing (Fig. 1.18 \rightarrow 1.20).

Note that the mixing H^\pm - W^\pm (Fig. 1.12, 1.13, 1.14) vanishes for an on-shell transverse W gauge boson. There is no contribution from the W^\pm - G^\mp mixing because $\gamma G^\pm H^\mp$ and $Z G^\pm H^\mp$ vertices are absent at the tree level. All the Feynman diagrams have been generated and computed using FeynArts and FormCalc ¹⁷ packages. We also used the fortran FF-package ¹⁸ in the numerical analysis.

Although the amplitude for our process is absent at the tree level, complications like tadpole contributions and vector boson-scalar boson mixing require a

^aAs it has been shown in Ref. [13], the top quark contribution does not decouple while squarks contributions does.


4 *A. Arhrib, R. Benbrik, M. Chabab, W.T. Chang and T.-C. Yuan*Fig. 1. Generic contributions to $H^\pm \rightarrow W^\pm V$

careful treatment of renormalization. We adopt, hereafter, the on-shell renormalization scheme of Ref. [19], for the Higgs sector. In this scheme, field renormalization is performed in the manifest-symmetric version of the Lagrangian. A field renormalization constant is assigned to each Higgs doublet $\Phi_{1,2}$. We follow closely the same approach adopted in Refs. [10, 20] for W^\pm - H^\mp and H^\pm - G^\mp mixing as well as renormalization of $\gamma W^\pm H^\mp$ and $ZW^\pm H^\mp$. We will use the following on shell renormalization conditions^{10,20}:

- The renormalized tadpoles, *i.e.* the sum of tadpole diagrams $T_{h,H}$ and tadpole counter-terms $\delta_{h,H}$ vanish:

$$T_h + \delta t_h = 0, \quad T_H + \delta t_H = 0.$$

These conditions guarantee that the vevs $v_{1,2}$ appearing in the renormalized

CP violation in Charged Higgs Bosons decays $H^\pm \rightarrow W^\pm(\gamma, Z)$ in MSSM 5Fig. 2. Generic contributions to $H^\pm \rightarrow W^\pm$ and $H^\pm \rightarrow G^\pm$ mixing as well as counter-terms needed.

Lagrangian $\mathcal{L}_{\mathcal{R}}$ are located at the minimum of the one-loop potential.

- The real part of the renormalized non-diagonal self-energy $\hat{\Sigma}_{H^\pm W^\pm}(k^2)$ vanishes for an on-shell charged Higgs boson. This renormalization condition determines the counter term for $\delta[A_\nu W_\mu^\pm H^\mp]$ and $\delta[Z_\nu W_\mu^\pm H^\mp]$.

Using the Slavnov–Taylor identity, which relates $H^\pm W^\pm$ and $H^\pm G^\pm$ self energies, the last renormalization condition about $H^\pm W^\pm$ mixing is sufficient to discard the real part of the $H^\pm G^\pm$ mixing contribution as well.

2.2. CP violation in $H^\pm \rightarrow W^\pm V$

The CP violating phases of MSSM can lead to CP-violating asymmetry in the decay rates $H^+ \rightarrow W^+ V$ and $H^- \rightarrow W^- V$. In what follow, we will study the following CP-violating asymmetry

$$\delta^{CP} = \frac{\Gamma(H^+ \rightarrow W^+ V) - \Gamma(H^- \rightarrow W^- V)}{\Gamma(H^+ \rightarrow W^+ V) + \Gamma(H^- \rightarrow W^- V)}, \quad V = \gamma, Z \quad (3)$$

and show that the MSSM with complex phases together with the absorptive part emerging from some loop integrals can lead to non-vanishing δ^{CP} . Generically, in

6 *A. Arhrib, R. Benbrik, M. Chabab, W.T. Chang and T.-C. Yuan*

the case of $H^\pm \rightarrow W^\pm \gamma$, the form factors $\mathcal{F}_{1,3}$ take the following form

$$\begin{aligned}\mathcal{F}_1(H^- \rightarrow W^- \gamma) &\propto \sum_{i,j,k} a_{1,ijk} \text{PV}_{1,ijk} + a_{2,ijk} \text{PV}_{2,ijk} \\ \mathcal{F}_3(H^- \rightarrow W^- \gamma) &\propto \sum_{i,j,k} b_{1,ijk} \widetilde{\text{PV}}_{1,ijk} + b_{2,ijk} \widetilde{\text{PV}}_{2,ijk} \\ \mathcal{F}_1(H^+ \rightarrow W^+ \gamma) &\propto \sum_{i,j,k} a_{1,ijk}^* \text{PV}_{1,ijk} + a_{2,ijk}^* \text{PV}_{2,ijk} \\ \mathcal{F}_3(H^+ \rightarrow W^+ \gamma) &\propto \sum_{i,j,k} b_{1,ijk}^* \widetilde{\text{PV}}_{1,ijk} + b_{2,ijk}^* \widetilde{\text{PV}}_{2,ijk}\end{aligned}$$

where $(a, b)_{\alpha,ijk}$ ($\alpha = 1, 2$ and i, j and k label the particles inside the loop) are combination of MSSM couplings, namely charged Higgs couplings to chargino-neutralinos and charged Higgs couplings to squarks. $\text{PV}_{\alpha,ijk}$ and $\widetilde{\text{PV}}_{\alpha,ijk}$ are Passarino-Veltman functions. The CP asymmetry (3) takes the following form

$$\begin{aligned}\delta^{CP} &\propto (|\mathcal{F}_1(H^-)|^2 - |\mathcal{F}_3(H^-)|^2 - |\mathcal{F}_1(H^+)|^2 + |\mathcal{F}_3(H^+)|^2)/\text{all} \\ &\propto \sum_{i,j,k} \sum_{\alpha\beta} \Im m(a_{\alpha,ijk} a_{\beta,ijk}^*) \Im m(\text{PV}_{\alpha,ijk} \text{PV}_{\beta,ijk}^*) + (\alpha \longleftrightarrow \beta, \text{PV} \longleftrightarrow \widetilde{\text{PV}})\end{aligned}$$

It is clear from the above equation that in order to have a non-vanishing δ^{CP} we need both complex couplings $a_{\alpha,ijk}$ as well an absorptive part from the Passarino-Veltman functions. For example, diagrams Fig.1.1c and Fig.1.18 (resp. Fig.1.2 and Fig.1.19) will develop some absorptive parts if the decay channels $H^\pm \rightarrow \widetilde{\chi}_i^\pm \widetilde{\chi}_j^0$ (resp. $H^\pm \rightarrow \widetilde{q}_i^\pm \widetilde{q}'_j^*$) is open, *i.e.* $m_{H^\pm} > m_{\widetilde{\chi}_i} m_{\widetilde{\chi}_j}$ (resp. $m_{H^\pm} > m_{\widetilde{q}_i} m_{\widetilde{q}'_j}$).

3. Numerics and discussions

In our numerical evaluations, we use the following experimental input quantities²¹: $\alpha^{-1} = 129$, m_Z , m_W , m_t , $m_b = 91.1875, 80.45, 174.3, 4.7$ GeV. In the MSSM, we specify the free parameters that will be used as follow: *i*) The MSSM Higgs sector is parameterized by the CP-odd mass m_{A^0} and $\tan\beta$, taking into account one-loop radiative corrections from [22], and we assume $\tan\beta \gtrsim 3$. *ii*) The chargino-neutralino sector can be parameterized by the gaugino-mass terms M_1 , M_2 , and the Higgsino-mass term μ . For simplification, GUT relation $M_1 \approx M_2/2$ is assumed. *iii*) Sfermions are characterized by a common soft-breaking sfermion mass $M_{SUSY} \equiv \widetilde{M}_L = \widetilde{M}_R$, μ the parameter and the soft trilinear couplings for third generation scalar fermions $A_{t,b,\tau}$. For simplicity, we will take $A_t = A_b = A_\tau$.

When varying the MSSM parameters, we take into account also the following constraints: *i*) The extra contributions to the $\delta\rho$ parameter from the Higgs scalars should not exceed the current limits from precision measurements²¹: $|\delta\rho| \lesssim 0.003$. *ii*) $b \rightarrow s\gamma$ constraint. The present world average for inclusive $b \rightarrow s\gamma$ rate is²¹ $\mathcal{B}(B \rightarrow X_s \gamma) = (3.3 \pm 0.4) \times 10^{-4}$. We keep the $B \rightarrow X_s \gamma$ branching ratio in the 3σ range of $(2.1-4.5) \times 10^{-4}$. The SM part of $B \rightarrow X_s \gamma$ is calculated up to NLO




Fig. 3. Branching ratios of $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of m_{H^\pm} in the MSSM and 2HDM for $M_{SUSY} = 500$ GeV, $M_2 = 175$ GeV, $\mu = -1.4$ TeV and $A_t = A_b = A_\tau = -\mu$ for various values of $\tan\beta$.

using the expression given in [23]. While for the MSSM part, the Wilson coefficients C_7 and C_8 are included at LO in the framework of MSSM with CKM as the only source of flavor violation and are taken from [24]. *iii*) We will assume that all SUSY particles sfermions and charginos are heavier than about 100 GeV; for the light CP even Higgs we assume $m_{h^0} \gtrsim 98$ GeV and $\tan\beta \gtrsim 3$ ²⁵. For charged Higgs boson we assume that $m_{H^\pm} \geq 78$ GeV from LEP experiments²⁶. As the experimental bound on m_{h^0} is concerned, care has to be taken. Since we are using only one-loop approximation for the Higgs spectrum, and as is well known, higher order corrections²⁷ may reduce the light CP-even Higgs mass in some cases. It may be possible that some parameter space region, shown in this analysis, which survives the experimental limit $m_{h^0} \gtrsim 98$ GeV with one loop calculation may disappear, once the higher order correction to the Higgs spectrum are included.

The total width of the charged Higgs is computed at tree level from [28] without any QCD improvement for its fermionic decays $H^\pm \rightarrow \bar{f}f'$. The SUSY channels like $H^+ \rightarrow \tilde{f}_i \tilde{f}'_j$ and $H^+ \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^+$ are included when kinematically allowed. In Fig. 3, we show branching ratios of $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of charged Higgs mass for $\tan\beta = 16$ and 25. In those plots, we have shown both the pure 2HDM^b and the full MSSM contributions. As it can be seen from those plots, both for $H^\pm \rightarrow W^\pm Z$ and $H^\pm \rightarrow W^\pm \gamma$ the 2HDM contribution is rather small. Once we include the SUSY particles, we can see that the branching fractions get enhanced and can reach 10^{-3} in the case of $H^\pm \rightarrow W^\pm Z$ and 10^{-5} in the case of $H^\pm \rightarrow W^\pm \gamma$. The source of this enhancement is mainly due to the presence of

^bPure 2HDM means that we include just the 2HDM part of the MSSM that contributes here in the loop, *i.e.* only SM fermions, gauge bosons and Higgs bosons with MSSM sum rules for the Higgs sector.


8 *A. Arhrib, R. Benbrik, M. Chabab, W.T. Chang and T.-C. Yuan*

Fig. 4. Branching ratios for $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of A_t in the MSSM with $M_{SUSY} = 500$ GeV, $M_2 = 200$ GeV, $m_{H^\pm} = 500$ GeV, $A_t = A_b = A_\tau = -\mu$ and $-2\text{ TeV} < \mu < -0.1$ TeV for various values of $\tan \beta$.

scalar fermions contributions in the loop which are amplified by threshold effects from the opening of the decay $H^\pm \rightarrow \tilde{t}_i \tilde{b}_j^*$. It turns out that the contribution of charginos neutralinos loops do not enhance the branching fractions significantly as compared to scalar fermions loops. The plots also show that, the branching fractions are more important for intermediate $\tan \beta = 16$ and are slightly reduced for larger $\tan \beta = 25$.

We now illustrate in Fig. 4 the branching fraction of $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of $A_t = A_b = A_\tau = -\mu$ for $M_{SUSY} = 500$ GeV and $M_2 = 200$ GeV. Since $b \rightarrow s \gamma$ favors A_t and μ to have opposite sign, we fix $\mu = -A_t$ and in this sense μ is also varied when A_t is varied. Both for $H^\pm \rightarrow W^\pm Z$ and $H^\pm \rightarrow W^\pm \gamma$, the chargino-neutralino contributions, which are rather small, decrease with $\mu = -A_t$: the larger A_t , the smaller chargino-neutralino contributions. As one can see from these figures, the plots stop at $A_t = 1.1$ TeV and $\tan \beta = 3$, because for larger A_t the $\delta \rho$ constraint will be violated. For $\tan \beta = 10$ and 20, the plots stop for the same reason. In case of $H^\pm \rightarrow W^\pm Z$, for $A_t \lesssim 1$ TeV it is the pure 2HDM contributions which dominate and that is why it is almost independent of A_t , while for larger A_t the branching ratios increase with A_t . It is clear that the larger A_t is the larger the branching ratios, which can be of the order of 10^{-3} for $H^\pm \rightarrow W^\pm Z$ with $\tan \beta = 10$, for example. As we know from $h^0 \rightarrow \gamma \gamma$ and $h^0 \rightarrow \gamma Z$ in MSSM³⁰, the squarks contributions decouple except in the light stop mass and large A_t limit³⁰. In $H^\pm \rightarrow W^\pm V$ case, the same situation happens. As we can see from Fig. 4 (left), for intermediate A_t , $300 < A_t < 1000$ GeV, the squarks are rather heavy and hence their contributions are small compared to 2HDM one. While for large A_t the stop becomes very light $\lesssim 200$ GeV and hence it can enhance the $H^\pm \rightarrow W^\pm V$ width. Of course this enhancement is also amplified by $H^\pm \tilde{t}_{L,R} \tilde{b}_{R,L}^*$ couplings which are directly proportional to $A_{t,b}$. In the case of




Fig. 5. Branching ratios for $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) in the MSSM as a function of $Arg(A_t)$: $M_{SUSY} = 500$ GeV, $M_2 = 150$ GeV, $m_{H^\pm} = 500$ GeV, $A_t = A_b = A_\tau = -\mu = 1$ TeV and for various values of $\tan\beta$.

$H^\pm \rightarrow W^\pm \gamma$ decay, the pure 2HDM and sfermions contribution are of comparable size, the branching ratio increases with A_t . We have also studied the effect of the MSSM CP violating phases on charged Higgs decays. For $H^\pm \rightarrow W^\pm Z$ and $H^\pm \rightarrow W^\pm \gamma$ decays which are sensitive to MSSM CP violating phases through squarks and charginos-neutralinos contributions, it turns out that the effect of MSSM CP violating phases is important and can enhance the rate by about one order of magnitude. For illustration we show in Fig. 5 the effect of $A_{t,b,\tau}$ CP violating phases for $M_{SUSY} = 500$ GeV, $A_{t,b,\tau} = -\mu = 1$ TeV and $M_2 = 150$ GeV. To avoid conflict with EDM constraint we set the μ parameter to be real⁵. As it is clear, the CP phase of $A_{t,b,\tau}$ can enhance the rates of both $H^\pm \rightarrow W^\pm(Z, \gamma)$ by more than one order of magnitude. The observed cuts in the plot are due to $b \rightarrow s\gamma$ constraint.

In Fig.(6) we show the CP asymmetry δ_{CP} for $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of charged Higgs mass. As one can see in both plots, for $m_{H^\pm} \lesssim 512$ GeV, δ_{CP} is very small, of the order 10^{-3} . In this case, the main contributions to δ_{CP} arise both from diagrams Fig.1.1 and Fig.1.18 with absorptive parts and real couplings and diagrams Fig.1.2 and Fig.1.19 with complex couplings and no absorptive part (the channel $H^\pm \rightarrow \tilde{t}_i \tilde{b}_j^*$ is not yet open!). However, once $m_{H^\pm} \gtrsim 512$ GeV, the channel $H^\pm \rightarrow \tilde{t}_i \tilde{b}_j^*$ is open, the diagrams Fig.1.2 and Fig.1.19 have an absorptive part and consequently they contribute to δ_{CP} , which can go up to several percent. The thresholds of $H^\pm \rightarrow \tilde{t}_1 \tilde{b}_1^*$ at $m_{H^\pm} \approx 512$ GeV and of $H^\pm \rightarrow \tilde{t}_1 \tilde{b}_2^*$ at $m_{H^\pm} \approx 740$ GeV are clearly visible in Fig.(6).

4. Conclusion

In the framework of MSSM, we have studied charged Higgs decays into a pair of gauge bosons, namely, $H^\pm \rightarrow W^\pm Z$ and $H^\pm \rightarrow W^\pm \gamma$. We have also studied


10 *A. Arhrib, R. Benbrik, M. Chabab, W.T. Chang and T.-C. Yuan*

Fig. 6. Total δ_{CP} in the $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of m_{H^\pm} in the MSSM, with $M_{SUSY} = 500$ GeV, $M_2 = 175$ GeV, $\tan \beta = 16$, $\mu = -1.4$ TeV, $A_t = A_b = A_{\tau} = -\mu$, for different values of $\text{Arg}(A_t)$.

the effects of MSSM CP violating phases in these processes. In contrast to previous studies, we have performed the calculation in the 't Hooft-Feynman gauge and used a renormalization prescription to deal with tadpoles, $W^\pm - H^\pm$ and $G^\pm - H^\pm$ mixings. The study has been carried out taking into account the experimental constraints on the ρ parameter as well as the inclusive b -decay $b \rightarrow s\gamma$. Numerical results for the branching ratios have been presented. In the MSSM, we have shown that the branching ratio of $H^\pm \rightarrow W^\pm Z$ can reach 10^{-3} in some cases while $H^\pm \rightarrow W^\pm \gamma$ never exceed 10^{-5} . Branching ratio of the order 10^{-3} might provide an opportunity to search for a charged Higgs boson at the LHC through $H^\pm \rightarrow W^\pm Z$. The effect of MSSM CP violating phases is also found to be important and can change the size of the branching ratios of $H^\pm \rightarrow W^\pm V$ by about an order of magnitude compared to the CP conserving case. The CP-violating asymmetry in the decays $H^+ \rightarrow W^+ V$ and $H^- \rightarrow W^- V$ can be rather important and can reach 80% in some region of parameter space [31].

Acknowledgments

This work is partially supported by the ICTP through the OEA-project-49 and by the Moroccan program PROTARS-III D16/04. AA, RB, WTC and TCY are supported by the National Science Council of R.O.C. under Grant Nos. NSC96-2811-M-008-020, NSC96-2811-M-033-002 and NSC 95-2112-M-007-001.

References

1. A. Pilaftsis, Phys. Lett. B **435** (1998) 88; S. Y. Choi, M. Drees and J. S. Lee, Phys. Lett. **B481** (2000) 57; S. Heinemeyer, Eur. Phys. J. C **22**, 521 (2001); M. Carena, J. Ellis, A. Pilaftsis and C. E. Wagner, Nucl. Phys. B **586** (2000) 92.

2. M. Carena, J. M. Moreno, M. Quiros, M. Seco and C. E. Wagner, Nucl. Phys. B **599** (2001) 158; M. Brhlik, G. J. Good and G. L. Kane, Phys. Rev. D **63** (2001) 035002.
3. S. W. Baek and P. Ko, Phys. Rev. Lett. **83**, 488 (1999); T. Goto, Y. Y. Keum, T. Nihei, Y. Okada and Y. Shimizu, Phys. Lett. B **460**, 333 (1999).
4. M. Dugan, B. Grinstein and L. J. Hall, Nucl. Phys. B **255**, 413 (1985).
5. P. Nath, Phys. Rev. Lett. **66** (1991) 2565; Y. Kizukuri and N. Oshimo, Phys. Rev. D **46**, 3025 (1992); S. Pokorski, J. Rosiek and C. A. Savoy, Nucl. Phys. B **570**, 81 (2000).
6. T. Ibrahim and P. Nath, Phys. Lett. B **418** (1998) 98; Phys. Rev. D **57** (1998) 478 [Erratum-ibid. D **58** (1998) 019901; D **60** (1999) 079903; D **60** (1999) 119901; M. Brhlik, G. J. Good and G. L. Kane, Phys. Rev. D **59** (1999) 115004.
7. S. A. Abel and J. M. Frere, Phys. Rev. D **55** (1997) 1623.
8. E. Christova *et al.*, Nucl. Phys. B **639**, 263 (2002) [Erratum-ibid. B **647**, 359 (2002)]; M. Frank and I. Turan, Phys. Rev. D **76**, 016001 (2007) [arXiv:hep-ph/0703184]; M. Frank and I. Turan, arXiv:0708.0026 [hep-ph].
9. See J. Williams in S. Kraml *et al.*, “Workshop on CP studies and non-standard Higgs physics,” arXiv:hep-ph/0608079.
10. A. Arhrib *et al.*, Phys. Lett. B **644**, 248 (2007); *ibid.*, J. Phys. G **34**, 907 (2007).
11. J. F. Gunion, R. Vega and J. Wudka, Phys. Rev. D **42**, 1673 (1990).
12. A. Mendez and A. Pomarol, Nucl. Phys. B **349**, 369 (1991).
13. M. Capdequi Peyranere, H. E. Haber and P. Irulegui, Phys. Rev. D **44**, 191 (1991).
14. S. Raychaudhuri and A. Raychaudhuri, Phys. Rev. D **50** (1994) 412; *ibid.*, Phys. Lett. B **297**, 159 (1992).
15. S. Kanemura, Phys. Rev. D **61**, 095001 (2000).
16. J. Hernandez-Sanchez, M. A. Perez, G. Tavares-Velasco and J. J. Toscano, Phys. Rev. D **69**, 095008 (2004).
17. T. Hahn, Comput. Phys. Commun. **140**, 418 (2001); T. Hahn, C. Schappacher, Comput. Phys. Commun. **143**, 54 (2002); T. Hahn, M. Perez-Victoria, Comput. Phys. Commun. **118**, 153 (1999); J. Küblbeck, M. Böhm, A. Denner, Comput. Phys. Commun. **60**, 165 (1990).
18. G. J. van Oldenborgh, Comput. Phys. Commun. **66**, 1 (1991); T. Hahn, Acta Phys. Polon. B **30**, 3469 (1999).
19. A. Dabelstein, Z. Phys. C **67**, 495 (1995).
20. A. Arhrib, M. Capdequi Peyranere, W. Hollik and G. Moultaka, Nucl. Phys. B **581**, 34 (2000); H. E. Logan and S. f. Su, Phys. Rev. D **66**, 035001 (2002).
21. S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592** (2004) 1.
22. J. F. Gunion and A. Turski, Phys. Rev. D **39**, 2701 (1989); M. S. Berger, Phys. Rev. D **41**, 225 (1990); Y. Okada, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. **85**, 1 (1991).
23. A. L. Kagan and M. Neubert, Phys. Rev. D **58**, 094012 (1998).
24. M. Aoki, G. C. Cho and N. Oshimo, Nucl. Phys. B **554**, 50 (1999).
25. S. Schael *et al.* [ALEPH Collaboration], Eur. Phys. J. C **47**, 547 (2006).
26. J. Abdallah *et al.* [DELPHI Collaboration], Eur. Phys. J. C **34**, 399 (2004).
27. S. Heinemeyer, W. Hollik and G. Weiglein, Phys. Lett. B **455**, 179 (1999).
28. A. Djouadi, arXiv:hep-ph/0503173.
29. A. Abulencia *et al.* [CDF Collaboration], Phys. Rev. Lett. **96**, 042003 (2006).
30. A. Djouadi, V. Driesen, W. Hollik and J. I. Illana, Eur. Phys. J. C **1**, 149 (1998); A. Djouadi, V. Driesen, W. Hollik and A. Kraft, Eur. Phys. J. C **1**, 163 (1998).
31. A. Arhrib, R. Benbrik, W.T. Chang, W.-Y. Keung and T. C. Yuan, work in progress.