

A Fixed Point Conjecture

Elemér E Rosinger

Department of Mathematics
and Applied Mathematics
University of Pretoria
Pretoria
0002 South Africa
eerosinger@hotmail.com

Abstract

Inverse limits, unlike direct limits, can in general be void, [1]. The existence of fixed points for arbitrary mappings $T : X \rightarrow X$ is conjectured to be equivalent with the fact that related direct limits of all finite partitions of X are not void.

1. The Setup

Let X be a nonvoid set and $T : X \rightarrow X$ a mapping. We denote by

$$(1) \quad \mathcal{FP}(X)$$

the set of all finite partitions of X .

Given $x \in X$ and $\Delta \in \mathcal{FP}(X)$, then obviously

$$(2) \quad \exists A \in \Delta : \{ n \in \mathbb{N}_+ \mid T^n(x) \in A \} \text{ is infinite}$$

Here and in the sequel, we use the notation $\mathbb{N} = \{0, 1, 2, 3, \dots\}$ and $\mathbb{N}_+ = \{1, 2, 3, \dots\}$.

Let us therefore denote

$$(3) \quad \Delta(x) = \{ A \in \Delta \mid \{ n \in \mathbb{N}_+ \mid T^n(x) \in A \} \text{ is infinite} \}$$

In view of (2) we obtain

$$(4) \quad \Delta(x) \neq \phi$$

In setting up the fixed point conjecture, it is useful to consider the following two simple instances.

Example 1

1) Let $T = id_X$, that is, the *identity* mapping on X . Then for $x \in X$ and $A \subseteq X$, we clearly have

$$(5) \quad \{ n \in \mathbb{N}_+ \mid T^n(x) \in A \} = \begin{cases} \phi & \text{if } x \notin A \\ \mathbb{N}_+ & \text{if } x \in A \end{cases}$$

hence for $\Delta \in \mathcal{FP}(X)$, we obtain

$$(6) \quad \Delta(x) = \{ A \}, \text{ where } x \in A$$

2) Let T be a *constant* mapping on X , that is, $T(x) = c$, for $x \in X$, where $c \in X$ is given. Then for $x \in X$ and $A \subseteq X$, we clearly have

$$(7) \quad \{ n \in \mathbb{N}_+ \mid T^n(x) \in A \} = \begin{cases} \phi & \text{if } c \notin A \\ \mathbb{N}_+ & \text{if } c \in A \end{cases}$$

hence for $\Delta \in \mathcal{FP}(X)$, we obtain

$$(8) \quad \Delta(x) = \{ A \}, \text{ where } c \in A$$

□

Let us recall now the following natural partial order structure on $\mathcal{FP}(X)$ given by the concept of *refinement* of partitions. Namely, if $\Delta, \Delta' \in \mathcal{FP}(X)$, then we denote

$$(9) \quad \Delta \leq \Delta'$$

if and only if

$$(10) \quad \forall A' \in \Delta' : \exists A \in \Delta : A' \subseteq A$$

and in such a case, we define the mapping

$$(11) \quad \psi_{\Delta', \Delta} : \Delta' \longrightarrow \Delta$$

by

$$(12) \quad A' \subseteq A = \psi_{\Delta', \Delta}(A')$$

Then obviously

$$(13) \quad \psi_{\Delta', \Delta}(\Delta'(x)) \subseteq \Delta(x)$$

Indeed, if $A' \in \Delta'(x)$, then (3) gives

$$\{ n \in \mathbb{N}_+ \mid T^n(x) \in A' \} \text{ is infinite}$$

but in view of (10), we have

$$A' \subseteq \psi_{\Delta', \Delta}(A')$$

hence

$$\{ n \in \mathbb{N}_+ \mid T^n(x) \in \psi_{\Delta', \Delta}(A') \} \text{ is infinite}$$

thus (13).

We note now that the finite partitions

$$(14) \quad \Delta \in \mathcal{FP}(X)$$

together with the mappings

$$(15) \quad \psi_{\Delta', \Delta}, \quad \Delta, \Delta' \in \mathcal{FP}(X), \quad \Delta \leq \Delta'$$

form an *inverse* family, [1, p.191].

Furthermore, in view of (13), we also have the following stronger version of the above. For every $x \in X$, let us define the mappings

$$(16) \quad \psi_{\Delta', \Delta, x} : \Delta'(x) \longrightarrow \Delta(x), \quad \Delta, \Delta' \in \mathcal{FP}(X), \Delta \leq \Delta'$$

by

$$(17) \quad \psi_{\Delta', \Delta, x} = \psi_{\Delta', \Delta} |_{\Delta'(x)}$$

Then again, for every $x \in X$, we obtain the *inverse* family

$$(18) \quad \Delta \in \mathcal{FP}(X)$$

$$(19) \quad \psi_{\Delta', \Delta, x} : \Delta'(x) \longrightarrow \Delta(x), \quad \Delta, \Delta' \in \mathcal{FP}(X), \Delta \leq \Delta'$$

Consequently, for each $x \in X$, we can consider the *inverse limit*

$$(20) \quad \varprojlim_{\Delta \in \mathcal{FP}(X)} \Delta(x)$$

Returning now to the two simple instances in Example 1 above, we further have

Example 2

1) In the case 1) of Example 1, it follows easily that, for $x \in X$, we have

$$(21) \quad \varprojlim_{\Delta \in \mathcal{FP}(X)} \Delta(x) = \{ (\xi_{\Delta} \mid \Delta \in \mathcal{FP}(X)) \}$$

where

$$(22) \quad \xi_{\Delta} = x, \quad \Delta \in \mathcal{FP}(X)$$

2) In the case 2) of Example 1, for $x \in X$, we easily obtain

$$(23) \quad \varprojlim_{\Delta \in \mathcal{FP}(X)} \Delta(x) = \begin{cases} \phi & \text{if } x \neq c \\ \{ (\xi_{\Delta} \mid \Delta \in \mathcal{FP}(X)) \} & \text{if } x = c \end{cases}$$

where

$$(24) \quad \xi_{\Delta} = c, \quad \Delta \in \mathcal{FP}(X)$$

Remark 1

At this stage, an important fact to note is that, in general, an inverse limit such as in (20) may be void, [1, (c) in Exercise 4, p. 252], even if none of the sets $\Delta(x)$ is void, and each the mappings $\psi_{\Delta', \Delta, x}$ is surjective. Therefore, the relations (21) and (23), even if easy to establish, are as inverse limits nontrivial, in view of the arbitrariness of the sets X and mappings T involved.

A common feature of the mappings $T : X \longrightarrow X$ in both cases above is that they have *fixed points*. Namely, for the identity mapping $T = id_X$, each point $x \in X$ is such a fixed point, while for the constant mapping $T = c$, the point $x = c \in X$ is the only fixed point.

Further, as suggested by Scott Kominers, Daniel Litt and Brett Harrison, in view of the fact that a fixed point of a mapping $T : X \longrightarrow X$ is but a *particular* case of a periodic point of that mapping, or equivalently, of a fixed point of the mapping T^n , for some $n \in \mathbb{N}_+$, we are led to the

Conjecture

Given a nonvoid set X and a mapping $T : X \longrightarrow X$, then for $x \in X$, we have

$$(28) \quad \varprojlim_{\Delta \in \mathcal{FP}(X)} \Delta(x) \neq \phi \iff (\exists n \in \mathbb{N}_+ : T^n(x) = x)$$

where the issue is whether the implication \implies holds, since the converse implication is easy to establish.

References

- [1] Bourbaki N : Elements of Mathematics, Theory of Sets. Springer, 2004