

Local distinguishability of orthogonal $2 \otimes 3$ pure states

Yu Xin^{1,2*} and Runyao Duan^{1†}

¹*State Key Laboratory of Intelligent Technology and Systems,
Tsinghua National Laboratory for Information Science and Technology,*

Department of Computer Science and Technology, Tsinghua University, Beijing 100084, China,

²*Department of Physics, Tsinghua University, Beijing 100084, China*

(Dated: February 9, 2022)

We present a complete characterization for the local distinguishability of orthogonal $2 \otimes 3$ pure states except for some special cases of three states. Interestingly, we find there is a large class of four or three states that are indistinguishable by local projective measurements and classical communication (LPCC), but can be perfectly distinguishable by LOCC. That indicates the ability of LOCC for discriminating $2 \otimes 3$ states is strictly more powerful than that of LPCC, which is strikingly different from the case of multi-qubit states. We also show that classical communication plays a crucial role for local distinguishability by constructing a class of $m \otimes n$ states which require at least $2 \min\{m, n\} - 2$ rounds of classical communication in order to achieve a perfect local discrimination.

PACS numbers: 03.67.-a, 03.65.Ud, 03.67.Hk

I. INTRODUCTION

The basic problem for distinguishing quantum states by local operations and classical communication (LOCC) can be formulated as follows. Suppose two spatially separated parties, say Alice and Bob, share a quantum state which is secretly chosen from a finite set of pre-specified quantum states. They want to figure out the identity of the unknown state, but are only allowed to manipulate their own quantum systems and to communicate with each other using classical channels. This problem has received considerable attentions and has been studied extensively. Numerous interesting results have been reported, see Refs. [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36] for an incomplete list. Despite these exciting progresses, it remains unknown how to determine the local distinguishability of a set of multipartite states.

For the convenience of the readers, we give a brief review for some of these results. Walgate and coworkers showed that any two orthogonal pure states, no matter entangled or not, can always be perfectly distinguishable by LOCC [5]. Furthermore, it has been shown that the local distinguishability and the global distinguishability of two pure states have the same efficiencies [6, 7, 8, 9, 10, 11]. However, the situation changes dramatically for a set of orthogonal states with three or more members, where a perfect discrimination is generally impossible. The most surprising discovery on this topic is that there exists a set of nine $3 \otimes 3$ orthogonal pure product states which are indistinguishable by LOCC, a phenomenon known as “nonlocality without entanglement” [2, 3, 4]. Inspired by this discovery, many researchers

devoted to the local distinguishability of product states. It is now clear that any set of $2 \otimes n$ orthogonal product pure states are perfectly distinguishable by LOCC, but a set of incomplete orthogonal product states which cannot be extended by adding some additional orthogonal product state (UPB) is indistinguishable by LOCC [3, 4]. The problem of distinguishing a complete basis has been completely solved [23, 24, 25], but only very recently a characterization for the locally distinguishable $3 \otimes 3$ product states was obtained by Feng and Shi [33].

One of the main difficulties in studying local distinguishability is that there is no effective characterization of LOCC operations. In order to partially overcome this obstacle, many researchers began to employ separable operations instead of LOCC operations to study the local distinguishability. The effectiveness of this method can be roughly understood as follows. First, the class of separable operations has a rather beautiful mathematical structure. It is much easier to work with separable operations rather than LOCC operations. Second, the class of LOCC operations is a subset of the class of separable operations [2]. So one can obtain useful necessary conditions about local distinguishability by applying separable operations. Third, any separable operation can be implemented by some LOCC operation with a nonzero success probability. In other words, separable operations and LOCC operations are probabilistically equivalent. Due to these reasons, separable operations have been widely used in studying local distinguishability. We shall briefly review two kinds of results: probabilistic discrimination and perfect discrimination. Chefles first studied the distinguishability of a set of general quantum states by probabilistic LOCC and presented a necessary and sufficient condition for the unambiguous distinguishability [22]. A simplified version of this condition when only pure states are under consideration was independently obtained by Bandyopadhyay and Walgate [28], with which it was demonstrated that any three pure states are distinguishable by stochastic

*Electronic address: xiny05@mails.tsinghua.edu.cn

†Electronic address: dry@tsinghua.edu.cn

LOCC. Based on these results, Duan *et al* studied the local distinguishability of an arbitrary basis of a multipartite state space and provided a universal tight lower bound on the number of locally unambiguously distinguishable members in an arbitrary basis [29]. Walgate and Scott further showed that this lower bound plays a crucial role in deciding the generic properties such as local unambiguous distinguishability of a set of randomly chosen states [30]. Separable operations were also used to show that certain set of states are not perfectly distinguishable by LOCC. More precisely, Nathanson showed that any $(d+1)$ maximally entangled bipartite states on $d \otimes d$ cannot be perfectly distinguishable by separable operations, thus also are indistinguishable by LOCC [17]. The same result was independently obtained by Owari and Hayashi using a slightly different method [18]. It is interesting that before this result Ghosh *et al* and Fan have respectively solved the special cases of $d = 2$ and $d = 3$ using a rather different approach [15, 16]. Watrous constructed a class of bipartite subspaces having no basis distinguishable by separable operations, thus solved an open problem concerning with the environment-assisted capacity of quantum channels [26]. Hayashi *et al* studied the relation between average entanglement degree and local distinguishability of a set of orthogonal states, and provided a very general bound on the number of states which can be locally distinguishable. In Ref. [32] we systematically studied the distinguishability of quantum states by separable operations and found a new characterization for the distinguishability of quantum states by separable operations. Notably, we showed that separable operations acting on two-qubit are strictly powerful than LOCC operations. A more general class of locally indistinguishable subspaces was also constructed.

All of the above works suggest that the problem of deciding the local distinguishability of a set of general quantum states is rather complicated. Interestingly, for some very simple cases such as two-qubit states, an analytical solution is possible. Ghosh *et al* first obtained some partial results on the local distinguishability of two-qubit states using some bounds on entanglement distillation [19]. Based on an idea of Groisman and Vaidman [20], Walgate and Hardy obtained a very simple characterization for a set of $2 \otimes n$ orthogonal pure states to be perfectly distinguishable by LOCC if the owner of the qubit makes the first nontrivial measurement [21]. Employing this condition, they finally settled the local distinguishability of $2 \otimes 2$ states [21]. Another immediate consequence is that local projective measurements and classical communication (LPCC) is sufficient for the local distinguishability of multi-qubit states [36], which greatly simplifies the local distinguishability of multi-qubit states. But this is not true in general. In Ref. [3] Bennett and coworkers constructed a set of five $3 \otimes 4$ pure product states which are perfectly distinguishable by LOCC but not by LPCC. Very recently a set of $3 \otimes 3$ states with similar property was obtained by Cohen [31]. However, we still don't know whether the general POVM

is required in order to distinguish $2 \otimes 3$ states. It seems somewhat strange that the local distinguishability of $2 \otimes 3$ states when the owner of the qutrit performs the first nontrivial measurement has never been touched yet since the work of Walgate and Hardy [21].

The purpose of this paper is to study the local distinguishability of $2 \otimes 3$ states. We assume the dimension of Alice's system is 2, and the dimension of Bob's system is 3. Due to the result in Ref. [21], we only consider the case when Bob goes first, which means that Bob first does a nontrivial measurement on his own system. We find that for the discrimination of six states and five states, LOCC and LPCC are equally powerful, i.e., a set of six or five $2 \otimes 3$ states is locally distinguishable if and only if they are distinguishable by LPCC. But for four states and three states, there exists a large class of states which can be distinguished by LOCC, but not by LPCC. Therefore, we conclude that local POVM is strictly powerful than local projective measurements even for $2 \otimes 3$ system. Furthermore, we obtain a complete characterization of four $2 \otimes 3$ states that are distinguishable by LOCC but not by LPCC. For three states, such a characterization is very difficult to obtain. Nevertheless, we construct a general class of three states which are distinguishable by LOCC but not by LPCC. A feasible procedure for determining the local distinguishability of three states is also presented.

We further study the effect of classical communication for discrimination. We show that in general many rounds of classical communication are necessary. We demonstrate this result by constructing a class of $m \otimes n$ orthogonal states which requires at least $2 \min\{m, n\} - 2$ rounds to achieve a perfect discrimination. In some sense, our result is in accordance with the recent result by Owari and Hayashi in Ref. [35], where they showed that two-way classical communication can effectively increase the local distinguishability. We would like to point out that the problem studied in Ref. [35] is quite different from ours. More precisely, Ref. [35] considers the discrimination between a pure state and a mixed state, and requires the detection of pure state can be achieved perfectly. The goal is to minimize the minimal error of detecting the mixed state. Here we only consider pure states and require each state to be identified perfectly.

The rest of the paper is organized as follows. In Section II we first give a characterization for the distinguishability of $2 \otimes 3$ states by LPCC. Then in Section III and Section IV we present in sequel our results about the local distinguishability of six and five $2 \otimes 3$ states. Section V and VI devote to the local distinguishability of four and three states, respectively. In Section VII we present a non-trivial set of bipartite pure states which requires multi-round of classical communication to achieve a perfect discrimination. We conclude the paper with a brief discussion in Section VIII.

For simplicity, in what follows we shall write $|\alpha\rangle = |\beta\rangle$ for any two states which are different from each other only with a nonzero factor. Sometimes we simply

say POVM or projective measurements instead of local POVM or local projective measurements, respectively.

II. DISTINGUISHABILITY OF $2 \otimes 3$ STATES BY LPCC

In Ref. [3] Bennett and coworkers showed that any finite set of $2 \otimes n$ orthogonal product states can be perfectly distinguishable using LPCC. For $2 \otimes 3$ states this interesting result has a converse as follows.

Theorem 1. A set of $2 \otimes 3$ states are distinguishable by LPCC only if there is a set of orthogonal product states such that each of the given states can be written as a disjoint summation of these product states.

Let us make the above theorem more transparent. Suppose $\{|\psi_k\rangle : k = 1, \dots, n\}$ is a set of $2 \otimes 3$ states. Then these states are distinguishable by LPCC if and only if there exists a set of orthogonal product states $\{|\phi_j\rangle : j = 1, \dots, m\}$ and a partition of $\{1, \dots, m\}$, say S_1, \dots, S_n , such that $|\psi_k\rangle \in \text{span}\{|\phi_j\rangle : j \in S_k\}$, where $\cup_{i=1}^m S_i = \{1, \dots, m\}$ and $S_i \cap S_j = \emptyset$ for any $i \neq j$.

Proof. If Alice goes first, then it has been proven in Ref. [21] that a set of $2 \otimes n$ states is distinguishable by LOCC can be written as the sum of states from a set of orthogonal product states.

Suppose now Bob goes first. If Bob's measurement operators can be written as $P_1 = |0\rangle\langle 0| + |1\rangle\langle 1|$ and $P_2 = |2\rangle\langle 2|$, then if measurement result is 1, we can write Alice's measurement as $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$. We construct a set of orthogonal product states as follows: $\{|\alpha\rangle_A|2\rangle_B, |\alpha^\perp\rangle_A|2\rangle_B, |0\rangle_A|\beta\rangle_B, |0\rangle_A|\beta^\perp\rangle_B, |1\rangle_A|\gamma\rangle_B, |1\rangle_A|\gamma^\perp\rangle_B\}$, where $|\beta\rangle, |\beta^\perp\rangle, |\gamma\rangle, |\gamma^\perp\rangle$ belong to $\text{span}\{|0\rangle, |1\rangle\}$. It is easy to see that if a set of states is distinguishable by the LPCC we write above, they can be rewritten as sum of states from the above set.

If Bob's measurement operators can be written as: $P_1 = |0\rangle\langle 0|$, $P_2 = |1\rangle\langle 1|$ and $P_3 = |2\rangle\langle 2|$. Then we construct a set of orthogonal product states as: $\{|\alpha\rangle_A|0\rangle_B, |\alpha^\perp\rangle_A|0\rangle_B, |\beta\rangle_A|1\rangle_B, |\beta^\perp\rangle_A|1\rangle_B, |\gamma\rangle_A|2\rangle_B, |\gamma^\perp\rangle_A|2\rangle_B\}$. Obviously, if a set of states can be distinguished by the LPCC we write above, these states can be rewritten as the disjoint sum of the states from the above set. \square

III. SIX STATES

Six states form a complete basis for $2 \otimes 3$ system. By a result of Horodecki *et al* [23], we know that six states are distinguishable by LOCC only if they are product states. Conversely, by the result of Bennett *et al* mentioned above, we conclude that any $2 \otimes 3$ product basis are perfectly distinguishable by LPCC. Thus we arrive at the following:

Theorem 2. Six orthogonal $2 \otimes 3$ states are perfectly distinguishable by LPCC if and only if they form a complete

orthogonal product basis. Furthermore, the condition for the LOCC distinguishability is the same for the LPCC distinguishability.

IV. FIVE STATES

Theorem 3. Five orthogonal $2 \otimes 3$ states are locally distinguishable if and only if at most one of them is entangled.

Proof. This result is a direct consequence of a more general result presented in Ref. [32]. Here we present a self-contained proof.

Suppose the nontrivial measurement performed by Bob is $\{M_m\}$. We consider $\text{rank}(M_m)$ and sort the condition according to the M_m 's rank.

If $\text{rank}(M_m) = 3$, then on the next step, Alice should distinguish 5 orthogonal states $I \otimes M_m|\psi_i\rangle$, which is possible only if at most one of the five states is entangled. Because M_m is of full rank, it does not change the property of being entangled or separable, so at most one of the five original states is entangled.

If $\text{rank}(M_m) = 2$, and $M_m|\beta\rangle = 0$. Let B' denote the subspace orthogonal to $|\beta\rangle$, and the system AB' is $2 \otimes 2$. As Alice can distinguish at most 4 orthogonal states in AB' , one state must be eliminated after Bob's measurement, denoted as $|\psi_0\rangle = |\alpha\rangle_A|\beta\rangle_B$. The other four states are:

$$\begin{aligned} |\psi_1\rangle &= |\eta_1\rangle_{AB'} + \lambda_1|\alpha_\perp\rangle_A|\beta\rangle, \\ |\psi_2\rangle &= |\eta_2\rangle_{AB'} + \lambda_2|\alpha_\perp\rangle_A|\beta\rangle, \\ |\psi_3\rangle &= |\eta_3\rangle_{AB'} + \lambda_3|\alpha_\perp\rangle_A|\beta\rangle, \\ |\psi_4\rangle &= |\eta_4\rangle_{AB'} + \lambda_4|\alpha_\perp\rangle_A|\beta\rangle. \end{aligned} \quad (1)$$

If one of $|\eta_i\rangle$ is 0, suppose it is $|\eta_1\rangle$, then to keep orthogonality, $\lambda_1 \neq 0$, and $|\psi_1\rangle$ is product state, while the rest $\lambda_i = 0$. As $I \otimes M_m|\eta_i\rangle$ can be distinguished by Alice, at most one of $I \otimes M_m|\eta_i\rangle$ is entangled state. On the system B' , M_m is of full rank, so at most one of the left three states $|\psi_i\rangle = |\eta_i\rangle$ is entangled. We then reach the conclusion that at most one state is entangled.

Suppose $|\eta_i\rangle \neq 0$ for each i . As Alice uses projective measurements $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$ to distinguish states after Bob's measurement, the five states can be rewritten as:

$$\begin{aligned} |\psi_0\rangle &= |\alpha\rangle_A|\beta\rangle_B, \\ |\psi_1\rangle &= |0\rangle_A|\gamma_1\rangle_B + \lambda_1|\alpha_\perp\rangle_A|\beta\rangle, \\ |\psi_2\rangle &= |0\rangle_A|\gamma_2\rangle_B + \lambda_2|\alpha_\perp\rangle_A|\beta\rangle, \\ |\psi_3\rangle &= |1\rangle_A|\gamma_3\rangle_B + \lambda_3|\alpha_\perp\rangle_A|\beta\rangle, \\ |\psi_4\rangle &= |1\rangle_A|\gamma_4\rangle_B + \lambda_4|\alpha_\perp\rangle_A|\beta\rangle. \end{aligned} \quad (3)$$

To keep orthogonality relation $\langle\psi_k|\psi_l\rangle = 0$, where $k \in \{1, 2\}$ and $l \in \{3, 4\}$, we have $\lambda_1 = \lambda_2 = 0$ or $\lambda_3 = \lambda_4 = 0$. Suppose $\lambda_3 = \lambda_4 = 0$, λ_1 and λ_2 are not 0, and $\langle\gamma_3|\gamma_4\rangle = 0$.

If $|\alpha\rangle = |1\rangle$, then the five states are all product states. We suppose $|\alpha\rangle \neq |1\rangle$. Then for an arbitrary E_m , the following condition is satisfied: $\langle\gamma_1|E_m|\beta\rangle = \langle\gamma_2|E_m|\beta\rangle = \langle\gamma_3|E_m|\gamma_4\rangle = 0$.

We choose E_m satisfying $E_m|\beta\rangle \neq 0$. After Bob's measurement, Alice does a projective measurement $\{|0'\rangle, |1'\rangle\}$. If $\{|0'\rangle, |1'\rangle\} = \{|\alpha\rangle, |\alpha^\perp\rangle\}$, then as $\langle\psi_1|(|\alpha\rangle\langle\alpha| \otimes E_m)|\psi_2\rangle = 0$, we have $\langle\gamma_1|E_m|\gamma_2\rangle = 0$. So $\langle\psi_1|(I \otimes E_m)|\psi_2\rangle = \lambda_1\lambda_2^*\langle\beta|E_m|\beta\rangle = 0$, one of λ_1 and λ_2 is 0. If $\{|0'\rangle, |1'\rangle\} \neq \{|\alpha\rangle, |\alpha^\perp\rangle\}$, then after Alice's measurement, at most three states are left as the dimension of Bob's system is 3. Because $E_m|\beta\rangle \neq 0$, the first three states are not 0. So the last two states are eliminated. Then we have $E_m|\gamma_3\rangle = 0$ and $E_m|\gamma_4\rangle = 0$. $|\gamma_3\rangle$ and $|\gamma_4\rangle$ form a basis of the subspace orthogonal to $|\beta\rangle$, thus $E_m = k|\beta\rangle\langle\beta|$. As $\langle\psi_2|I \otimes E_m|\psi_3\rangle = \langle\gamma_1|E_m|\gamma_2\rangle + \lambda_1\lambda_2^*\langle\beta|E_m|\beta\rangle = k\lambda_1\lambda_2^* = 0$, that indicates one of λ_1 and λ_2 is also 0. So one of $|\psi_2\rangle$ and $|\psi_3\rangle$ is product state. Noticing that we already have three product states: $|\psi_1\rangle$, $|\psi_4\rangle$ and $|\psi_5\rangle$, we now have four product states.

If $\text{rank}(E_m) = 1$ for each m , then $E_m = \lambda_m|\beta\rangle\langle\beta|$ for some $\lambda_m > 0$. Let B' denote the subspace orthogonal to $|\beta\rangle$. Then five states can be rewritten as

$$\begin{aligned} &|\alpha\rangle_A|\beta\rangle_B + |\eta_0\rangle_{AB'}, \\ &|\alpha^\perp\rangle_A|\beta\rangle_B + |\eta_1\rangle_{AB'}, \\ &|\eta_2\rangle_{AB'}, |\eta_3\rangle_{AB'}, |\eta_4\rangle_{AB'}. \end{aligned} \quad (4)$$

where $\langle\eta_i|\eta_j\rangle = 0$.

The system AB' is $2 \otimes 2$, so at most four states can be orthogonal to each other, then one of $|\eta_0\rangle_{AB'}$ and $|\eta_1\rangle_{AB'}$ should vanish. Therefore, one of $|\psi_0\rangle$ and $|\psi_1\rangle$ is a product state, and can be written as $|\alpha\rangle_A|\beta\rangle_B$ corresponds to $E_m = \lambda_m|\beta\rangle\langle\beta|$. The number of measurement operators is at least 3 in order to satisfy $\sum E_m = I$. If the number is 3, then the measurement is actually a projective measurement. By the result in Sec. II, at most one entangled state exists. We only need to consider the case when the number is larger than 3. In this case we have at least 4 different measurement operators and each of them corresponds to a different product state. So there must be at least 4 product states.

From the discussion above, we conclude that at most one entangled state exists in the set of five orthogonal states which are distinguishable by LOCC. \square

From the above proof it is obvious that five states are perfectly distinguished by LOCC if and only if they are perfectly distinguished by LPCC.

V. FOUR STATES

Now we consider the LOCC distinguishability of four states. We have the following theorem:

Theorem 4. Four orthogonal $2 \otimes 3$ states are perfectly distinguishable by LOCC only if at least two of them are product states.

Proof. We consider the rank of measurement operators performed by Bob. There are three different cases:

Case 1: One of the measurement operators has rank 3. After Bob's measurement, none of the states is eliminated. So Alice has to distinguish 4 orthogonal states, which is possible only when at least two states are product states. As a full rank measurement operator does not change the property of being entangled or separable. Thus, at least two original states are product states.

Case 2: One of the measurement operator M_1 has rank 2. Let us assume $M_1|2\rangle_B = 0$. There are three subcases we need to consider:

Case 2.1: Two states can be written as $|\alpha\rangle|2\rangle$ and $|\beta\rangle|2\rangle$. Then there are already two product states.

Case 2.2: Only one state can be written as $|\alpha\rangle|2\rangle$. Then after Bob's measurement with outcome 1, three states are left. As Alice's measurement can only be a projective measurement of the form $\{|0\rangle, |1\rangle\}$, we can rewrite three post-measurement states as follows:

$$\begin{aligned} &|0\rangle_A|\xi_1\rangle_B + |\eta_1\rangle_A|2\rangle_B, \\ &|1\rangle_A|\xi_2\rangle_B + |\eta_2\rangle_A|2\rangle_B, \\ &|0\rangle_A|\xi_3\rangle_B + |1\rangle_A|\xi_4\rangle_B + |\eta_3\rangle_A|2\rangle_B. \end{aligned}$$

To keep orthogonality, we have $\langle\eta_1|\eta_2\rangle = \langle\alpha|\eta_2\rangle = \langle\alpha|\eta_1\rangle = 0$. As the dimension of Alice's system is 2, from the equation above, we have one of $|\eta_1\rangle$ and $|\eta_2\rangle$ is 0. Then there are two product states.

Case 2.3: No state can be written as $|\alpha\rangle|2\rangle$. The four original states must be written as:

$$\begin{aligned} &|0\rangle_A|\alpha_1\rangle_B + |\eta_1\rangle_A|2\rangle_B, \\ &|0\rangle_A|\alpha_2\rangle_B + |\eta_2\rangle_A|2\rangle_B, \\ &|1\rangle_A|\alpha_3\rangle_B + |\eta_3\rangle_A|2\rangle_B, \\ &|1\rangle_A|\alpha_4\rangle_B + |\eta_4\rangle_A|2\rangle_B. \end{aligned}$$

We choose another measurement operator M_2 satisfying $M_2|2\rangle \neq 0$, which always exists. If M_2 's condition can be sorted into above cases 2.1 and 2.2, then we reach the conclusion that two states are product states. So we only have to prove the case that $I \otimes M_2|\psi_i\rangle \neq 0$ for each i , which suggests that if the result is 2, all post-measurement states are product states.

If $|\eta_i\rangle \neq 0$, then only under the condition that $M_2|\alpha_i\rangle = \lambda_i M_2|2\rangle$ could $I \otimes M_2|\psi_i\rangle$ be product state. The states after measurement can be written as: $I \otimes M_2|\psi_i\rangle = (\lambda_i|0\rangle + |\eta_i\rangle)M|2\rangle$ or $(\lambda_i|1\rangle + |\eta_i\rangle)M|2\rangle$. Suppose at most one of $|\eta_i\rangle$ is 0, which means that there are at most one product state. If one of $|\eta_i\rangle$ is 0, then the state is already product state. To keep orthogonality, one of the rest three states satisfying $\lambda_i|0\rangle + |\eta_i\rangle$ or $\lambda_i|1\rangle + |\eta_i\rangle$ is 0, which means the state is also product state. We then have two product states. If none of $|\eta_i\rangle$ is 0, then to keep orthogonality, two $\lambda_i|0\rangle + |\eta_i\rangle$ or $\lambda_i|1\rangle + |\eta_i\rangle$ are 0, and the two states are product states.

Case 3: Every measurement operator has rank 1, and can be written as $E_i = |e_i\rangle\langle e_i|$ (unnormalized). Let $|\psi_i\rangle = |0\rangle_A|\alpha_i\rangle + |1\rangle_A|\beta_i\rangle$. After the measurement, two

states must be eliminated to keep orthogonality. It means for each $|e_i\rangle$, there are two states $|\psi_{i1}\rangle$ and $|\psi_{i2}\rangle$ satisfying

$$(I \otimes |e_i\rangle\langle e_i|)|\psi_{ik}\rangle = 0, k = 1, 2,$$

where i can take at least three different values as there are at least three measurement operators. So we have at least six orthogonality equations. Then there are two states $|\psi_i\rangle$ such that both of them have two orthogonality equations, i.e., each of them is orthogonal to two $|e_k\rangle$. We can write these orthogonality equations explicitly as follows: $\langle\alpha_i|e_{i1}\rangle = \langle\alpha_i|e_{i2}\rangle = \langle\beta_i|e_{i1}\rangle = \langle\beta_i|e_{i2}\rangle = 0$. As $|e_{i1}\rangle$ and $|e_{i2}\rangle$ are linearly independent, $|\alpha_i\rangle = |\beta_i\rangle$. It follows that two states should be product states. \square

But different from the condition of five or six states, we find two classes of four states which can only be distinguished by LOCC but not by projective measurements. The result suggests that LOCC are more powerful than LPCC. We list our results as the two following theorems and each theorem discusses one class of states. We then prove that these two classes consist of all states which can be distinguished by LOCC but not by LPCC.

Theorem 5. The following four orthogonal $2 \otimes 3$ states can be distinguishable by LOCC but not by LPCC.

$$\begin{aligned} &|0\rangle_A|0\rangle_B, \quad |1\rangle_A|\alpha\rangle_B, \\ &|0\rangle_A(a_1|1\rangle + b_1|2\rangle)_B + |1\rangle_A(c_1|\alpha^\perp\rangle + d_1|2\rangle)_B, \\ &|0\rangle_A(a_2|1\rangle + b_2|2\rangle)_B + |1\rangle_A(c_2|\alpha^\perp\rangle + d_2|2\rangle)_B, \end{aligned} \quad (5)$$

where $a_1a_2^* + c_1c_2^* = b_1b_2^* + d_1d_2^* = 0$, $|\alpha\rangle$ and $|\alpha^\perp\rangle$ belong to $\text{span}\{|0\rangle, |1\rangle\}$, $|\alpha\rangle \neq |0\rangle$, $k = -\frac{a_1a_2^*}{b_1b_2^*} = -\frac{c_1c_2^*}{d_1d_2^*}$ is real number and satisfies $0 < k < 1$.

Proof. First, to prove the states can be distinguished by LOCC, we give a set of Bob's measurement operators:

$$M_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \sqrt{k} \end{pmatrix}, \quad M_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \sqrt{1-k} \end{pmatrix}.$$

If the measurement outcome is 1, then four post-measurement states would be:

$$\begin{aligned} &|0\rangle_A|0\rangle_B, \quad |1\rangle_A|\alpha\rangle_B, \\ &|0\rangle_A(a_1|1\rangle + \sqrt{k}b_1|2\rangle)_B + |1\rangle_A(c_1|\alpha^\perp\rangle + \sqrt{k}d_1|2\rangle)_B, \\ &|0\rangle_A(a_2|1\rangle + \sqrt{k}b_2|2\rangle)_B + |1\rangle_A(c_2|\alpha^\perp\rangle + \sqrt{k}d_2|2\rangle)_B. \end{aligned} \quad (6)$$

The above four states then can be distinguished by Alice with a projective measurement $\{|0\rangle, |1\rangle\}$.

If the measurement result is 2, then two left states are:

$$\begin{aligned} &\sqrt{1-k}(b_1|0\rangle + d_1|1\rangle)_A|2\rangle_B, \\ &\sqrt{1-k}(b_2|0\rangle + d_2|1\rangle)_A|2\rangle_B. \end{aligned} \quad (7)$$

We can verify that the above two states are orthogonal, and thus can be perfectly distinguished. As a result,

the original four states can be perfectly distinguished by LOCC.

Next we shall show that the above four states cannot be distinguished by LPCC. Suppose Bob goes first. Since $\sum E_m = I$, there is a rank one projective measurement operator which can be written as: $P_1 = |\theta\rangle\langle\theta|$. If $\langle\theta|0\rangle \neq 0$, then $(I \otimes |\theta\rangle\langle\theta|)|\psi_0\rangle = \langle\theta|0\rangle|0\rangle|\theta\rangle$. To keep orthogonality between $|\psi_0\rangle$, $|\psi_2\rangle$, and $|\psi_3\rangle$, $|\theta\rangle$ should be orthogonal to $a_1|1\rangle + b_1|2\rangle$ and $a_2|1\rangle + b_2|2\rangle$. The above two states are linear independent because if $a_1|1\rangle + b_1|2\rangle = \lambda(a_2|1\rangle + b_2|2\rangle)$, then $k = -\frac{a_1a_2^*}{b_1b_2^*} = -\frac{a_1a_2^*}{b_1b_1^*} < 0$. Then $\langle\theta|1\rangle = \langle\theta|2\rangle = 0$, $|\theta\rangle = |0\rangle$. However, the projector $|0\rangle\langle 0|$ cannot keep orthogonality, because we can prove that orthogonality require $k = 1$. Thus the assumption of $\langle\theta|0\rangle \neq 0$ is incorrect. So we should have $\langle\theta|0\rangle = 0$.

Similarly, we can prove $\langle\theta|\alpha\rangle = 0$. As $|\alpha\rangle \neq |0\rangle$, $|\theta\rangle = |2\rangle$. So the rank one projective measurement operator is $|2\rangle\langle 2|$. The left projective measurement operator is $P_2 = |0\rangle\langle 0| + |1\rangle\langle 1|$. Unfortunately, by a similar argument we can show that $P_2|\psi_i\rangle$ cannot be distinguished by Alice. So the original four states cannot be distinguished by projective measurements if Bob goes first.

If instead of Bob going first, Alice goes first, then after Alice's measurement at most three states are left. In order to eliminate one state, Alice's measurement operators must be $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$. But these operators cannot keep orthogonality between the four states. So the original four states cannot be distinguished by LOCC if Alice goes first. Thus we finish our proof.

An explicit example is as follows:

$$\begin{aligned} &|0\rangle_A|0\rangle_B, \quad |1\rangle_A|1\rangle_B, \\ &|0\rangle_A(|1\rangle + |2\rangle)_B + |1\rangle_A(|0\rangle - 2|2\rangle)_B, \\ &|0\rangle_A(|1\rangle - 2|2\rangle)_B - |1\rangle_A(|1\rangle + |2\rangle)_B. \end{aligned} \quad (8)$$

The measurement operators performed by Bob are:

$$M_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \sqrt{1/2} \end{pmatrix}, \quad M_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \sqrt{1/2} \end{pmatrix}.$$

Another different class of states is as follows:

Theorem 6. The following four orthogonal states can be distinguished by LOCC but not by LPCC.

$$\begin{aligned} &|0\rangle_A|0\rangle_B, \quad |\alpha\rangle_A|1\rangle_B, \\ &a_1|1\rangle_A|0\rangle_B + b_1|\alpha^\perp\rangle_A|1\rangle_B + c_1|1\rangle_A|2\rangle_B, \\ &a_2|1\rangle_A|0\rangle_B + b_2|\alpha^\perp\rangle_A|1\rangle_B + c_2|\alpha^\perp\rangle_A|2\rangle_B, \end{aligned} \quad (9)$$

where $a_1a_2^* + b_1b_2^* + c_1c_2^*\langle\alpha^\perp|1\rangle = 0$, and $k = -\frac{a_1a_2^*}{c_1c_2^*\langle\alpha^\perp|1\rangle}$ is real number which satisfies $0 < k < 1$. $|\alpha\rangle \neq |0\rangle$ and $|1\rangle$.

Proof. Consider the following general measurement:

$$M_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \sqrt{k} \end{pmatrix}, \quad M_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \sqrt{1-k} \end{pmatrix}.$$

If the measurement outcome is 1, then after the measurement, three left states are:

$$\begin{aligned} & |0\rangle_A |0\rangle_B, \\ & a_1 |1\rangle_A |0\rangle_B + c_1 \sqrt{k} |1\rangle_A |2\rangle_B, \\ & a_2 |1\rangle_A |0\rangle_B + c_2 \sqrt{k} |\alpha^\perp\rangle_A |2\rangle_B. \end{aligned} \quad (10)$$

Alice can distinguish the states using projective measurements $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$. If the measurement result is 2, then three left states after the measurement are:

$$\begin{aligned} & |\alpha\rangle_A |1\rangle_B, \\ & b_1 |\alpha^\perp\rangle_A |1\rangle_B + \sqrt{1-k} c_1 |1\rangle_A |2\rangle_B, \\ & b_2 |\alpha^\perp\rangle_A |1\rangle_B + \sqrt{1-k} c_2 |\alpha^\perp\rangle_A |2\rangle_B. \end{aligned} \quad (11)$$

Alice can distinguish the three states using $|\alpha\rangle\langle\alpha|$ and $|\alpha^\perp\rangle\langle\alpha^\perp|$.

The next part is to prove that the four states cannot be distinguished by LPCC. As $\sum P_m = I$, there is $P_1 = |\theta\rangle\langle\theta|$. If $\langle\theta|0\rangle \neq 0$, then as $\langle\alpha|0\rangle \neq 0$, to keep orthogonality between $|\psi_1\rangle$ and $|\psi_2\rangle$ we have $\langle\theta|1\rangle = 0$. Because $\langle\alpha^\perp|0\rangle \neq 0$, to keep orthogonality between $|\psi_1\rangle$ and $|\psi_4\rangle$ we also have $\langle\theta|2\rangle = 0$, therefore $|\theta\rangle = |0\rangle$. But $|0\rangle\langle 0|$ cannot distinguish the four states, so $\langle\theta|0\rangle$ must be 0.

Using the same method, we can also first assume $\langle\theta|1\rangle \neq 0$, then we prove that $|\theta\rangle\langle\theta|$ cannot distinguish states, so $\langle\theta|1\rangle = 0$. Therefore $|\theta\rangle = |2\rangle$. But $\langle\psi_3|(I \otimes |2\rangle\langle 2|)|\psi_4\rangle \neq 0$, so the four states cannot be distinguished by projective measurements if Bob goes first.

Suppose Alice goes first. After Alice does any operator $|\theta\rangle\langle\theta|$ there are at most three states left as the dimension of Bob's system is 3. As $|\psi_3\rangle$ and $|\psi_4\rangle$ are entangled states, $(I \otimes |\theta\rangle\langle\theta|)|\psi_{3(4)}\rangle \neq 0$, so one of $(I \otimes |\theta\rangle\langle\theta|)|\psi_{1(2)}\rangle = 0$. $|\theta\rangle$ must be $|1\rangle$ or $|\alpha^\perp\rangle$. But $|1\rangle\langle 1|$ and $|\alpha^\perp\rangle\langle\alpha^\perp|$ cannot keep orthogonality. So the four states cannot be distinguished if Alice goes first. Hence we finish the proof. \square

We also give an explicit example:

$$\begin{aligned} & |0\rangle_A |0\rangle_B, \quad \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}}\right)_A |1\rangle_B, \\ & -\frac{1}{2} |1\rangle_A |0\rangle_B + \left(\frac{|0\rangle - |1\rangle}{2\sqrt{2}}\right)_A |1\rangle_B + |1\rangle_A |2\rangle_B, \\ & \frac{1}{\sqrt{2}} |1\rangle_A |0\rangle_B - \left(\frac{|0\rangle - |1\rangle}{2}\right)_A |1\rangle_B - \left(\frac{|0\rangle - |1\rangle}{\sqrt{2}}\right)_A |2\rangle_B. \end{aligned}$$

The measurement performed by Bob is given by

$$M_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \sqrt{1/2} \end{pmatrix}, \quad M_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \sqrt{1/2} \end{pmatrix}.$$

Interestingly, the above two classes of states completely characterize the local distinguishability of four $2 \otimes 3$ states.

Theorem 7. Any four $2 \otimes 3$ orthogonal states can be distinguished by LOCC but not by LPCC if and only if they can be written as one of the form in the above two theorems.

Proof. We have proved in Lemma 2 that two of the four states should be product states if they are distinguishable by LOCC.

Case 1: If there are two states that can be written as $|0\rangle_A |0\rangle_B$ and $|1\rangle_A |\eta_0\rangle_B$, where $|\eta_0\rangle$ and $|\eta_0^\perp\rangle$ belong to $\text{span}\{|0\rangle, |1\rangle\}$. Then the other two states can be written as:

$$\begin{aligned} & |0\rangle_A |\alpha_1\rangle_B + |1\rangle_A |\eta_1\rangle_B, \\ & |0\rangle_A |\alpha_2\rangle_B + |1\rangle_A |\eta_2\rangle_B, \end{aligned} \quad (12)$$

where $|\alpha_1\rangle$ and $|\alpha_2\rangle$ belong to $\text{span}\{|1\rangle, |2\rangle\}$, $|\eta_1\rangle$ and $|\eta_2\rangle$ belong to $\text{span}\{|\eta_0^\perp\rangle, |2\rangle\}$.

We assume that $|\alpha_1\rangle \neq |\alpha_2\rangle$ and $|\eta_1\rangle \neq |\eta_2\rangle$ and $|\eta_0\rangle \neq |0\rangle$. Other cases such as $|\alpha_1\rangle = \lambda|\alpha_2\rangle$ or $|\eta_1\rangle = \lambda|\eta_2\rangle$ or $|\eta_0\rangle = |0\rangle$ will be discussed later. To keep orthogonality after measurement, $\langle 0|E_m|\alpha_1\rangle = \langle 0|E_m|\alpha_2\rangle = 0$, as $|\alpha_1\rangle \neq |\alpha_2\rangle$, we have $\langle 0|E_m|1\rangle = \langle 0|E_m|2\rangle = 0$. For the same reason, $\langle \eta_0|E_m|\eta_0^\perp\rangle = \langle \eta_0|E_m|2\rangle = 0$, as $|\eta_0\rangle \neq |0\rangle$, $\langle 1|E_m|2\rangle = 0$. We obtain that E_m is diagonal under the bases $\{|0\rangle, |1\rangle, |2\rangle\}$, $E_m = \text{diag}(\lambda_0, \lambda_1, \lambda_2)$. We rewrite $|\psi_3\rangle$ and $|\psi_4\rangle$ as:

$$\begin{aligned} & |0\rangle_A (a_1 |1\rangle + a_2 |2\rangle)_B + |1\rangle_A (a_3 |0\rangle + a_4 |1\rangle + a_5 |2\rangle)_B, \\ & |0\rangle_A (b_1 |1\rangle + b_2 |2\rangle)_B + |1\rangle_A (b_3 |0\rangle + b_4 |1\rangle + b_5 |2\rangle)_B. \end{aligned}$$

To keep orthogonality, we should have $\langle\psi_3|I \otimes E_m|\psi_4\rangle = 0$, which is equivalent to

$$a_3 b_3^* \lambda_0 + (a_1 b_1^* + a_4 b_4^*) \lambda_1 + (a_2 b_2^* + a_5 b_5^*) \lambda_2 = 0.$$

There are only two linearly independent solutions to the above equation. Suppose $E_1 = \text{diag}(\lambda_0, \lambda_1, \lambda_2)$ and $E_2 = \text{diag}(\lambda'_0, \lambda'_1, \lambda'_2)$ are two independent solutions. If we have another operator E_3 , it must be written as $E_3 = aE_1 + bE_2$, so we can use only E_1 and E_2 to distinguish the states instead of using three or more operators. Therefore, there are only two measurement operators $E_1 = \text{diag}(\lambda_0, \lambda_1, \lambda_2)$ and $E_2 = \text{diag}(1-\lambda_0, 1-\lambda_1, 1-\lambda_2)$.

If $\lambda_0 \neq 0$, $E_1|0\rangle \neq 0$. After Alice's measurement $|\theta\rangle\langle\theta|$, $\langle\psi_1|(|\theta\rangle\langle\theta| \otimes E_1)|\psi_3\rangle = \lambda_0 \langle 0|\theta\rangle\langle\theta|1\rangle\langle 0|\eta_1\rangle = 0$. If $|\theta\rangle \neq |0\rangle$ or $|1\rangle$, then as $|\eta_1\rangle$ belongs to $\text{span}\{|\eta_0^\perp\rangle, |2\rangle\}$, $|\eta_1\rangle = |2\rangle$. For the same reason $|\eta_2\rangle = |\eta_1\rangle = |2\rangle$, but we have assumed in the beginning that $|\eta_2\rangle \neq |\eta_1\rangle$. So $|\theta\rangle = |0\rangle$ or $|1\rangle$ and Alice's measurement operators are: $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$. Similarly, if $\lambda_1 \neq 0$, then Alice's measurement operators must also be $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$.

From $\langle\psi_3|(|0\rangle\langle 0| \otimes E_m)|\psi_4\rangle = 0$, and $\langle\psi_3|1\rangle\langle 1| \otimes E_m|\psi_4\rangle = 0$, we have $a_1 b_1^* \lambda_2 + a_2 b_2^* \lambda_3 = 0$ and $a_3 b_3^* \lambda_1 + a_4 b_4^* \lambda_2 + a_5 b_5^* \lambda_3 = 0$. If $1-\lambda_1 \neq 0$ or $1-\lambda_2 \neq 0$, we also have $a_1 b_1^* (1-\lambda_2) + a_2 b_2^* (1-\lambda_3) = 0$ and $a_3 b_3^* (1-\lambda_1) + a_4 b_4^* (1-\lambda_2) + a_5 b_5^* (1-\lambda_3) = 0$. Then from those equations above, $a_1 b_1^* + a_2 b_2^* = 0$ and $a_3 b_3^* + a_4 b_4^* + a_5 b_5^* = 0$ stand, therefore the four states can be distinguished by

Alice first doing measurements $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$. As we have assumed these four states cannot be distinguished by projective measurements, we have either $\lambda_1 = \lambda_2 = 0$ or $1 - \lambda_1 = 1 - \lambda_2 = 0$. So the POVM consists of $E_1 = \text{diag}(1, 1, k)$ and $E_2 = \text{diag}(0, 0, 1 - k)$ which is just the case in theorem 4.

Here we will discuss other conditions we mentioned in the beginning. First, if $|\eta_0\rangle = |0\rangle$, then $|0\rangle\langle 0|$ can distinguish $|\psi_1\rangle$ and $|\psi_2\rangle$. As $|\alpha_i\rangle$ and $|\eta_i\rangle$ belong to $\text{span}\{|1\rangle, |2\rangle\}$, $|1\rangle\langle 1| + |2\rangle\langle 2|$ can distinguish $|\psi_3\rangle$ and $|\psi_4\rangle$. Therefore, the four states can be distinguished by projective measurements.

Secondly, if $|\eta_2\rangle = \lambda|\eta_1\rangle$, we choose M_1 which satisfies $M_1|\eta_1\rangle \neq 0$. Then Alice's measurement cannot be $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$, because $\langle\psi_3|(1|1\rangle\langle 1| \otimes E_1)|\psi_4\rangle = \lambda\langle\eta_1|E_1|\eta_1\rangle \neq 0$. Then to keep orthogonality after measurements $|0'\rangle\langle 0'| \otimes M_1$ and $|1'\rangle\langle 1'| \otimes M_1$, we have $\langle\psi_1|(|0'\rangle\langle 0'| \otimes E_1)|\psi_2\rangle = \langle 0|0'\rangle\langle 0'|1\rangle\langle 0|E_1|\eta_0\rangle = 0$, so $E_1|0\rangle \perp |\eta_0\rangle$. Similarly, we have $E_1|0\rangle \perp \{|\eta_0\rangle, |\alpha_1\rangle, |\alpha_2\rangle, |\eta_1\rangle\}$ and $E_1|\eta_0\rangle \perp \{|0\rangle, |\alpha_1\rangle, |\alpha_2\rangle, |\eta_1\rangle\}$.

As $|\eta_0\rangle = a_1|0\rangle + a_2|1\rangle$, $|\alpha_i\rangle = b_1|1\rangle + b_2|2\rangle$ and $|\eta_1\rangle = c_1|\eta_0^\perp\rangle + c_2|2\rangle$, at most one of the next two equations stands: $|\alpha_i\rangle = \lambda_i|\eta_0\rangle + \mu_i|\eta_1\rangle$ and $|\alpha_i\rangle = \lambda_i|0\rangle + \mu_i|\eta_1\rangle$.

If both of them does not stand, then the dimension of each set is 3. So $E_1|0\rangle = E_1|1\rangle = 0$, $E_1 = |2\rangle\langle 2|$, and the four states can be distinguishable by Bob's projective measurements $|0\rangle\langle 0| + |1\rangle\langle 1|$ and $|2\rangle\langle 2|$.

Without loss of generality, we suppose the second one does not stand, then we have $E_1|\eta_0\rangle = 0$, and the four states can be rewritten as:

$$\begin{aligned} &|0\rangle|0\rangle, |1\rangle|\eta_0\rangle, \\ &\lambda_1|0\rangle|\eta_0\rangle + |\beta_1\rangle|\eta_1\rangle, \\ &\lambda_2|0\rangle|\eta_0\rangle + |\beta_2\rangle|\eta_1\rangle. \end{aligned} \quad (13)$$

$\langle\psi_3|(I \otimes E_1)|\psi_4\rangle = \lambda_1\lambda_2\langle\eta_0|E_1|\eta_0\rangle + \langle\beta_1|\beta_2\rangle\langle\eta_1|E_1|\eta_1\rangle = 0$, so $\langle\beta_1|\beta_2\rangle = 0$, then $\langle\psi_3|\psi_4\rangle = \lambda_1\lambda_2 = 0$. One of λ_1 and λ_2 is 0, then the four states can be distinguished by projective measurements. The condition $|\alpha_2\rangle = \lambda|\alpha_1\rangle$ can be discussed similarly.

Case 2: If two product states can be written as $|0\rangle_A|0\rangle_B$ and $|\alpha\rangle_A|1\rangle_B$, then the other two are

$$\begin{aligned} &a_1|1\rangle_A|0\rangle_B + b_1|\alpha^\perp\rangle_A|1\rangle_B + |\theta_1\rangle_A|2\rangle_B, \\ &a_2|1\rangle_A|0\rangle_B + b_2|\alpha^\perp\rangle_A|1\rangle_B + |\theta_2\rangle_A|2\rangle_B. \end{aligned} \quad (14)$$

As the condition that $|\alpha\rangle = |1\rangle$ can be count into case 1, we suppose here $|\alpha\rangle \neq |1\rangle$. If one of $|\theta_i\rangle$ is 0, then the four states can only be distinguished by projective measurements $|2\rangle\langle 2|$ and $|0\rangle\langle 0| + |1\rangle\langle 1|$. So we suppose none of $|\theta_i\rangle$ is 0. We also assume that at least one of a_i , or b_i is not 0, because otherwise the four states can be distinguished by projective measurements.

To keep orthogonality between $|\psi_1\rangle$ and $|\psi_2\rangle$ after Bob's measurement, $\langle 0|E_m|1\rangle = 0$. And as $\langle\psi_1|I \otimes E_m|\psi_3\rangle = \langle 0|\theta_1\rangle\langle 0|E_m|2\rangle = 0$ and $\langle\psi_1|I \otimes E_m|\psi_3\rangle = \langle 0|\theta_2\rangle\langle 0|E_m|2\rangle = 0$. If $|\theta_1\rangle$ or $|\theta_2\rangle$ is not $|1\rangle$, then $\langle 0|E_m|2\rangle = 0$. Similarly, if $|\theta_1\rangle$ or $|\theta_2\rangle$ is not $|\alpha^\perp\rangle$, then $\langle 1|E_m|2\rangle = 0$. So E_m is diagonal, $E_m = (\lambda_1, \lambda_2, \lambda_3)$.

We also suppose $|\alpha\rangle \neq |0\rangle$. The conditions such as $|\theta_1\rangle = |\theta_2\rangle = |1\rangle$ or $|\alpha^\perp\rangle$ and $|\alpha\rangle = |0\rangle$ will be discussed later. We choose M_1 satisfying $M_1|0\rangle \neq 0$, and denote Alice's measurement operators as $|0'\rangle\langle 0'|$ and $|1'\rangle\langle 1'|$. As we suppose one of a_i is not 0, without losing generality, $a_1 \neq 0$, then $\langle\psi_1|(|0'\rangle\langle 0'| \otimes E_1)|\psi_2\rangle = \langle 0|0'\rangle\langle 0'|1\rangle\langle 0|E_1|0\rangle = 0$. $\langle 0|E_1|0\rangle \neq 0$, so either $\langle 0|0'\rangle = 0$ or $\langle 0'|1\rangle = 0$, then Alice's measurements should be $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$.

Similarly, we consider the distinguishability between $|\psi_2\rangle$, $|\psi_3\rangle$ and $|\psi_4\rangle$. If $M_1|1\rangle$ is also not 0, then Alice's measurements should be $|\alpha\rangle\langle\alpha|$ and $|\alpha^\perp\rangle\langle\alpha^\perp|$. Notice that the two sets $\{|0\rangle\langle 0|, |1\rangle\langle 1|\}$ and $\{|\alpha\rangle\langle\alpha|, |\alpha^\perp\rangle\langle\alpha^\perp|\}$ are different as we suppose $|\alpha\rangle$ is not equal to $|0\rangle$ or $|1\rangle$, Alice cannot distinguish the four states after Bob's measurement. Therefore, only one of $M_1|0\rangle$ and $M_1|1\rangle$ is not 0. As E_m is diagonal, there are at most two linear independent solutions of $(\lambda_1, \lambda_2, \lambda_3)$ which results from similar discussion as in case 1. The two measurements can be written as: $E_1 = \text{diag}(1, 0, k)$ and $E_1 = \text{diag}(0, 1, 1 - k)$. If the result is 1, then Alice's measurements should be $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$. $\langle\psi_3|(|0\rangle\langle 0| \otimes E_1)|\psi_4\rangle = k\langle\theta_1|0\rangle\langle\theta_2|0\rangle = 0$, so one of $|\theta_i\rangle$ is $|1\rangle$. For the same reason, the other is $|\alpha^\perp\rangle$. It is the case in theorem 5.

We discuss other conditions here. First, if $|\theta_1\rangle = |\theta_2\rangle = |1\rangle$, then $|\psi_3\rangle$ and $|\psi_4\rangle$ are:

$$\begin{aligned} &|1\rangle_A(a_1|0\rangle + b_1|2\rangle)_B + c_1|\alpha^\perp\rangle_A|1\rangle_B, \\ &|1\rangle_A(a_2|0\rangle + b_2|2\rangle)_B + c_2|\alpha^\perp\rangle_A|1\rangle_B. \end{aligned} \quad (15)$$

If here $|\alpha\rangle = |0\rangle$, then the four states are all product states, and can be distinguished by projective measurements. We then suppose here $|\alpha\rangle \neq |0\rangle$. To keep orthogonality between $M_m|\psi_1\rangle$ and $M_m|\psi_2\rangle$, $\langle 0|E_m|1\rangle = 0$. $\langle\psi_2|(I \otimes E_m)|\psi_3\rangle = b_1\langle\alpha|1\rangle\langle 1|E_m|2\rangle = 0$. As at least one of b_i is not 0, we have $\langle 1|E_m|2\rangle = 0$.

We choose M_1 satisfying $M_1|1\rangle \neq 0$. After Bob's measurement, Alice should distinguish four states $I \otimes M_1|\psi_i\rangle$. Suppose one of Alice's measurement operators is $|0'\rangle\langle 0'|$, then $\langle\psi_2|(|0'\rangle\langle 0'| \otimes E_1)|\psi_3\rangle = \langle\alpha|0'\rangle\langle 0'|\alpha^\perp\rangle^* b_1\langle 1|E_1|1\rangle = 0$. The equation is also satisfied for b_2 . As we suppose one of b_i is not 0, $\langle 0'|\alpha\rangle = 0$ or $\langle 0'|\alpha^\perp\rangle = 0$. So Alice's measurement should be $|\alpha\rangle\langle\alpha|$ and $|\alpha^\perp\rangle\langle\alpha^\perp|$. As $\langle 1|E_1|0\rangle = \langle 1|E_1|2\rangle = 0$, $\langle\psi_3|(|\alpha\rangle\langle\alpha| \otimes E_1)|\psi_4\rangle = \langle 1|\alpha\rangle\langle 1|\alpha\rangle^*(a_1\langle 0| + c_1\langle 2|)E_1(a_2|0\rangle + c_2|2\rangle) = 0$, therefore $(a_1\langle 0| + c_1\langle 2|)E_1(a_2|0\rangle + c_2|2\rangle) = 0$. It results in $\langle\psi_3|(|\alpha^\perp\rangle\langle\alpha^\perp| \otimes E_1)|\psi_4\rangle = \langle 1|\alpha^\perp\rangle\langle 1|\alpha^\perp\rangle^*(a_1\langle 0| + c_1\langle 2|)E_1(a_2|0\rangle + c_2|2\rangle) + b_1b_2^*\langle 1|E_1|1\rangle = b_1b_2^*\langle 1|E_1|1\rangle = 0$, so one of b_i is 0. Then the four states can be distinguished by Bob's measurement operators $|1\rangle\langle 1|$ and $|0\rangle\langle 0| + |2\rangle\langle 2|$. Similarly, the condition that $|\theta_1\rangle = |\theta_2\rangle = |\alpha^\perp\rangle$ can be discussed using the above method.

Secondly, we discuss the condition that $|\alpha\rangle = |0\rangle$ while one of $|\theta_i\rangle$ is not $|1\rangle$. The four states are:

$$\begin{aligned} &|0\rangle_A|0\rangle_B, |0\rangle_A|1\rangle_B, \\ &|1\rangle_A(a_1|0\rangle + b_1|1\rangle)_B + |\theta_1\rangle_A|2\rangle_B, \\ &|1\rangle_A(a_2|0\rangle + b_2|1\rangle)_B + |\theta_2\rangle_A|2\rangle_B. \end{aligned} \quad (16)$$

If one of $|\theta_i\rangle = |1\rangle$, we will have three product states, then the four states can be distinguished by Alice first

doing measurement $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$, so we suppose none of $|\theta_i\rangle$ is equal to $|1\rangle$. From orthogonality, we can get E_m is diagonal. We choose M_1 satisfying $M_1|0\rangle \neq 0$ or $M_1|1\rangle \neq 0$, then as we proved above, Alice's measurement should be $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$. As $\langle\psi_3|(|0\rangle\langle 0| \otimes E_1)|\psi_4\rangle = \langle\theta_1|0\rangle\langle\theta_2|0\rangle\langle 2|E_1|2\rangle = 0$, if $M_1|0\rangle \neq 0$ or $M_1|1\rangle \neq 0$, then $M_1|2\rangle = 0$. As $\sum E_m = I$, one of the measurement operator must be $|2\rangle\langle 2|$, so $\langle\theta_1|\theta_2\rangle = 0$. The four states can be distinguished by Bob's measurement $|2\rangle\langle 2|$ and $|0\rangle\langle 0| + |1\rangle\langle 1|$. \square

VI. THREE STATES

We can easily construct a class of three states that can be exactly distinguishable by LOCC but not by LPCC as follows:

Theorem 8. Three orthogonal $2 \otimes 3$ states $|\psi_i\rangle = |0\rangle_A|\eta_i\rangle_B + |1\rangle_A|\xi_i\rangle_B$, which have the following forms can be distinguishable by LOCC but not by LPCC:

$$\begin{aligned} & |0\rangle_A|0\rangle_B, \\ & |0\rangle_A(a_1|1\rangle + a_2|2\rangle)_B + |1\rangle_A(a_3|0\rangle + a_4|1\rangle + a_5|2\rangle)_B, \\ & |0\rangle_A(b_1|1\rangle + b_2|2\rangle)_B + |1\rangle_A(b_3|0\rangle + b_4|1\rangle + b_5|2\rangle)_B, \end{aligned}$$

where a_i and b_i satisfy $a_1b_1^* + a_2b_2^* + a_3b_3^* + a_4b_4^* + a_5b_5^* = 0$, $\alpha a_1b_1^* + \beta a_2b_2^* = 0$, $a_3b_3^* + \alpha a_4b_4^* + \beta a_5b_5^* = 0$, $\langle\eta_2|\eta_3\rangle \neq 0$, $\langle\xi_2|\xi_3\rangle \neq 0$, $\langle\eta_2|\eta_3\rangle + \langle\xi_2|\eta_2\rangle\langle\eta_2|\xi_3\rangle \neq 0$, $\langle\eta_2|\eta_3\rangle + \langle\xi_2|\eta_3\rangle\langle\eta_3|\xi_3\rangle \neq 0$, $|\eta_2\rangle \neq |\eta_3\rangle$, $a_3b_3^* \neq 0$ and $0 < \alpha, \beta < 1$.

Proof. The POVM consists of two parts:

$$M_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sqrt{\alpha} & 0 \\ 0 & 0 & \sqrt{\beta} \end{pmatrix}, \quad M_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \sqrt{1-\alpha} & 0 \\ 0 & 0 & \sqrt{1-\beta} \end{pmatrix}.$$

If the measurement result is 1, then after Bob's measurement the three states are:

$$\begin{aligned} & |0\rangle|0\rangle, \\ & a_3|1\rangle|0\rangle + (a_1|0\rangle + a_4|1\rangle)\sqrt{\alpha}|1\rangle + (a_2|0\rangle + a_5|1\rangle)\sqrt{\beta}|2\rangle, \\ & b_3|1\rangle|0\rangle + (b_1|0\rangle + b_4|1\rangle)\sqrt{\alpha}|1\rangle + (b_2|0\rangle + b_5|1\rangle)\sqrt{\beta}|2\rangle, \end{aligned} \quad (17)$$

Because of the relationship given above, three (unnormalized) states $I \otimes M_1|\psi_i\rangle$ are orthogonal to each other and can be distinguished if Alice performs a measurement $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$.

If measurement result is 2, then there are only two orthogonal states $I \otimes M_2|\psi_2\rangle$ and $I \otimes M_2|\psi_3\rangle$ left, and $\langle\psi_2|I \otimes E_2|\psi_3\rangle = 0$. So the two states can be distinguished by LOCC.

The next part of the proof is to prove the three states cannot be distinguished by projective measurements. Let $P_1 = |\theta\rangle\langle\theta|$. We assume $\langle\theta|0\rangle \neq 0$, then, to keep orthogonality between the three states, one state should

be eliminated if the measurement result is 1. Without losing generality, we can suppose $I \otimes P_1|\psi_3\rangle = 0$, then $\langle\theta|\eta_3\rangle = \langle\theta|\xi_3\rangle = 0$. From $\langle\psi_1|I \otimes P_1|\psi_2\rangle = 0$, we have $\langle\theta|\eta_2\rangle = 0$. The conditions in the theorem indicates that $|\eta_2\rangle, |\eta_3\rangle, |\xi_3\rangle$ are linear independent, therefore $|\theta\rangle$ does not exist. Then $\langle\theta|0\rangle$ must be equal to 0.

The left projective measurement is $P_2 = |0\rangle\langle 0| + |\theta^\perp\rangle\langle\theta^\perp|$, where $|\theta^\perp\rangle$ belongs to $\text{span}(|1\rangle, |2\rangle)$. Notice that the necessary condition for Alice to distinguish three states is at most one state is entangled, then one of $I \otimes (|0\rangle\langle 0| + |\theta^\perp\rangle\langle\theta^\perp|)|\psi_{2(3)}\rangle$ must be product state. As $P_2|\eta_i\rangle \neq P_2|\xi_i\rangle$ and $P_2|\xi_i\rangle \neq 0$, we have $P_2|\eta_i\rangle = 0$ if the state is product state. It indicates that one of $|\eta_2\rangle$ or $|\eta_3\rangle$ must be orthogonal to $|\theta^\perp\rangle$. Suppose $|\theta^\perp\rangle$ is orthogonal to $|\eta_2\rangle$, then $|\eta_2\rangle = |\theta\rangle$. Because the condition $\langle\eta_2|\eta_3\rangle + \langle\xi_2|\eta_2\rangle\langle\eta_2|\xi_3\rangle \neq 0$ is satisfied, $P_1 = |\theta\rangle\langle\theta| = |\eta_2\rangle\langle\eta_2|$ cannot keep orthogonality between $|\psi_2\rangle$ and $|\psi_3\rangle$. Thus, if Bob goes first, these states cannot be distinguished by LPCC.

On the other hand, if Alice goes first, suppose Alice's measurement is: $|0'\rangle\langle 0'|$ and $|1'\rangle\langle 1'|$. As $\langle\psi_1|(|0'\rangle\langle 0'| \otimes I)|\psi_2\rangle = a_3\langle 0|0'\rangle\langle 0'|1\rangle = 0$, Alice's measurement must be: $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$. Because $\langle\eta_2|\eta_3\rangle \neq 0$, Alice's measurement $|0\rangle\langle 0|$ cannot distinguish the four states. Therefore, the three states cannot be distinguished by LPCC. Here, we finish our proof.

We give a specific example of three states which have the form in the theorem:

$$\begin{aligned} & |0\rangle_A|0\rangle_B, \\ & |0\rangle_A(3|0\rangle + 3|2\rangle)_B + |0\rangle_A(|0\rangle + 3|1\rangle - 2|2\rangle)_B, \\ & |0\rangle_A(3|0\rangle - 2|2\rangle)_B + |0\rangle_A(2|0\rangle - 1|1\rangle + |2\rangle)_B. \end{aligned} \quad (18)$$

The POVM performed by Bob is as follows:

$$M_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sqrt{1/3} & 0 \\ 0 & 0 & \sqrt{1/2} \end{pmatrix}, \quad M_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \sqrt{2/3} & 0 \\ 0 & 0 & \sqrt{1/2} \end{pmatrix}.$$

It is easy to prove that the above three states can be distinguished by the above POVM but not by projective measurements.

For three states, to determine whether they can be distinguished by LOCC is much harder. We will give a protocol to determine whether three given orthogonal states can be distinguished.

First, the three states $|\psi_i\rangle$ are denoted as: $|0\rangle_A|\alpha_i\rangle_B + |1\rangle_A|\beta_i\rangle_B$. After the measurement, the states are $I \otimes M|\psi_i\rangle$. Taking the condition for Alice to distinguish three states into consideration, the three states after Bob's measurement can be written as: $|0^*\rangle_A|\xi_i\rangle_B + |1^*\rangle_A|\theta_i\rangle_B$, where $|\xi_i\rangle$ and $|\theta_i\rangle$ are two sets of orthogonal states of Bob's system, $|0^*\rangle$ and $|1^*\rangle$ are two specific bases of Alice's. In spite of coefficients, we have $\sum |\xi_i\rangle\langle\xi_i| = I$ and $\sum |\theta_i\rangle\langle\theta_i| = I$.

Suppose $|0^*\rangle = a|0\rangle + b|1\rangle$ and $|1^*\rangle = -b^*|0\rangle + a^*|1\rangle$. Then we have $|0^*\rangle\langle 0^*| \otimes M_m|\psi_i\rangle = |0^*\rangle_A M_m(a|\alpha_i\rangle + b|\beta_i\rangle)_B = |0^*\rangle_A|\xi_i\rangle_B$ and $|1^*\rangle\langle 1^*| \otimes M_m|\psi_i\rangle = |1^*\rangle_A M_m(-b^*|\alpha_i\rangle + a^*|\beta_i\rangle)_B = |1^*\rangle_A|\theta_i\rangle_B$.

Let $|\phi_i\rangle$ denote $a|\alpha_i\rangle + b|\beta_i\rangle$, then we can construct another set of states $|\eta_i\rangle$ satisfying $\langle\eta_i|\phi_j\rangle = 0$ for any $j \neq i$. Because $|\xi_i\rangle$ is a set of orthogonal states, $\langle\xi_i|\xi_j\rangle = \langle\xi_i|M|\phi_j\rangle = 0$. Comparing to the definition of $|\eta_i\rangle$, we have $|\eta_i\rangle = M|\xi_i\rangle$. So we can choose positive numbers λ_i , to have the following equation satisfied: $\sum \lambda_i |\eta_i\rangle \langle\eta_i| = \sum M^\dagger |\xi_i\rangle \langle\xi_i| M = M^\dagger M = E_m$.

If we let $|\varphi_i\rangle = -b^*|\xi_i\rangle + a^*|\theta_i\rangle$, then using the same method, we can find $\langle\mu_i|\varphi_j\rangle = 0$, for any $j \neq i$. We can also choose proper positive numbers ν_i to have the following equation satisfied: $\sum \nu_i |\mu_i\rangle \langle\mu_i| = E_m$.

We finally have the equation

$$\sum \nu_i |\mu_i\rangle \langle\mu_i| = \sum \lambda_i |\eta_i\rangle \langle\eta_i| = E_m.$$

There are eight independent variables a, b, λ_i, ν_i and nine equations. Getting value of the variables which satisfying the above equation, we can construct a set of POVM to distinguish the three given states. From the equation, we can see that it is much more difficult than four states' condition to judge whether the three states can be distinguished by LOCC. Actually we cannot provide an analytical characterization. Nevertheless, we can still get some results qualitatively.

If the equation is satisfied for any a and b , we can adjust a and b to make E_m satisfy $\sum E_m = I$. If the equation is satisfied for a certain value a_0 and b_0 , then we only have an $E_0 = I$. Therefore, Bob can only do a trivial operation on his system. Then we only need to judge whether these states can be distinguished if Alice goes first, which is much easier. If there is no solution to the equation, then the three states cannot be locally distinguished.

VII. A NONTRIVIAL EXAMPLE REQUIRING MULTI-ROUND CLASSICAL COMMUNICATION

Now we turn to discuss the role of classical communication in local discrimination. We find a set of $m \otimes n$ states needs at least $2 \min\{m, n\} - 2$ rounds to be distinguished using LOCC.

First, suppose $m = n$, where m is the dimension of the first system, Alice's system, and n is the dimension of the second system, Bob's system. We construct a set of states as follows:

$$\begin{array}{llll} |0\rangle|\eta_{00}\rangle + & |\alpha_{00}\rangle|0\rangle + & |1\rangle|\eta_{10}\rangle + & \cdots + |n-1\rangle|n-2\rangle \\ |0\rangle|\eta_{01}\rangle, & |\alpha_{01}\rangle|0\rangle, & |1\rangle|\eta_{11}\rangle, & \cdots, |n-1\rangle|n-1\rangle \\ |0\rangle|\eta_{02}\rangle, & |\alpha_{02}\rangle|0\rangle, & |1\rangle|\eta_{12}\rangle, & \cdots \\ \vdots & \vdots & \vdots & \cdots \\ |0\rangle|\eta_{0n-1}\rangle, & |\alpha_{0n-2}\rangle|0\rangle, & |1\rangle|\eta_{10}\rangle, & \cdots \end{array}$$

where $\{|\eta_{ki}\rangle, 0 \leq i \leq n-k-1\}$ is an orthonormal basis for the orthogonal complement of $\text{span}\{|0\rangle, \dots, |k-1\rangle\}$ and $\{|\alpha_{li}\rangle, 0 \leq i \leq n-l-2\}$ is an orthonormal basis for the orthogonal complement of $\text{span}\{|0\rangle, \dots, |l\rangle\}$. $\langle\eta_{k1}|\eta_{l1}\rangle \neq 0$, $\langle\alpha_{k1}|\alpha_{l1}\rangle \neq 0$, and $\langle\eta_{k0}|i\rangle \neq 0$ for $k \leq i \leq n-1$,

$\langle\alpha_{l0}|j\rangle \neq 0$ for $l+1 \leq j \leq n-1$. The total number of the states is $n^2 - 2n + 3$.

Theorem 9. The above $n^2 - 2n + 3$ states need at least $2n - 2$ rounds classical communication to be distinguishable by LOCC.

Proof. The key idea is to prove that measurement operators should be projective measurements. Suppose Alice goes first, and let E_m denote Alice's POVM operator with outcome m . As $\langle\eta_{01}|\eta_{k1}\rangle \neq 0$ and the orthogonality between $|0\rangle_A|\eta_{01}\rangle$ and $|k\rangle_A|\eta_{k1}\rangle$ should be kept after the measurement, we have $\langle 0|E_m|k\rangle = 0$, similarly, $\langle j|E_m|k\rangle = 0$. Therefore E_m is diagonal, $E_m = \text{diag}(\lambda_0, \lambda_1, \dots, \lambda_{n-1})$.

To keep orthogonality of $|\psi_0\rangle$ and $|\alpha_{0i}\rangle_A|0\rangle_B$, E_m should also be diagonal under the bases $\{|\alpha_{0i}\rangle, |0\rangle\}$, then $E_m = \lambda'_0|\alpha_{00}\rangle\langle\alpha_{00}| + \lambda'_1|\alpha_{01}\rangle\langle\alpha_{01}| + \cdots + \lambda'_{n-2}|\alpha_{0n-2}\rangle\langle\alpha_{0n-2}| + \lambda'_{n-1}|0\rangle\langle 0|$.

From the restriction in the theorem, we have $\langle\alpha_{00}|j\rangle \neq 0$, for any $j \neq 0$. Therefore $\langle\alpha_{00}|E_m|j\rangle = \lambda_j\langle\alpha_{00}|j\rangle = \lambda'_0\langle\alpha_{00}|j\rangle$, $\lambda_j = \lambda'_0$, so $E_m = \text{diag}(\lambda_0, \lambda'_0, \dots, \lambda'_0)$.

If λ'_0 and λ_0 are both not 0, then after Alice's measurement, Bob should do a nontrivial operation on his own system according to Alice's result. We denote F_n as Bob's operator. As we discussed above, we can conclude that F_n is diagonal on bases $\{|0\rangle, |1\rangle, \dots, |n-1\rangle\}$. To keep orthogonality of $|\psi_0\rangle$ and $|0\rangle_A|\eta_{0j}\rangle_B$, we can also rewrite $F_n = \mu'_0|\eta_{00}\rangle\langle\eta_{00}| + \mu'_1|\eta_{01}\rangle\langle\eta_{01}| + \cdots + \mu'_{n-1}|\eta_{0n-1}\rangle\langle\eta_{0n-1}|$. Following the steps above, as $\langle\eta_{00}|j\rangle \neq 0$, we have $\mu_j = \mu'_0$ for arbitrary j . Thus $F_n = \mu'_0 I$ is a trivial operator. Finally, either $\lambda_0 = 0$ or $\lambda'_0 = 0$.

Notice that this result also suggest that these states cannot be distinguished if Bob goes first. As we can see the process as Alice first does a diagonal operator on her system, $\lambda_0 = \lambda'_0 = 1$. As they are both not 0, we have proved that in the above paragraph that after Alice's measurement, these states cannot be distinguished.

We go back to Alice's first nontrivial measurement. Due to the above result, Alice's measurement only has two measurement operators: $E_1 = \text{diag}(1, 0, \dots, 0)$ and $E_2 = \text{diag}(0, 1, \dots, 1)$. If the measurement outcome is 1, Bob only needs to do projective measurements to distinguish the left states. If the measurement outcome is 2, the system is then $(n-1) \times n$.

It is then Bob's turn to do measurement. Following the method we used above, we can similarly prove that Bob's measurement must be $E_1 = \text{diag}(1, 0, \dots, 0)$ and $E_2 = \text{diag}(0, 1, \dots, 1)$. By induction, we find the number of rounds needed for distinguishing is $2n - 2$. Hence we complete the proof. \square

In general case, $m \neq n$, we can suppose $m < n$, then to distinguish the set of states we give in the theorem $2m - 2$ rounds are needed. We can also construct a set of states which requires $2m - 1$ rounds to achieve a perfect

discrimination. An explicit construction is as follows:

$$\begin{aligned}
& |\alpha_{00}\rangle|0\rangle + |0\rangle|\eta_{00}\rangle + |\alpha_{10}\rangle|1\rangle + \cdots + |m-1\rangle|m-2\rangle \\
& |\alpha_{01}\rangle|0\rangle, |0\rangle|\eta_{01}\rangle, |\alpha_{11}\rangle|1\rangle, \cdots, |m-1\rangle|m-1\rangle \\
& |\alpha_{02}\rangle|0\rangle, |1\rangle|\eta_{02}\rangle, |\alpha_{12}\rangle|1\rangle, \cdots, \\
& \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \cdots \\
& |\alpha_{0m-1}\rangle|0\rangle, |1\rangle|\eta_{0m-2}\rangle, |\alpha_{10}\rangle|1\rangle, \cdots,
\end{aligned}$$

where $\{|\eta_{ki}\rangle, 0 \leq i \leq n-k-2\}$ is an orthonormal basis for the orthogonal complement of $\text{span}\{|0\rangle, \dots, |k\rangle\}$ and $\{|\alpha_{li}\rangle, 0 \leq i \leq m-l-1\}$ is an orthonormal basis for the orthogonal complement of $\text{span}\{|0\rangle, \dots, |l-1\rangle\}$. $\langle\eta_{k1}|\eta_{l1}\rangle \neq 0$, $\langle\alpha_{k1}|\alpha_{l1}\rangle \neq 0$, and $\langle\eta_{k0}|i\rangle \neq 0$ for $k+1 \leq i \leq n-1$, $\langle\alpha_{l0}|j\rangle \neq 0$ for $l \leq j \leq m-1$.

The proof of the example above is almost the same as the previous one.

VIII. CONCLUSION

We have studied the local distinguishability of $2 \otimes 3$ states when the owner of the qutrit performs the first nontrivial measurement. We surprisingly find that for certain four or three states we need to perform the general

local POVM in order to achieve a perfect discrimination, only LPCC is not sufficient. We have almost completely characterized the local distinguishability of $2 \otimes 3$ states except for some special case when only three states are under consideration. It would be of great interest to extend these results to $2 \otimes n$ states where $n > 3$.

We further construct a special set of $m \otimes n$ states which require at least $2 \min\{m, n\} - 2$ rounds classical communication to finish the discrimination. Our result indicates that classical communication plays a crucial role in local discrimination. An interesting open problem is to construct a set of states which may require more rounds to achieve a perfect discrimination.

We are indebted to the colleagues in the Quantum Computation and Quantum Information Research Group for many enjoyable conversations. In particular, we sincerely thank Prof. M. Ying for his numerous encouragement and constant support on this research. R. Duan is also grateful to Y. Feng for sharing his insight on LOCC discrimination. This work was partly supported by the National Natural Science Foundation of China (Grant Nos. 60621062 and 60503001) and the Hi-Tech Research and Development Program of China (863 project) (Grant No. 2006AA01Z102).

-
- [1] A. Peres and W. K. Wootters, Phys. Rev. Lett. **66**, 1119 (1991).
 - [2] C. H. Bennett, D. P. DiVincenzo, C. A. Fuchs, T. Mor, E. Rains, P. W. Shor, J. A. Smolin, and W. K. Wootters, Phys. Rev. A **59**, 1070 (1999).
 - [3] C. H. Bennett, D. P. DiVincenzo, T. Mor, P. W. Shor, J. A. Smolin, B. M. Terhal, Phys. Rev. Lett. **82**, 5385 (1999).
 - [4] D. P. DiVincenzo, T. Mor, P. W. Shor, J. A. Smolin, and B. M. Terhal, Comm. Math. Phys. **238**, 379 (2003).
 - [5] J. Walgate, A. J. Short, L. Hardy, and V. Vedral, Phys. Rev. Lett. **85**, 4972 (2000).
 - [6] S. Virmani, M. F. Sacchi, M. B. Plenio, and D. Markham, Phys. Lett. A **288**, 62 (2001).
 - [7] Y.-X. Chen and D. Yang, Phys. Rev. A **64**, 064303 (2001); Phys. Rev. A **65**, 022320 (2002).
 - [8] M. Hillery and J. Mimih, Phys. Rev. A **67**, 042304 (2003).
 - [9] Z. Ji, H. Cao, and M. Ying, Phys. Rev. A **71**, 032323 (2005).
 - [10] A. Acín, E. Bagan, M. Baig, Ll. Masanes, and R. Muñoz-Tapia, Phys. Rev. A **71**, 032338 (2005).
 - [11] Y. Ogata, J. Phys. A: Math. Gen. **39**, 3059 (2006).
 - [12] B. M. Terhal, D. P. DiVincenzo, and D. W. Leung, Phys. Rev. Lett. **86**, 5807 (2001).
 - [13] D. P. DiVincenzo, D. W. Leung, and B. M. Terhal, IEEE Trans. Inf. Theory **48**, 580 (2002).
 - [14] T. Eggeling and R. F. Werner, Phys. Rev. Lett. **89**, 097905 (2002).
 - [15] S. Ghosh, G. Kar, A. Roy, A. Sen(De), and U. Sen, Phys. Rev. Lett. **87**, 277902 (2001).
 - [16] H. Fan, Phys. Rev. Lett. **92**, 177905 (2004).
 - [17] M. Nathanson, J. Math. Phys. **46**, 062103 (2005).
 - [18] M. Owari and M. Hayashi, Phys. Rev. A **74**, 032108 (2006), see also quant-ph/0411143 for a preliminary version.
 - [19] S. Ghosh, G. Kar, A. Roy, D. Sarkar, A. Sen(De), and U. Sen, Phys. Rev. A **65**, 062307 (2002).
 - [20] B. Groisman, L. Vaidman, J. Phys. A **34** 6881 (2001), quant-ph/0103084.
 - [21] J. Walgate and L. Hardy, Phys. Rev. Lett. **89**, 147901 (2002).
 - [22] A. Chefles, Phys. Rev. A **69**, 050307(R) (2004).
 - [23] M. Horodecki, A. Sen(De), U. Sen, and K. Horodecki, Phys. Rev. Lett. **90**, 047902 (2003).
 - [24] P.-X. Chen and C.-Z. Li, Phys. Rev. A **70**, 022306 (2004).
 - [25] S. De Rinaldis, Phys. Rev. A **70**, 022309 (2004).
 - [26] J. Watrous, Phys. Rev. Lett. **95**, 080505 (2005).
 - [27] M. Hayashi, D. Markham, M. Murao, M. Owari, and S. Virmani, Phys. Rev. Lett. **96**, 040501 (2006).
 - [28] S. Bandyopadhyay and J. Walgate, quant-ph/0612013.
 - [29] R. Y. Duan, Y. Feng, Z. F. Ji, and M. S. Ying, Phys. Rev. Lett. **98**, 230502 (2007).
 - [30] J. Walgate and A. Scott, arXiv:0709.4238 [quant-ph].
 - [31] S. M. Cohen, Phys. Rev. A **75**, 052313 (2007).
 - [32] R. Y. Duan, Y. Feng, Y. Xin, and M. S. Ying, arXiv:0705.0795 [quant-ph].
 - [33] Y. Feng and Y. Y. Shi, arXiv:0707.3581[quant-ph].
 - [34] S. M. Cohen, arXiv 0708.2396 [quant-ph].
 - [35] M. Owari, M. Hayashi, arXiv:0708.3154[quant-ph].
 - [36] R. Y. Duan and M. S. Ying, arXiv:0708.3559 [quant-ph].