# Pseudoclassical description of scalar particle in non-Abelian background and path-integral representations

R. Fresneda\*and D. Gitman<sup>†</sup>

February 9, 2022

#### Abstract

Path-integral representations for a scalar particle propagator in non-Abelian external backgrounds are derived. To this aim, we generalize the procedure proposed by Gitman and Schvartsman 1993 of path-integral construction to any representation of SU(N) given in terms of antisymmetric generators. And for arbitrary representations of SU(N), we present an alternative construction by means of fermionic coherent states. From the path-integral representations we derive pseudoclassical actions for a scalar particle placed in non-Abelian backgrounds. These actions are classically analyzed and then quantized to prove their consistency.

Insituto de Física, Universidade de São Paulo, São Paulo, Brasil.

## 1 Introduction

QFT with external backgrounds is a good approach for describing many physical situations and effects. If the external background is strong enough it has to be taken into account non-perturbatively. The corresponding methods for QED are well developed and were fruitfully applied for a number of calculations, see e.g. [1] and citations therein. The external background concept in non-Abelian QFT is less developed and meets some difficulties (there is no gauge invariant way of introducing a non-Abelian external field). However, the undeniable existence of physical situations where there is a sufficiently strong quantized non-Abelian field often serves as a physical justification for treating this field as an external classical field, in spite of the above mentioned problem. Interesting physically meaningful results obtained in this conceptual framework serve as an additional justification for it. We can point out calculations of one-loop effective actions in constant non-Abelian external fields [5, 8, 9] that were used for constructing the true QCD vacuum, see [9, 10, 13, 19]. One also ought to mention the description of phase-transitions in cosmological QCD [20], non-pertubative parton production from vacuum by a classical SU(3) [21] and SU(2) [22] chromoelectric field, boundary conditions and topological effects of the vacuum in the presence of a non-homogenous external magnetic field in the form of a flux tube [23, 24], and so on.

The key objects in nonperturbative (with respect to the background) QFT with a non-Abelian background are scalar and spinning particle propagators in the corresponding non-Abelian external

<sup>\*</sup>fresneda@fma.if.usp.br

<sup>&</sup>lt;sup>†</sup>gitman@dfn.if.usp.br

field. Exact solutions for such objects allow one to obtain by an integration one-loop results for various physical quantities. Moreover, path-integral representations for the propagators may be useful in obtaining exact solutions, which could then be used in calculations. Manifold path-integral representations for scalar and spinning particle propagators were constructed and calculated for various Abelian backgrounds in [25, 28, 29, 38, 39, 40, 41]. It turned out that such representations are also useful for deriving the so-called pseudoclassical actions for spinning particles, see [38, 39, 45. Some path-integral representations for propagators in non-Abelian backgrounds and problems related to the pseudoclassical description of isospin were studied in [47, 48, 28]. We recall that a classical theory for a Yang-Mills particle was first constructed from the classical limit of the Yang-Mills field equations by Wong [49]. Afterwards, Chen and Dresden [50] showed that the Yang-Mills field equations imply the equations of motion for a test particle with isotopic spin in a way similar as the Einstein equations imply the equations of a massive test particle. Casalbuoni et.al. [48] obtained a gauge-invariant Lagrangian description of scalar and spinning particles with isotopic spin, where Grassmann variables describe the internal degrees of freedom at the classical level, so that quantization gives finite-dimensional representations of the gauge group. Balachandran et.al. [47] applied Dirac quantization to a pseudoclassical Lagrangian formulation of scalar and spinning particles interacting with a non-Abelian gauge field, and additionally developed the method we use here to obtain the irreducible representations of isospin. In [40], the isospinor structure of the propagator of a scalar relativistic particle in the fundamental representation of SU(2) is derived from a path-integral representation using methods developed for the case of the spinning particle.

In the present article we return once again to these problems for the case of a scalar particle with isospin placed in various non-Abelian external backgrounds. We point out that a quantized scalar field in a non-Abelian background has been put forward as a tentative explanation of QCD confinement by means of a massive scalar particle (dilaton) [51], and also appears in the form of fundamental scalars coupled to gauge curvature terms in string theory [54].

We construct path-integral representations for the scalar particle propagator from two approaches: one is a generalization of the procedure proposed in [40] to any representation of SU(N) given in terms of antisymmetric generators, while the other is a constructed using fermionic coherent states valid for arbitrary representations of SU(N). The latter approach is a modification of the path-integral representation of the Dirac propagator by means of fermionic coherent states presented in [28]. In both cases we derive the pseudoclassical actions for a scalar particle in non-Abelian backgrounds, and quantize them to prove their consistency. In the Appendix, we put some technical details and proofs. The developed techniques can be easily generalized to the case of a spinning particle in non-Abelian and gravitational backgrounds. Such a generalization is the subject of our next publication.

### 2 Propagator representations

The causal propagator for a relativistic scalar particle interacting with a su(N) valued external field  $\mathcal{A}_{\mu}$  in Minkowski spacetime (in natural units  $\hbar = c = 1$ ) is described by the equation

$$\left(\mathcal{P}^2 - m^2\right)^{\alpha}_{\ \beta} D^{\beta}_{\ \gamma}\left(x, y\right) = -\delta^{\alpha}_{\gamma} \delta^4\left(x - y\right) , \ \mathcal{P}_{\mu} = i\partial_{\mu} - q\mathcal{A}_{\mu} \,, \tag{1}$$

where  $\mathcal{A}_{\mu} = A^{a}_{\mu} t^{\alpha}_{a\beta}$  is a linear combination of the traceless hermitian matrices  $t^{\alpha}_{a\beta}$ ,  $a = 1, ..., N^{2} - 1$ which are the generators of the Lie algebra su(N) in an  $n \times n$  irreducible matrix representation whose indices are labeled by greek letters from the beginning of the alphabet,  $\alpha, \beta, \gamma, \text{etc.}$ ,  $\alpha = 1, ..., N$ . Since SU(N) is a compact group, there is a basis where its structure constants are totally antisymmetric and purely imaginary,

$$[t_a, t_b] = f_{ab}^c t_c, \ f_{ab}^c \equiv f_{[abc]}, \tag{2}$$

and the generators can be normalized as  $\operatorname{tr}(t_a t_b) = 1/2\delta_{ab}$ .

In the following we will consider two different realizations of the Lie algebra (2) of su(N). The first realization will be in terms of creation and annihilation operators defined on a suitable Fock space, and the second realization will be in terms of the generators of a suitable Clifford algebra.

I. Let us consider the first realization. Consider an abstract Hilbert space  $\mathcal{H}$  which is the direct product of the usual representation space for the Heisenberg algebra, whose basis vectors are denoted as  $|x\rangle$ ,

$$\hat{x}^{\mu} |x\rangle = x^{\mu} |x\rangle , \ \langle x |y\rangle = \delta^{4} (x - y) , \ \int d^{4}x |x\rangle \langle x| = I ,$$
$$[\hat{x}^{\mu}, \hat{p}_{\nu}] = i\delta^{\mu}_{\nu} , \ \langle x| \, \hat{p}_{\mu} \, |y\rangle = -i\partial_{\mu}\delta^{4} (x - y) , \qquad (3)$$

and an abstract Hilbert space V which we do not specify for the time being, but whose orthonormal basis vectors are  $|\alpha\rangle$ ,  $\alpha = 1, ..., n$ ,

$$\langle \alpha | \beta \rangle = \delta_{\alpha\beta} , \sum_{\alpha=1}^{n} | \alpha \rangle \langle \alpha | = I .$$
 (4)

Thus, the abstract Hilbert space  $\mathcal{H} = H \otimes V$  has the orthonormal basis  $|x, \alpha\rangle = |x\rangle \otimes |\alpha\rangle$ ,  $\langle x, \alpha | y, \beta \rangle = \delta^4 (x - y) \delta_{\alpha\beta}$ .

Next, we interpret the matrix operators appearing in (1), as matrix elements of operators in  $\mathcal{H}$ . With this in mind, the propagator  $D^{\alpha}_{\beta}(x,y)$  is the matrix element of an abstract operator  $\hat{D}$ ,

$$D(x,y)^{\alpha}_{\ \beta} = \langle x, \alpha | \, \hat{D} \, | y, \beta \rangle , \qquad (5)$$

and the generators  $t^{\alpha}_{a\beta}$  are matrix elements of the operators  $\hat{t}_a$ ,

$$\langle \alpha | \hat{t}_a | \beta \rangle = t^{\alpha}_{a\beta}$$

We note that if the matrix elements of the operators  $\hat{t}_a$  are generators of a representation the algebra su(N), then so are the operators themselves:

$$\left[t_a, t_b\right]^{\alpha}_{\ \beta} = f^c_{ab} t^{\alpha}_{c\beta} \Leftrightarrow \left[\hat{t}_a, \hat{t}_b\right] = f^c_{ab} \hat{t}_c \,. \tag{6}$$

Using the operators just defined, one can write (1) in operator form,

$$\left(\hat{P}^2 - m^2\right)\hat{D} = -I\,,$$

where

$$\hat{P}_{\mu} = -\hat{p}_{\mu} - q\hat{\mathcal{A}}_{\mu} , \ \hat{\mathcal{A}}_{\mu} = A^{a}_{\mu}\left(\hat{x}\right)\hat{t}_{a} , \ \langle x,\alpha | \ \hat{P}_{\mu} | y,\beta \rangle = \left(i\partial_{\mu}\delta_{\alpha\beta} - qA_{\mu}\left(x\right)t^{\alpha}_{a\beta}\right)\delta^{4}\left(x-y\right) .$$

Thus, one can formally write the inverse of the operator  $\hat{D}$ ,

$$\hat{D} = -\left(\hat{P}^2 - m^2 + i\varepsilon\right)^{-1},\,$$

by means of the proper time representation

$$\hat{D} = i \int_0^\infty d\lambda e^{-i\hat{H}(\lambda)} , \ \hat{H} = -\lambda \left( \hat{P}^2 - m^2 + i\varepsilon \right) .$$
<sup>(7)</sup>

Let us now further specify  $\mathcal{H}$  by defining V as the one-particle sector of the Fock space for the fermionic creation and annihilation operators  $a^{\dagger}$  and a,

$$\hat{a}_{\alpha} \left| 0 \right\rangle = 0, \ \hat{a}_{\alpha}^{\dagger} \left| 0 \right\rangle = \left| \alpha \right\rangle$$

which satisfy the algebra

$$\left[\hat{a}^{\dagger}_{\alpha},\hat{a}_{\beta}\right]_{+} = \delta_{\alpha\beta}, \ \left[\hat{a}^{\dagger}_{\alpha},\hat{a}^{\dagger}_{\beta}\right]_{+} = \left[\hat{a}_{\alpha},\hat{a}_{\beta}\right]_{+} = 0.$$
(8)

Then it is possible to represent the operators  $\hat{t}_a$  as

$$\hat{t}_a = \hat{a}^{\dagger}_{\alpha} t^{\alpha}_{a\beta} \hat{a}_{\beta} , \ t^{\alpha}_{a\beta} = \langle \alpha | \, \hat{t}_a \, | \beta \rangle \ . \tag{9}$$

Here it is important to observe that  $\hat{t}_a$  are generators of a representation of su(N),

$$\left[\hat{t}_a, \hat{t}_b\right] = f_{ab}^c \hat{t}_c,$$

since their matrix elements  $t^{\alpha}_{a\beta}$  satisfy the su(N) commutation relations (6). In addition, tracelessness and hermiticity of  $t^{\alpha}_{a\beta}$  imply the same for the operators  $\hat{t}_a$ ,

$$\begin{aligned} \mathrm{tr} \hat{t}_a &\equiv \sum_{\alpha=1}^M \left< \alpha \right| \hat{t}_a \left| \alpha \right> = t^{\alpha}_{a\alpha} = 0 \\ \hat{t}^{\dagger}_a &= \left( \hat{a}^{\dagger}_{\alpha} t^{\alpha}_{a\beta} \hat{a}_{\beta} \right)^{\dagger} = \hat{a}^{\dagger}_{\beta} \bar{t}^{\alpha}_{a\beta} \hat{a}_{\alpha} = \hat{t}_a \,, \end{aligned}$$

where the  $\dagger$ -involution of the abstract operator algebra complex-conjugates the matrix entries of  $t^{\alpha}_{a\beta}$ in the above. Finally, we note that  $\hat{t}_a$  conserves the number of particles. Using the representation (9) for the generators  $t_a$  and following [55, 28], we now introduce coherent states  $|\chi\rangle$  and  $\langle \bar{\chi}|$  defined by the exponential of the fermion operators  $\hat{a}$  and  $\hat{a}^{\dagger}$  acting on the vacuum:

$$|\chi\rangle = D(\chi)|0\rangle , \ \langle \bar{\chi}| = |\chi\rangle^{\dagger} , \ D(\chi) = e^{\hat{a}^{\dagger}\chi - \hat{a}\bar{\chi}} , \ [\hat{a}_{\alpha}, D(\chi)]_{-} = \chi_{\alpha}D(\chi) ,$$

where  $\chi_{\alpha}$  and  $\bar{\chi}_{\alpha} = \chi_{\alpha}^{\dagger}$  are Grassmann numbers that commute with the vacuum state. Consequently, these states satisfy

$$\begin{aligned} \hat{a}^{\alpha} |\chi\rangle &= \chi^{\alpha} |\chi\rangle \ , \ \langle \bar{\chi} | \ \hat{a}^{\dagger}_{\alpha} &= \langle \bar{\chi} | \ \bar{\chi}_{\alpha} \ , \\ \langle \bar{\chi} | \ \xi\rangle &= e^{\frac{1}{2} \left( \chi \bar{\chi} + \xi \bar{\xi} - 2\xi \bar{\chi} \right)} \ , \ \int \prod_{\alpha=1}^{N} d\bar{\chi}_{\alpha} d\chi^{\alpha} |\chi\rangle \ \langle \bar{\chi} | = \hat{1}_{V} \ , \ \int d\chi \chi = \int d\bar{\chi} \bar{\chi} = 1 \end{aligned}$$

Using the above identity resolutions, it is possible to relate matrix elements from the one-particle sector fock-space basis  $|\alpha\rangle$  to the coherent basis  $|\chi\rangle$ ,

$$\langle \alpha | \cdot | \beta \rangle = \int \prod_{\sigma,\kappa=1}^{N} d\bar{\chi}'_{\sigma} d\chi'^{\sigma} d\bar{\chi}_{\kappa} d\chi^{\kappa} e^{\frac{1}{2} \left( \chi' \bar{\chi}' + \chi \bar{\chi} \right)} \chi'^{\alpha} \langle \bar{\chi}' | \cdot | \chi \rangle \bar{\chi}_{\beta} , \qquad (10)$$

where we have used  $\langle \bar{\chi} | \alpha \rangle = \bar{\chi}_{\alpha} \exp \frac{1}{2} \chi \bar{\chi}$ . As a consequence, we are able to recast the original form of the propagator (5), as matrix elements of one-particle Fock states, in terms of matrix elements of the coherent states,

$$D(x,y)^{\alpha}_{\ \beta} = \int \prod_{\sigma,\kappa=1}^{N} d\bar{\chi}'_{\sigma} d\chi'^{\sigma} d\bar{\chi}_{\kappa} d\chi^{\kappa} e^{\frac{1}{2} \left(\chi' \bar{\chi}' + \chi \bar{\chi}\right)} \chi'^{\alpha} \langle x, \bar{\chi}' | \hat{D} | y, \chi \rangle \bar{\chi}_{\beta} .$$
(11)

In the next section, the matrix elements  $\langle x, \bar{\chi}' | \hat{D} | y, \chi \rangle$  will be used to to obtain a path-integral representation for the propagator.

II. Another possible interpretation of the propagator D(x, y) appearing in (1) can be simply as the matrix elements

$$D(x,y)_{j}^{i} = \langle x | \hat{D}_{j}^{i} | y \rangle$$

of the basis elements  $|x\rangle$  of the abstract Hilbert space H. The abstract operator  $\hat{D}$  acquires indices directly from the matrices of the generators of su(N). Notice we have relabeled the indices of the matrix representation of su(N). The new indices i and j denote the matrix entries of a new set of generators  $T_a$ ,

$$T_a = \frac{1}{4} \Gamma_{\alpha} t^{\alpha}_{a\beta} \Gamma_{\beta} , \ [\Gamma_{\alpha}, \Gamma_{\beta}] = 2\delta_{\alpha\beta} .$$
<sup>(12)</sup>

These generators are very convenient for obtaining path-integral representations of the propagator using techniques adapted from the spinning particle case. However, for  $T_a$  satisfying (2), this is a representation only if the matrices  $t_a$  are antisymmetric,  $t_a^T = -t_a$ . This drawback can be circumvented if we take the  $t_a$  matrices in the adjoint representation  $t_{ab}^c = f_{ab}^c$ . Besides, there are be situations where it is possible to choose antisymmetric  $t_a$  for different irreducible representations. For instance, in the case of SU(2), it is always possible to choose antisymmetric  $t_a$  for the integer spin *s* representations. In this case,  $\alpha, \beta = 1, ..., 2s + 1$  and  $i, j = 1, ..., 2^s$ . In the general case, in the adjoint representation,  $\alpha, \beta = 1, ..., N^2 - 1$  and thus  $i, j = 1, ..., 2^{\left[ (N^2 - 1)/2 \right]}$ . In the familiar case of the adjoint representation of su(2), one has

$$T_i = \frac{i}{4} \varepsilon_{ijk} \Gamma_k \Gamma_j = -\frac{i}{4} \varepsilon_{ijk} \Gamma_j \Gamma_k \,, \ i, j, k = 1, 2, 3 \,,$$

where the  $\Gamma$ 's satisfy  $[\Gamma_i, \Gamma_j]_+ = 2\delta_{ij}$  and are order 2 matrices, so they can be chosen to be the Pauli matrices,  $\Gamma_i = \sigma_i^{-1}$ ,

$$T_i = -\frac{i}{4}\varepsilon_{ijk}\sigma_j\sigma_k = \frac{1}{2}\sigma_i.$$
(13)

The generators  $T_i$  are hermitian and traceless, and they satisfy the su(2) algebra

$$[T_i, T_j] = i\varepsilon_{ijk}T_k$$

This case is special, because the choice of the adjoint representation for  $t_a$  gives  $T_a$  in the fundamental representation. Another special situation occurs with SU(4), where we can choose  $t_a$  to be antisymmetric matrices of order 6, since  $su(4) \simeq so(6)$ . Thus one has even or odd spinors of so(6)with 4 components, giving by means of a method described in sections 3.1 and 4.1, the fundamental representation of SU(4).

$$^{1}\sigma_{1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \ \sigma_{2} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \ \sigma_{3} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

# 3 Path integral in coherent states representation

### 3.1 Path integral

Our goal in this section is to write a path-integral representation for

$$D_{\chi}(x,\bar{\chi}';y,\chi) \equiv \langle x,\bar{\chi}'|\,\hat{D}|y,\chi\rangle = i \int_{0}^{\infty} d\lambda \,\langle x,\bar{\chi}'|\,e^{-i\hat{H}(\lambda)}\,|y,\chi\rangle \tag{14}$$

We insert N-1 identity resolutions  $I = \int dx d\bar{\chi} d\chi |x, \chi\rangle \langle x, \bar{\chi}|$  and N integration over  $\lambda$ :

$$D_{\chi}(x,\bar{\chi}';y,\chi) = \lim_{N \to \infty} i \int_{0}^{\infty} d\lambda_{0} \int \left(\prod_{k=1}^{N-1} dx_{k} d\bar{\chi}_{k} d\chi_{k}\right) d\lambda_{1} \cdots d\lambda_{N}$$
$$\prod_{k=1}^{N} \langle x_{k}, \bar{\chi}_{k} | e^{-i\hat{H}(\lambda_{k})/N} | x_{k-1}, \chi_{k-1} \rangle \,\delta\left(\lambda_{k} - \lambda_{k-1}\right) \,, \tag{15}$$

where  $x_N = x$ ,  $\bar{\chi}_N = \bar{\chi}'$ ,  $x_0 = y$  and  $\chi_0 = \chi$ . In order to evaluate the general matrix element appearing in (15), one must choose a definite ordering prescription for the operators in  $\hat{H}$ . In particular, one must solve the ordering ambiguity of the four-fermion term in  $\hat{P}^2$ . In [28], an additional identity resolution is inserted between the  $\hat{P}$  operators as a solution to the ordering problem. We do not know to which ordering prescription this corresponds, and conventional ordering prescriptions such as Weyl ordering and normal ordering are not gauge-invariant. In the sequel we show that Weyl ordering is not gauge-invariant, and compute the resulting effective action. As shown in the Appendix (39), the Hamiltonian operator differs from the Weyl-ordered<sup>2</sup> expression by the term  $\lambda \frac{q^2}{4} \operatorname{tr}(t_a t_b) \hat{A}^{\mu}_{\mu} \hat{A}^{\mu b}$ . This action, apart from the gauge-breaking term, is identical to the one that would be obtained by doubling the time partition.

Applying the midpoint rule (41) for the general matrix elements gives

$$\begin{split} \langle x_k, \bar{\chi}_k | \, \hat{H}\left(\lambda_k\right) | x_{k-1}, \chi_{k-1} \rangle &= \int \frac{dp_k}{(2\pi)^4} d\bar{\eta}_k d\eta_k \left\langle x_k, \bar{\chi}_k | \, p_k, \eta_k \right\rangle \left( H_W\left(\lambda_k\right) + Q\left(\lambda_k\right) \right) \left\langle p_k, \bar{\eta}_k | \, x_{k-1}, \chi_{k-1} \right\rangle \,, \\ H_W\left(\lambda_k\right) &\equiv H_W\left(\lambda_k, \frac{x_k + x_{k-1}}{2}, p_k, \bar{\eta}_k, \frac{\eta_k + \chi_{k-1}}{2}\right) \,, \\ Q\left(\lambda_k\right) &\equiv \lambda_k \frac{q^2}{4} \mathrm{tr}\left(t_a t_b\right) A_\mu^a \left(\frac{x_k + x_{k-1}}{2}\right) A^{b\mu} \left(\frac{x_k + x_{k-1}}{2}\right) \,, \end{split}$$

where  $H_W$  is the Weyl-symbol of  $\hat{H}_W$ . Substituting the delta functions  $\delta(\lambda_k - \lambda_{k-1})$  by their integral representations and using the integral representations of the fermionic delta (43) for the  $\chi$  and  $\bar{\chi}$  integrations, we have

$$D_{\chi}\left(x,\bar{\chi}';y,\chi\right) = \lim_{N\to\infty} i\int_{0}^{\infty} d\lambda_{0} \int \left(\prod_{k=1}^{N-1} dx_{k}\right) \left(\prod_{k=1}^{N} \frac{dp_{k}}{(2\pi)^{4}} d\lambda_{k} \frac{d\pi_{k}}{(2\pi)} d\bar{\eta}_{k} d\eta_{k}\right) \exp\frac{1}{2} \left(\chi'\bar{\chi}' - \eta_{N}\bar{\eta}_{N} + 2\bar{\chi}'\eta_{N}\right)$$
$$\exp i\sum_{k=1}^{N} \left\{ p_{k} \frac{(x_{k} - x_{k-1})}{\Delta t} + \pi_{k} \frac{(\lambda_{k} - \lambda_{k-1})}{\Delta t} - \frac{i}{2} \frac{(\eta_{k} - \eta_{k-1})}{\Delta t} \bar{\eta}_{k} - \frac{i}{2} \frac{(\bar{\eta}_{k} - \bar{\eta}_{k-1})}{\Delta t} \eta_{k-1} - H_{W}\left(\lambda_{k}\right) - Q\left(\lambda_{k}\right) \right\} \Delta t ,$$

 $^{2}$ Weyl ordering here means total symmetrization in bosonic degrees of freedom, and total antisymmetrization in fermionic degrees of freedom.

where  $H_W(\lambda_k) = H_W\left(\lambda_k, \frac{x_k + x_{k-1}}{2}, p_k, \bar{\eta}_k, \frac{\eta_k + \eta_{k-1}}{2}\right), \eta_0 = \chi$ . The term  $\chi' \bar{\chi}' - \eta_N \bar{\eta}_N + 2\bar{\chi}' \eta_N$  comes from  $\langle \bar{\chi}' | \eta_N \rangle$ , and in the limit  $N \to \infty$  will reduce to  $2\bar{\chi}' \eta$  (1). Taking the limit  $N \to \infty$  ( $\Delta t \to 0$ ) and renaming  $\eta \to \chi$  and  $\bar{\eta} \to \bar{\chi}$ , one has

$$D_{\chi}\left(x,\bar{\chi}';y,\chi\right) = i\int_{0}^{\infty}d\lambda_{0}\int DxDpD\lambda D\pi D\bar{\chi}D\chi\exp iS_{eff}\exp\bar{\chi}\left(1\right)\chi\left(1\right),$$
$$S_{eff} = \int_{0}^{1}dt\left(p\dot{x}+\pi\dot{\lambda}+\frac{i}{2}\left(\bar{\chi}\dot{\chi}-\dot{\chi}\chi\right)+\lambda\left(\left(p_{\mu}+qA_{\mu}^{a}I_{a}\right)^{2}-m^{2}\right)-\frac{q^{2}}{4}\lambda\mathrm{tr}\left(t_{a}t_{b}\right)A_{\mu}^{a}A^{b\mu}\right), \quad (16)$$

where  $I_a = \bar{\chi} t_a \chi$ , and the functional integration is performed over the paths  $x^{\mu}(t)$ ,  $p_{\mu}(t)$ ,  $\lambda(t)$ ,  $\pi(t)$ ,  $\bar{\chi}(t)$  and  $\chi(t)$ , with boundary values  $x^{\mu}(0) = y^{\mu}$ ,  $x^{\mu}(1) = x^{\mu}$ ,  $\lambda(0) = \lambda_0$ ,  $\bar{\chi}(1) = \bar{\chi}'$  and  $\chi(0) = \chi$ .

Since the path integral is translation-invariant, one can integrate over the momenta  $p_{\mu}$  by shifting  $p \mapsto p + \tilde{p}$ , where  $\tilde{p} = -\dot{x}/2\lambda - qA^aI_a$  is the solution to the classical equation  $\dot{x} = \partial H_{eff}/\partial p$ . One finds after making the substitution  $2\lambda = e$ , the Lagrangian form of the path integral:

$$D_{\chi}(x,\bar{\chi}';y,\chi) = i \int_{0}^{\infty} d\lambda_{0} \int Dx De D\pi D\bar{\chi} D\chi M[e,x] \exp i \left(S_{eff} + S_{G}\right) \exp\left(\bar{\chi}\left(1\right)\chi\left(1\right)\right) ,$$
$$S_{eff} = \int_{0}^{1} dt \left(-\frac{\dot{x}^{2}}{2e} - \frac{e}{2}m^{2} - q\dot{x}^{\mu}A_{\mu}^{a}I_{a} + \frac{i}{2}\left(\bar{\chi}\dot{\chi} - \dot{\chi}\chi\right)\right) , \tag{17}$$

with the Lagrangian measure and reparametrization gauge-fixing term  $S_G$ 

$$M[e,x] = \int Dp \exp\frac{i}{2} \int_0^1 e\left(p^2 - \frac{q^2}{4} \operatorname{tr} t_a t_b A^a_\mu A^{\mu b}\right) dt$$
(18)

$$S_G = \int_0^1 \pi \dot{e} d\tau \tag{19}$$

Thus, the path-integral representation for the propagator can be derived with an unambiguous ordering prescription (Weyl-ordering) at the cost of defining a gauge non-invariant measure.

#### **3.2** Pseudoclassical action

The action functional  $S_{eff}$  in (17),

$$S_{eff} = \int_0^1 dt \left( -\frac{\dot{x}^2}{2e} - \frac{e}{2} m^2 - q \dot{x}^{\mu} A^a_{\mu} I_a + \frac{i}{2} \left( \bar{\chi} \dot{\chi} - \dot{\bar{\chi}} \chi \right) \right) , \ I_a = \bar{\chi} t_a \chi \,, \tag{20}$$

is reparametrization invariant,

$$\delta_{\epsilon}S_{eff} = 0 , \ \delta_{\epsilon}x = \epsilon \dot{x} , \ \delta_{\epsilon}e = \frac{d}{dt} \left(\epsilon e\right) , \ \delta_{\epsilon}\chi = \epsilon \dot{\chi} , \ \delta_{\epsilon}\bar{\chi} = \epsilon \dot{\bar{\chi}} .$$
<sup>(21)</sup>

In the gauge  $e = \sqrt{\dot{x}^2}/m$  it coincides with the action given in [48, 47] describing a scalar relativistic particle with anticommuting coordinates in a representation of a symmetry group G, whose equations of motion are

$$m\frac{d}{dt}\frac{\dot{x}_{\mu}}{\sqrt{\dot{x}^2}} = q\dot{x}^{\nu}F^a_{\mu\nu}I_a , \ D_t\chi^{\alpha} \equiv \frac{d}{dt}\chi^{\alpha} + iq\dot{x}^{\mu}A^a_{\mu}t^{\alpha}_{a\beta}\chi^{\beta} = 0 ,$$
 (22)

where  $F^a_{\mu\nu} = \partial_{\mu}A^a_{\nu} - \partial_{\nu}A^a_{\mu} + iqf^a_{bc}A^b_{\mu}A^c_{\nu}$  is the field-strength and  $D_t$  is the covariant derivative. For canonical analysis purposes<sup>3</sup>, however, it is better to start from the reparametrization invari-

For canonical analysis purposes", however, it is better to start from the reparametrization invariant action (20). Since this action does not contain derivatives of the einbein, it is best to consider it as a velocity (see [57]), and not introduce its conjugate momentum. One thus arrives at the following Hamiltonian,

$$H = -\frac{e}{2}T - \dot{\chi}_{\alpha}\phi_{\alpha} - \dot{\bar{\chi}}_{\alpha}\bar{\phi}_{\alpha} \,,$$

where the set of constraints  $\Phi = \{T, \phi, \bar{\phi}\},\$ 

$$T = (p_{\mu} + qA_{\mu}^{a}I_{a})^{2} - m^{2}, \ \phi_{\alpha} = \pi_{\alpha} - \frac{i}{2}\bar{\chi}_{\alpha}, \ \bar{\phi}_{\alpha} = \bar{\pi}_{\alpha} - \frac{i}{2}\chi_{\alpha},$$

defines a degenerate supermatrix  $\{\Phi, \Phi\}$ . The constraint algebra is simplified if we consider an equivalent set of constraints  $\{\tilde{T}, \phi, \bar{\phi}\}$ , where  $\tilde{T}$  is obtained from T through the shifts  $\chi \to \chi - i\bar{\phi}$  and  $\bar{\chi} \to \bar{\chi} - i\phi$ ,

$$\left\{\tilde{T},\phi_{\alpha}\right\} = \left\{\tilde{T},\bar{\phi}_{\alpha}\right\} = 0, \ \left\{\phi_{\alpha},\bar{\phi}_{\beta}\right\} = -i\delta_{\alpha\beta}.$$

The new Hamiltonian with redefined Lagrange multipliers is

$$\tilde{H} = \Lambda \tilde{T} + \Lambda_{\alpha} \phi_{\alpha} + \bar{\Lambda}_{\alpha} \bar{\phi} \,,$$

giving the following time-evolution for the constraints,

$$\frac{d}{dt}\tilde{T} = 0 , \ \frac{d}{dt}\phi_{\alpha} = i\bar{\Lambda}_{\alpha} , \ \frac{d}{dt}\bar{\phi}_{\alpha} = i\Lambda_{\alpha} ,$$

so the condition of conservations of the constraints in time simply determines  $\Lambda$  and  $\bar{\Lambda}$ . The equations of motion for the independent variables  $\eta = (x^{\mu}, p_{\mu}, \chi_{\alpha}, \bar{\chi}_{\alpha})$  are given by

$$\dot{\eta} = \left\{\eta, \Lambda \tilde{T}\right\}_{D(\phi)}, \ \phi_{\alpha} = \bar{\phi}_{\alpha} = \tilde{T} = 0,$$

where the Dirac brackets have been constructed with regard to the second-class constraint set  $\{\phi; \bar{\phi}\}$ . Using well known properties of the Dirac brackets, the equations of motion become

$$\dot{\eta} = \{\eta, \Lambda T\}_{D(\phi)}, \ \phi_{\alpha} = \bar{\phi}_{\alpha} = T = 0,$$

And the nonzero brackets between independent variables are

$$\{x^{\mu}, p_{\nu}\}_{D(\phi)} = \delta^{\mu}_{\nu}, \ \{\chi_{\alpha}, \bar{\chi}_{\beta}\}_{D(\phi)} = -i\delta_{\alpha\beta}.$$
 (23)

Moreover, the  $I_a$  are covariantly constant generators of SU(N),

$$\{I_a, I_b\}_{D(\phi)} = -if_{ab}^c I_c , \ D_t I_a \equiv \frac{d}{dt} I_a + iq\dot{x}^{\mu} A^b_{\mu} f^c_{ab} I_c = 0 ,$$
(24)

hence are called isospin.

<sup>&</sup>lt;sup>3</sup>Definitions and conventions are those used in [56].

From (23), we see that the Grassmann operators will generate a creation-annihilation operator algebra,

$$\chi_{\alpha} \to a_{\alpha} , \ \bar{\chi}_{\alpha} \to a_{\alpha}^{\dagger} , \ \left[ a_{\alpha}, a_{\beta}^{\dagger} \right]_{+} = \delta_{\alpha\beta} .$$
 (25)

The Hilbert space  $\mathcal{H}$  can be realized as the direct product of a representation space for the Heisenberg algebra and the  $2^n$ -dimensional Fock space of the creation and annihilation operators,

$$|x;\alpha_1\cdots\alpha_p\rangle = a^{\dagger}_{\alpha_1}\cdots a^{\dagger}_{\alpha_p} |x;0\rangle \in \mathcal{H}, \ p = 0,..,n.$$
<sup>(26)</sup>

As is well-known, the group SO(2n) preserves the commutation relations (25), and the so(2n) generators in the above representation are given by  $c_{\alpha\beta} = \left[a_{\alpha}, a_{\beta}^{\dagger}\right]/2$ ,  $a_{\alpha}a_{\beta}$  and  $a_{\alpha}^{\dagger}a_{\beta}^{\dagger}$ . The  $c_{\alpha\beta}$  belong to the u(n) subalgebra of so(2n). The *n* operators  $c_{\alpha\beta}$  for  $\alpha = \beta$  form the Cartan subalgebra of so(2n).

The representation (26) is a  $2^n$ -dimensional spinor representation of so(2n), and its irreducible representations are given by states with an even or odd number of creation operators, corresponding to the  $2^{n-1}$ -dimensional Weyl (semi-spinor) representations of so(2n). These states can be further decomposed in irreducible representations of su(N), since the isospin generators  $\hat{t}_a$  are a linear combination of the so(2n) generators,

$$\hat{t}_a = t_{a\alpha\beta} a^{\dagger}_{\alpha} a_{\beta} = t_{a\alpha\beta} \left( -2c_{\beta\alpha} + \delta_{\alpha\beta} \right) = -2t_{a\alpha\beta}c_{\beta\alpha} \,.$$

Therefore, we see that the  $\hat{t}_a$  generate a su(N) subalgebra of so(2n).

In general, in order to determine the SU(N) content of the wave function, one proceeds as in [47]: given  $t_a$  an irreducible representation of su(N) in terms of  $n \times n$  matrices, the wave function belongs to a (Weyl) semi-spinor representation of so(2n). Then, one decomposes the set of Cartan generators of su(N) (a maximal set of commuting generators) in terms of the nCartan generators of so(2n). In the special case of the representation (26), one can choose the operators  $c_{\alpha} = [a_{\alpha}, a_{\alpha}^{\dagger}]/2$ ,  $\alpha = 1, ..., n$  as the maximum set of commuting generators of so(2n). For instance, in the case of SU(2) one can take the isospin projection  $I_1$  and for SU(3) one can take the isospin projection  $I_1$  and the hypercharge Y to characterize irreducible representations. One then decomposes isospin generators in terms of the  $c_{\alpha}$  to obtain their eigenvalues for the spinor representation of so(2n)to which the wave function belongs. The range of these eigenvalues gives the irreducible representations of SU(N). Therefore, to each given n-dimensional irreducible representation  $t_a$  of SU(N), the wave function will belong to a  $2^{n-1}$ -dimensional representation of SO(2n) (a semi-spinor representation), which decomposes into irreducible representations of SU(N) as determined by the isospin generators  $\hat{t}_a$ .

In the special case of SU(2), since it is of rank 1, the Cartan subalgebra is generated by a single element, say  $t_3$ , whose matrix representation in a basis of isospin s eigenstates is of the form  $t_3 = \text{diag}(s, s - 1, ..., -s + 1, -s)$ . The decomposition in Cartan generators of so(4s + 2) of the isospin operator  $\hat{t}_3$  is as follows,

$$\hat{t}_3 = sc_1 + (s-1)c_2 + \dots + (-s)c_{2s+1}$$
.

Each  $c_{\alpha}$  can take either of the values plus or minus 1/2. However, the wave function is in a state of either an even number of plus +1/2 (even Weyl spinor) or an odd number of +1/2 (odd Weyl spinor). For instance, for s = 1/2, the possible eigenvalues of  $t_3$  for the even representation is twice 0, giving two scalar representations; and for the odd representation is  $\pm 1/2$ , giving the isospin 1/2

representation. For integer spin, even and odd representations decompose in the same way, and the largest representation is of spin (s + 1) s/2. For example, s = 1 gives the eigenvalues 1, 0, -1 and again 0, giving the isospin 1 representation plus a scalar. We summarize the results for some values of isospin in the table below,

isospin	symmetry group	representation dimension	decomposition(even; odd)
0	$SO\left(2 ight)$	1	<u>0</u>
1/2	$SO\left(4 ight)$	2	$2 \times \underline{0}; \underline{\frac{1}{2}}$
1	$SO\left(6 ight)$	4	$\underline{0} + \underline{1}$
3/2	$SO\left(8 ight)$	8	$3 \times \underline{0} + \underline{2}; 2 \times \underline{3}$
2	SO(10)	16	$\underline{0} + \underline{1} + \underline{2} + \underline{3}$

Thus, in order to obtain the fundamental representation of SU(2) upon quantization, one must choose the Hilbert space to be the odd Weyl spinor representation of SO(4) of two-component spinors. In this case, one gets from the constraint T the Dirac quantization condition

$$\hat{T}\phi = \left[ \left( \hat{p}_{\mu} + qA_{\mu}^{a}t_{a} \right)^{2} - m^{2} \right] \phi \left( x \right) = 0, \qquad (27)$$

which is precisely the wave equation in (1) for  $t_a = \frac{1}{2}\sigma_a$ .

It also possible to arrive at these results starting from the classical action (20). In the following, it will be convenient to express the Grassmann variables  $\chi$  in terms of their real and imaginary parts,

$$\chi_{\alpha} = \frac{1}{\sqrt{2}} \left( \chi_{1\alpha} + i \chi_{2\alpha} \right) \,,$$

so that we are left with the real variables

$$\chi_{1\alpha} = \frac{1}{\sqrt{2}} \left( \chi_{\alpha} + \bar{\chi}_{\alpha} \right) , \ \chi_{2\alpha} = \frac{1}{i\sqrt{2}} \left( \chi_{\alpha} - \bar{\chi}_{\alpha} \right) .$$
(28)

In this way, the Grassmanian kinetic term becomes

$$L_{kin} = \frac{i}{4} \left( \chi_1 \dot{\chi}_1 - \dot{\chi}_1 \chi_1 + \chi_2 \dot{\chi}_2 - \dot{\chi}_2 \chi_2 \right) \,.$$

 $L_{kin}$  is invariant under transformations induced by  $R_{\alpha\beta} = -i(\chi_{1\alpha}\chi_{1\beta} + \chi_{2\alpha}\chi_{2\beta})$  and  $S_{\alpha\beta} = -i(\chi_{1\alpha}\chi_{2\beta} + \chi_{1\beta}\chi_{2\alpha})$ 

$$\delta_{\omega}\chi_{i\alpha} \equiv \left\{\frac{1}{2}\omega_{\beta\gamma}R_{\beta\gamma},\chi_{i\alpha}\right\}_{D(\phi)} = \omega_{\alpha\beta}\chi_{i\beta},$$
$$\delta_{\lambda}\chi_{i\alpha} \equiv \left\{\frac{1}{2}\lambda_{\beta\gamma}S_{\beta\gamma},\chi_{i\alpha}\right\}_{D(\phi)} = (-1)^{i+1}\lambda_{\alpha\beta}\chi_{i\beta}$$

where the Dirac brackets for the real variables follows from the old variables' brackets (23) and their expression in terms of the real variables (28),

$$\{\chi_{1\alpha}, \chi_{1\beta}\}_{D(\phi)} = \{\chi_{2\alpha}, \chi_{2\beta}\}_{D(\phi)} = -i\delta_{\alpha\beta}, \ \{\chi_{1\alpha}, \chi_{2\beta}\}_{D(\phi)} = 0.$$

The symmetry generators  $R_{\alpha\beta}$  and  $S_{\alpha\beta}$  satisfy the Lie algebra

$$\begin{split} \{R_{\alpha\beta}, R_{\gamma\delta}\}_{D(\phi)} &= \delta_{\alpha\gamma}R_{\beta\delta} + \delta_{\beta\delta}R_{\alpha\gamma} - \delta_{\alpha\delta}R_{\beta\gamma} - \delta_{\beta\gamma}R_{\alpha\delta} \,, \\ \{S_{\alpha\beta}, S_{\gamma\delta}\}_{D(\phi)} &= \delta_{\alpha\gamma}R_{\beta\delta} + \delta_{\beta\delta}R_{\alpha\gamma} + \delta_{\alpha\delta}R_{\beta\gamma} + \delta_{\beta\gamma}R_{\alpha\delta} \,, \\ \{R_{\alpha\beta}, S_{\gamma\delta}\}_{D(\phi)} &= \delta_{\alpha\gamma}S_{\beta\delta} - \delta_{\beta\delta}S_{\alpha\gamma} + \delta_{\alpha\delta}S_{\beta\gamma} - \delta_{\beta\gamma}S_{\alpha\delta} \,. \end{split}$$

Above we recognize the commutation relations of the combination of the o(2n) generators  $L_{ij}$ , i, j = 1, ..., 2n,

$$R_{\alpha\beta} = L_{2\alpha-1,2\beta-1} + L_{2\alpha,2\beta} \,, \ S_{\alpha\beta} = L_{2\alpha,2\beta-1} - L_{2\alpha-1,2\beta} - \delta_{\alpha\beta} \,.$$

Moreover, from the following decomposition of the generators  $I_a$  in terms of the symmetric and antisymmetric part of  $t_a$ ,

$$I_{a} = t_{a(\alpha\beta)} \left( \bar{\chi}_{\alpha} \chi_{\beta} + \bar{\chi}_{\beta} \chi_{\alpha} \right) + t_{a[\alpha\beta]} \left( \bar{\chi}_{\alpha} \chi_{\beta} - \bar{\chi}_{\beta} \chi_{\alpha} \right) = \frac{i}{2} t_{a(\alpha\beta)} \left( \chi_{1\alpha} \chi_{2\beta} + \chi_{1\beta} \chi_{2\alpha} \right) + \frac{1}{2} t_{a[\alpha\beta]} \left( \chi_{1\alpha} \chi_{1\beta} + \chi_{2\alpha} \chi_{2\beta} \right) , = -\frac{1}{2} t_{a(\alpha\beta)} S_{\alpha\beta} + \frac{i}{2} t_{a[\alpha\beta]} R_{\alpha\beta}$$

$$(29)$$

we again find the  $I_a$  are a linear combination of  $R_{\alpha\beta}$  and  $S_{\alpha\beta}$ , which is to say that the  $I_a$  are the generators of the subalgebra su(N) of so(2n). It is so(2n) and not o(2n), because the trace part of  $S_{\alpha\beta}$  in the expansion of  $I_a$  gives no contribution, since the  $t_a$  are traceless.

# 4 Path integral in Clifford algebra representation

### 4.1 Path integral

We use the representation (12) for the generators  $T_a$  and standard techniques [38, 58] from the spinning particle case, adapted to our present problem, to represent the causal propagator. In this case, the indices  $\alpha$ ,  $\beta$  and  $\gamma$  label the matrix entries of the  $\Gamma$ -matrices, that is, they label the representation space for the Clifford algebra. The propertime representation for the operator  $\hat{D}$  (7) in the position representation is

$$D(x_{out}, x_{in}) = i \int_0^\infty \langle x_{out} | e^{-i\hat{H}(\lambda)} | x_{in} \rangle \, d\lambda \,.$$
(30)

Next a discretization is made inserting N-1 identity resolutions  $I = \int dx |x\rangle \langle x|$  in the above expression,

$$D(x_{out}, x_{in}) = \lim_{N \to \infty} i \int_0^\infty d\lambda_0 \int_{-\infty}^\infty \left(\prod_{i=1}^{N-1} dx_i\right) d\lambda_1 \cdots d\lambda_N$$
$$\prod_{i=1}^N \langle x_i | e^{-i\hat{H}(\lambda_i)/N} | x_{i-1} \rangle \,\delta(\lambda_i - \lambda_{i-1})$$
(31)

where  $x_N = x_{out}$  and  $x_0 = x_{in}$ . Applying the symmetric or Weyl correspondence to the general matrix element, one has

$$\langle x_i | e^{-i\hat{H}(\lambda_i)/N} | x_{i-1} \rangle = \int \frac{dp_i}{(2\pi)^4} \exp\left(-\frac{i}{N} H\left(\lambda_i, \frac{x_1 + x_2}{2}, p_i\right)\right) e^{i(x_i - x_{i-1})p_i}, \quad (32)$$

where H is the Weyl symbol of  $\hat{H}$ ,

$$H(\lambda, x, p) = \lambda \left[ m^2 - \left( p_i^2 + q p_i^{\mu} A_{\mu}^a(x) T_a \right)^2 \right] \,.$$

As in the spinning particle case [38], one assigns to each matrix  $T_a$  its 'time'  $\tau_j = j\Delta\tau$ , so that the time-ordered (31) becomes, for  $1/N \equiv \Delta\tau$ ,

$$D(x_{out}, x_{in}) = \lim_{\Delta \tau \to 0} iT \int_0^\infty d\lambda \int_{-\infty}^\infty \left(\prod_{i=1}^{N-1} dx_i\right) \left(\prod_{i=1}^N \frac{\mathrm{d}p_i}{(2\pi)^4} d\lambda_i \frac{d\pi_i}{2\pi}\right)$$
$$\times \exp i \sum_{i=1}^N S_i(x_i, x_{i-1}, p_i, \lambda_i, \pi_i) , \qquad (33)$$

where

$$S_{i} = \left(\frac{x_{i} - x_{i-1}}{\Delta \tau} \cdot p_{i} - H\left(\lambda_{i}, \frac{x_{i} + x_{i-1}}{2}, p_{i}\right) + \pi_{i} \frac{\lambda_{i} - \lambda_{i-1}}{\Delta \tau}\right) \Delta \tau .$$
(34)

In the limit  $\Delta \tau \to 0$ ,  $S_i \to S_H[x, p; \tau_{in}, \tau_{out}]$  is the Hamiltonian action, a functional of the trajectory (x(t), p(t)) in phase space, defined in the proper-time interval  $[\tau_{in}, \tau_{out}]$ , and (33) is the discrete version of the following path integral in the Hamiltonian form:

$$D(x_{out}, x_{in}) = iT \int_0^\infty d\lambda_0 \int_{x_{in}}^{x_{out}} Dx \int Dp \int_{\lambda_0} D\lambda D\pi \exp i \int_{\tau_{in}}^{\tau_{out}} \left( \dot{x} \cdot p - H(\lambda, x, p) + \pi \dot{\lambda} \right) d\tau \,.$$
(35)

Following [38], we introduce odd sources  $\rho_a(\tau)$ , anticommuting with the  $\Gamma$ -matrices, and rewrite 35 as

$$D(x_{out}, x_{in}) = i \int_0^\infty d\lambda_0 \int_{x_{in}}^{x_{out}} Dx \int Dp \int_{\lambda_0} D\lambda D\pi \exp i \int_0^1 \left[ \lambda \left( \left( p_\mu + \frac{q}{4} t^\alpha_{\alpha\beta} A^a_\mu \frac{\delta_l}{\delta\rho_\alpha} \frac{\delta_l}{\delta\rho_\beta} \right)^2 - m^2 \right) p \cdot \dot{x} + \pi \dot{\lambda} d\tau \right] \times T \int_0^1 \rho_\alpha(\tau) \Gamma^\alpha d\tau \Big|_{\rho=0},$$

where for simplicity we have made  $\tau_{in} = 0$  and  $\tau_{out} = 1$ . It is possible to present the last term on the right-hand side of the above equation as a path integral [38, 58],

$$T \int_{0}^{1} \rho_{\alpha}(\tau) \Gamma^{\alpha} d\tau = \exp\left(i\Gamma^{\alpha} \frac{\partial_{l}}{\partial\theta^{\alpha}}\right)$$

$$\times \int_{\psi(0)+\psi(1)=\theta} \exp\left[\int_{0}^{1} \left(\psi^{\alpha}(\tau) \dot{\psi}_{\alpha}(\tau) - i2\rho_{\alpha}(\tau) \psi^{\alpha}(\tau)\right) d\tau + \psi^{\alpha}(1) \psi_{\alpha}(0)\right] \mathcal{D}\psi\right|_{\theta=0}$$

$$\mathcal{D}\psi = D\psi\left[\int_{\psi(0)+\psi(1)=0} \exp\int_{0}^{1} \psi^{\alpha}(\tau) \dot{\psi}_{\alpha}(\tau) d\tau\right]^{-1},$$

where  $\theta$  are odd constants, anticommuting with the  $\Gamma$ -matrices. Then, we arrive at the Hamiltonian path-integral representation for the propagator:

$$D(x_{out}, x_{in}) = i \exp\left(i\Gamma^{\alpha} \frac{\partial_{l}}{\partial\theta^{\alpha}}\right) \int_{0}^{\infty} d\lambda_{0} \int_{x_{in}}^{x_{out}} Dx \int Dp \int_{\lambda_{0}} D\lambda D\pi$$
$$\int \exp\left\{i \int_{0}^{1} \left[\lambda \left(\left(p_{\mu} - qt_{\alpha\beta}^{\alpha} A_{\mu}^{a} \psi_{\alpha} \psi_{\beta}\right)^{2} - m^{2}\right)\right. \\\left. -i\psi^{\alpha} \dot{\psi}_{\alpha} + p \cdot \dot{x} + \pi \dot{\lambda}\right] d\tau + \psi^{\alpha} (1) \psi_{\alpha} (0) \right\} \mathcal{D}\psi\Big|_{\theta=0},$$
$$x(0) = x_{in}, \ x(1) = x_{out}, \ \lambda(0) = \lambda_{0}, \ \psi(0) + \psi(1) = \theta.$$

Integrating over the momenta, one finds the Lagrangian path-integral representation:

$$D(x_{out}, x_{in}) = i \exp\left(i\Gamma^{\alpha} \frac{\partial_l}{\partial \theta^{\alpha}}\right) \int_0^{\infty} de_0 \int \exp\left\{i\left(S_{eff} + S_G\right) + \psi^{\alpha}\left(1\right)\psi_{\alpha}\left(0\right)\right\} M\left[e, x\right] Dx De D\pi \mathcal{D}\psi|_{\theta=0}$$

$$S_{eff} = i \int_0^1 \left(-\frac{\dot{x}^2}{2e} - \frac{e}{2}m^2 + qt^{\alpha}_{a\beta}\dot{x}^{\mu}A^a_{\mu}\psi_{\alpha}\psi_{\beta} - i\psi^{\alpha}\dot{\psi}_{\alpha}\right)$$

$$x\left(0\right) = x_{in}, \ x\left(1\right) = x_{out}, \ e\left(0\right) = e_0, \ \psi\left(0\right) + \psi\left(1\right) = \theta,$$
(36)

where the measure M[e, x] and  $S_G$  are

$$M[e,x] = \int Dp \exp{\frac{i}{2} \int_0^1 ep^2 d\tau}, \ S_G = \int_0^1 \pi \dot{e}.$$

### 4.2 Pseudoclassical action

Let us consider the reparametrization invariant action from (36) with the rescaling  $\psi \to i/\sqrt{2}\psi$ ,

$$S_{eff} = \int dx^4 \left( -\frac{\dot{x}^2}{2e} - \frac{e}{2}m^2 - q\dot{x}^{\mu}A^a_{\mu}I_a + \frac{i}{2}\psi^{\alpha}\dot{\psi}_{\alpha} \right) , \ I_a = \frac{1}{2}t^{\alpha}_{\alpha\beta}\psi_{\alpha}\psi_{\beta} .$$
(37)

The above action is essentially the one written in [48] in the case the set of Grassmann variables  $\psi$  belong to the adjoint representation of a compact simple group  $G(t_{a\beta}^{\alpha} = f_{ac}^{b})$ , and in [47] for  $\psi$  in a representation with antisymmetric generators  $t_{a}$ . The equations of motion in the gauge  $e = \sqrt{\dot{x}^2}/m$  are

$$\begin{split} \frac{d}{dt} \left( m \frac{\dot{x}^{\mu}}{\sqrt{\dot{x}^2}} \right) &= q F^a_{\mu\nu} \dot{x}^{\nu} I_a \,, \ D_t \psi^{\alpha} \equiv \frac{d}{dt} \psi^{\alpha} + i q \dot{x}^{\mu} A^a_{\mu} t^{\alpha}_{a\beta} \psi^{\beta} = 0 \,, \\ F^a_{\mu\nu} &= \partial_{\mu} A^a_{\nu} - \partial_{\nu} A^a_{\mu} + i q f^a_{bc} A^b_{\mu} A^c_{\nu} \,, \end{split}$$

where  $F^a_{\mu\nu}$  is the non-Abelian field strength.

Next, we follow a similar canonical analysis path than the one taken in the case of the coherent representation, this time with  $t_a$  denoting  $n \times n$  antisymmetric matrices. As expected, the Hamiltonian is proportional to constraints,

$$H = -\frac{e}{2}T - \dot{\psi}^{\alpha}\phi_{\alpha}$$

where

$$T = (p_{\mu} + qA_{\mu}^{a}I_{a})^{2} - m^{2}, \ \phi_{\alpha} = \pi_{\alpha} - \frac{i}{2}\psi_{\alpha}$$

After redefining T through the shift  $\psi \to \psi - i\phi$ ,  $T \to \tilde{T}$ , the constraint algebra becomes

$$\left\{\tilde{T},\phi_{\alpha}\right\} = 0, \ \left\{\phi_{\alpha},\phi_{\beta}\right\} = -i\delta_{\alpha\beta}.$$

The set  $\Phi = \left\{\tilde{T}, \phi\right\}$  is first-class, and the evolution of the independent variables  $\eta = (x, p, \psi)$  is

$$\dot{\eta} = \{\eta, \Lambda T\}_{D(\phi)} = 0, \ T = \phi_{\alpha} = 0$$

where the Dirac brackets are defined with respect to the second-class constraint set  $\{\phi\}$ . The Dirac commutator of the independent variables is

$$\{x^{\mu}, p_{\nu}\}_{D(\phi)} = \delta^{\mu}_{\nu}, \ \{\psi_{\alpha}, \psi_{\beta}\}_{D(\phi)} = -i\delta_{\alpha\beta},$$

The isospin quantities  $I_a$  satisfy the Lie algebra of SU(N) after quantization and are covariantly constant:

$$\{I_a, I_b\}_{D(\phi)} = -if^c_{ab}I_c , \ D_\tau I^c = \frac{d}{d\tau}I^c + iq\dot{x}^{\mu}A^a_{\mu}f^c_{ab}I^b = 0 \,.$$

It is clear that upon quantization the Grassmann variables  $\psi_{\alpha}$  generate Clifford algebra with n generators and positive-definite inner product. And thus the physical states  $\phi$  are  $2^{[n/2]}$ -component vectors satisfying

$$\left[ \left( \hat{p}_{\mu} + q A^{a}_{\mu} \hat{I}_{a} \right)^{2} - m^{2} \right] \phi \left( x \right) = 0 , \qquad (38)$$

where the quantum-mechanical isospin operators  $\hat{I}_a = \frac{1}{4} t^{\alpha}_{a\beta} \Gamma_{\alpha} \Gamma_{\beta}$  are precisely those introduced in (12) and they satisfy the su(N) algebra (2)

$$\left[\hat{I}_a, \hat{I}_b\right] = f_{ab}^c \hat{I}_c \,.$$

Let us draw a similar analysis of the isospin content for the classical theory as the one given in section 3.1. Here, the Grassmannian kinetic terms in the action are invariant under the transformations generated by  $R_{\alpha\beta} = -i\psi_{\alpha}\psi_{\beta}$ ,

$$\delta_{\omega}\psi_{\alpha} = \left\{\frac{1}{2}\omega_{\beta\gamma}R_{\beta\gamma},\psi_{\alpha}\right\}_{D(\phi)} = \omega_{\alpha\beta}\psi_{\beta}, \ \omega_{\alpha\beta} = -\omega_{\beta\alpha},$$

which give a representation for the Lie algebra so(n):

$$\{R_{\alpha\beta}, R_{\gamma\delta}\}_{D(\phi)} = \delta_{\alpha\gamma}R_{\beta\delta} - \delta_{\beta\gamma}R_{\alpha\delta} - \delta_{\alpha\delta}R_{\beta\gamma} + \delta_{\beta\delta}R_{\alpha\gamma} \,.$$

So the the generators  $I_a$  are a linear combination of the generators  $L_{\alpha\beta}$  of so(n), and therefore they generate an su(N) subalgebra of so(n). In order to determine the SU(N) content of the wave function, one proceeds as in [47]: given  $t_a$  an irreducible representation of su(N) in terms of  $n \times n$  **antisymmetric** matrices, it is clear  $\psi_{\alpha}$  gives rise to a Clifford algebra in the quantum theory, so the wave function can be taken to belong to a spinor representation of so(n). One then calculates the eigenvalues of a maximal set of commuting generators  $I_a$  for this representation, and thus determines how it decomposes in irreducible representations of SU(N). Therefore, to each given *n*-dimensional irreducible representation  $t_a$  of SU(N), the wave function will belong to a  $2^{[n/2]}$ -dimensional representation of SO(n), which decomposes into irreducible representations of SU(N) as determined by the isospin generators  $I_a$ .

For example, in the case of SU(2), one can find a basis for which  $t_a$  are antisymmetric, and in which  $I_1$  decomposes as

$$I_1 = L_{23} + 2L_{45} + \dots + sL_{2s,2s+1}$$
.

The wave function is a  $2^s$ -component spinor of SO(2s+1). Below we give the SU(2) decomposition of the spinor SO(2s+1) representation for some values of isospin:

isospin	symmetry group	representation dimension	decomposition
1	$SO\left(3 ight)$	2	$\frac{1}{2}$
2	$SO\left(5 ight)$	4	$\frac{3}{2}$
3	$\overline{SO}(7)$	8	$\underline{0} + \underline{3}$

### 5 Summary

We have described two methods of generating classical actions for a scalar particle with isospin via path-integral representations of the causal propagator. Dirac quantization of these actions produce the corresponding wave equations for various possible representations of SU(N). By means of a judicious choice of the pseudo-classical action and the representation of the su(N) algebra in the action, it is possible to obtain the wave action for any desired isospin.

Acknowledgment: R.F. thanks FAPESP for support and D.M.G acknowledges FAPESP and CNPq for permanent support.

# A Weyl ordering of operators and functions in the Berezin algebra

Let us write the Hamiltonian operator (7) explicitly:

$$\hat{H} = -\lambda \left( \hat{p}^2 + q t_{a\alpha\beta} \left( \hat{p}^{\mu} \hat{A}^a_{\mu} + \hat{A}^a_{\mu} \hat{p}^{\mu} \right) \hat{a}^{\dagger}_{\alpha} \hat{a}_{\beta} + q^2 t_{a\alpha\beta} t_{b\gamma\delta} \hat{A}^a_{\mu} \hat{A}^{\mu b} \hat{a}^{\dagger}_{\alpha} \hat{a}_{\beta} \hat{a}^{\dagger}_{\gamma} \hat{a}_{\delta} - m^2 \right) \,.$$

Total symmetrization in  $\hat{x}$  and  $\hat{p}$ , and total antisymmetrization in  $a^{\dagger}$  and a gives the Weyl-ordered Hamiltonian operator  $\hat{H}_W$ :

$$\hat{H}_W = -\lambda \left( \hat{p}^2 + \frac{q}{2} t_{a\alpha\beta} \left( \hat{p}^{\mu} \hat{A}^a_{\mu} + \hat{A}^a_{\mu} \hat{p}^{\mu} \right) \left[ \hat{a}^{\dagger}_{\alpha}, \hat{a}_{\beta} \right] + q^2 t_{a\alpha\beta} t_{b\gamma\delta} \hat{A}^a_{\mu} \hat{A}^{\mu b} \left( \hat{a}^{\dagger}_{\alpha} \hat{a}_{\beta} \hat{a}^{\dagger}_{\gamma} \hat{a}_{\delta} \right)_W - m^2 \right) , \quad (39)$$

where the four-fermion term is given by

$$\hat{a}_{\alpha}^{\dagger}\hat{a}_{\beta}\hat{a}_{\gamma}^{\dagger}\hat{a}_{\delta} = \left(\hat{a}_{\alpha}^{\dagger}\hat{a}_{\beta}\hat{a}_{\gamma}^{\dagger}\hat{a}_{\delta}\right)_{W} + \frac{1}{2}\delta_{\gamma\delta}\left(\hat{a}_{\alpha}^{\dagger}\hat{a}_{\beta}\right)_{W} - \frac{1}{2}\delta_{\delta\alpha}\left(\hat{a}_{\gamma}^{\dagger}\hat{a}_{\beta}\right)_{W} + \delta_{\alpha\beta}\left(\hat{a}_{\gamma}^{\dagger}\hat{a}_{\delta}\right)_{W} + \frac{1}{2}\delta_{\gamma\beta}\left(\hat{a}_{\alpha}^{\dagger}\hat{a}_{\delta}\right)_{W} - \frac{1}{4}\delta_{\delta\alpha}\delta_{\gamma\beta} - \delta_{\alpha\beta}\delta_{\gamma\delta}$$

Using the tracelessness of the matrices  $t_a$  and antisymmetry of the structure constants  $f_{abc}$ , we have

$$\hat{H} = \hat{H}_W + \lambda \frac{q^2}{4} \operatorname{tr}\left(t_a t_b\right) \hat{A}^a_\mu \hat{A}^{\mu l}$$

Thus, the Hamiltonian is the sum of a Weyl-ordered expression plus a gauge non-invariant contribution. The Weyl-symbol corresponding to  $\hat{H}_W$  is

$$H_W = -\lambda \left( p^2 + 2q t_{a\alpha\beta} \left( p^{\mu} A^a_{\mu} \right) \bar{\chi}_{\alpha} \chi_{\beta} + q^2 t_{a\alpha\beta} t_{b\gamma\delta} A^a_{\mu} A^{\mu b} \bar{\chi}_{\alpha} \chi_{\beta} \bar{\chi}_{\gamma} \chi_{\delta} - m^2 \right)$$
(40)

# A.1 Proof<sup>4</sup> of fermionic midpoint rule.

If  $F(\hat{a}, \hat{a}^{\dagger})$  is any Weyl-ordered polynomial in  $\hat{a}$  and  $\hat{a}^{\dagger}$ , then

$$\langle \bar{\chi} | F\left(\hat{a}, \hat{a}^{\dagger}\right) | \chi \rangle = \int d\bar{\eta} d\eta \, \langle \bar{\chi} | \eta \rangle F\left(\frac{\chi + \eta}{2}, \bar{\eta}\right) \langle \bar{\eta} | \chi \rangle , \qquad (41)$$

$$= \int d\bar{\eta} d\eta \left\langle \bar{\chi} \right| \left. \eta \right\rangle F\left(\chi, \frac{\bar{\chi} + \bar{\eta}}{2}\right) \left\langle \bar{\eta} \right. \left| \chi \right\rangle \,, \tag{42}$$

Let us prove the identity (41). The proof of the second identity is analogous. First, consider  $F(\hat{a}^{\dagger})$  a polynomial in creation operators. Clearly, F is Weyl-ordered, and (41) is trivially satisfied,

$$\left\langle \bar{\chi} \right| F\left(\hat{a}^{\dagger}\right) \left| \chi \right\rangle = \int d\bar{\eta} d\eta \left\langle \bar{\chi} \right| \eta \right\rangle F\left( \bar{\chi} \right) \left\langle \bar{\eta} \left| \chi \right\rangle \,.$$

Now, for  $F(\hat{a}, \hat{a}^{\dagger}) = \frac{1}{2} \left( \hat{a}_{\alpha} f(\hat{a}^{\dagger}) + (-1)^{\varepsilon(f)} f(\hat{a}^{\dagger}) \hat{a}_{\alpha} \right)$ , where the plus or minus sign depends on the parity of  $f(\hat{a}^{\dagger})$ , (41) is easily seen to hold. Any Weyl-ordered polynomial can be obtained by repeated antisymmetrizations of the form  $F = \frac{1}{2} \left( \hat{a}_{\alpha} f \pm f \hat{a}_{\alpha} \right)$  where  $f(\hat{a}, \hat{a}^{\dagger})$  is Weyl-ordered. Therefore, let us prove (41) inductively, by assuming it holds for  $f(\hat{a}, \hat{a}^{\dagger})$  and proving it is also true for  $F = \frac{1}{2} \left( \hat{a}_{\alpha} f \pm f \hat{a}_{\alpha} \right)$ ,

$$\begin{split} \langle \bar{\chi} | \frac{1}{2} \left( \hat{a}_{\alpha} f \pm f \hat{a}_{\alpha} \right) | \chi \rangle &= \int d\bar{\eta} d\eta \frac{1}{2} \left( \langle \bar{\chi} | \hat{a}_{\alpha} | \eta \rangle \langle \bar{\eta} | f | \chi \rangle \pm \langle \bar{\chi} | \eta \rangle \langle \bar{\eta} | f \hat{a}_{\alpha} | \chi \rangle \right) \\ &= \int d\bar{\eta} d\eta \langle \bar{\chi} | \eta \rangle \frac{\eta_{\alpha} + \chi_{\alpha}}{2} \langle \bar{\eta} | f | \chi \rangle \\ &= \int d\bar{\eta} d\eta d\bar{\xi} d\xi \langle \bar{\chi} | \eta \rangle \langle \bar{\eta} | \xi \rangle \frac{\chi_{\alpha} + \eta_{\alpha}}{2} f\left( \frac{\chi + \xi}{2}, \bar{\xi} \right) \langle \bar{\xi} | \chi \rangle \\ &= d\bar{\xi} d\xi \langle \bar{\chi} | \xi \rangle \frac{\chi_{\alpha} + \xi_{\alpha}}{2} f\left( \frac{\chi + \xi}{2}, \bar{\xi} \right) \langle \bar{\xi} | \chi \rangle \end{split}$$

where in the last equality we used the identity

$$\int d\bar{\eta} d\eta \langle \bar{\alpha} | \eta \rangle \langle \bar{\eta} | \beta \rangle f(\eta) = \langle \bar{\alpha} | \beta \rangle f(\beta) .$$
(43)

# References

 Lebedev Phys. Inst. Quantum Electrodynamics of Phenomena in Intense Fields, 111. Nauka Moscow, 1979.

<sup>&</sup>lt;sup>4</sup>Adapted from [59]

- [2] A. A. Grib, S.G. Mamaev, and V.M. Mostepanenko. Vacuum Quantum Effects in Strong Fields. Friedmann Laboratory Publishing, St. Petersburg, 1994.
- [3] E. S. Fradkin, D. M. Gitman, and Sh. M. Shvartsman. Quantum electrodynamics with unstable vacuum. Springer, 1991. Berlin, Germany: Springer (1991) 288 p. (Springer series in nuclear and particle physics).
- [4] W. Greiner, B. Muller, and J. Rafelski. QUANTUM ELECTRODYNAMICS OF STRONG FIELDS. Springer, 1985. Berlin, Germany: Springer (1985) 594 P. (Texts and Monographs In Physics).
- [5] Jan Ambjorn, R. J. Hughes, and N. K. Nielsen. Action principle of bogolyubov coefficients. Ann. Phys., 150:92, 1983.
- [6] Jan Ambjorn and Richard J. Hughes. Particle creation in color electric fields. *Phys. Lett.*, B113:305, 1982.
- [7] Jan Ambjorn and Richard J. Hughes. Canonical quantization in nonabelian background fields.
   1. Ann. Phys., 145:340, 1983.
- [8] A. I. Vainshtein, Valentin I. Zakharov, V. A. Novikov, and Mikhail A. Shifman. Calculations in external fields in qcd: An operator method. (in russian). Sov. J. Nucl. Phys., 39:77, 1984.
- [9] Martin Reuter, Michael G. Schmidt, and Christian Schubert. Constant external fields in gauge theory and the spin 0, 1/2, 1 path integrals. Annals Phys., 259:313–365, 1997.
- [10] G. K. Savvidy. Infrared instability of the vacuum state of gauge theories and asymptotic freedom. *Phys. Lett.*, B71:133, 1977.
- [11] N. K. Nielsen and P. Olesen. An unstable yang-mills field mode. Nucl. Phys., B144:376, 1978.
- [12] C. Ragiadakos. A stable symmetrized savvidy vacuum. Phys. Lett., B100:471, 1981.
- [13] Asim Yildiz and Paul H. Cox. Vacuum behavior in quantum chromodynamics. Phys. Rev., D21:1095, 1980.
- [14] M. Claudson, A. Yildiz, and P. H. Cox. Vacuum behavior in quantum chromodynamics. ii. *Phys. Rev.*, D22:2022–2026, 1980.
- [15] Stephen L. Adler. An overrelaxation method for the monte carlo evaluation of the partition function for multiquadratic actions. *Phys. Rev.*, D23:2901, 1981.
- [16] Walter Dittrich and Martin Reuter. Effective qcd lagrangian with zeta function regularization. Phys. Lett., B128:321, 1983.
- [17] Curt A. Flory. A selfdual gauge field, its quantum fluctuations, and interacting fermions. *Phys. Rev.*, D28:1425, 1983.
- [18] Y. M. Cho and D. G. Pak. Dynamical symmetry breaking and magnetic confinement in qcd. hep-th, 2000.
- [19] Steven K. Blau, Matt Visser, and Andreas Wipf. Analytical results for the effective action. Int. J. Mod. Phys., A6:5409-5433, 1991.

- [20] Abhijit Bhattacharyya et al. Cosmological qcd phase transition and dark matter. Nucl. Phys., A661:629-632, 1999.
- [21] Gouranga C. Nayak and Peter van Nieuwenhuizen. Soft-gluon production due to a gluon loop in a constant chromo-electric background field. *Phys. Rev.*, D71:125001, 2005.
- [22] Dmitri Kharzeev, Eugene Levin, and Kirill Tuchin. Multi-particle production and thermalization in high-energy qcd. Phys. Rev., C75:044903, 2007.
- [23] E. M. Serebryanyi. Vacuum polarization by magnetic flux: The aharonov-bohm effect. Theor. Math. Phys., 64:846-855, 1985.
- [24] Pawel Gornicki. Aharonov-bohm effect and vacuum polarization. Ann. Phys., 202:271–296, 1990.
- [25] B.M. Barbashov. JETP, 48:607, 1965.
- [26] I. A. Batalin and E. S. Fradkin. Quantum electrodynamics in external fields. 1. Teor. Mat. Fiz., 5:190-218, 1970.
- [27] Marc Henneaux and Claudio Teitelboim. Relativistic quantum mechanics of supersymmetric particles. Ann. Phys., 143:127, 1982.
- [28] N. v. Borisov and P. p. Kulish. Path integral in superspace for a relativistic spinor particle in an external gauge field. *Theor. Math. Phys.*, 51:535–540, 1982.
- [29] V. Ya. Fainberg and A. V. Marshakov. Local supersymmetry and dirac particle propagator as a path integral. Nucl. Phys., B306:659, 1988.
- [30] Alexander M. Polyakov. GAUGE FIELDS AND STRINGS. CHUR, SWITZERLAND: HAR-WOOD, 1987.
- [31] A. V. Marshakov and V. Ya. Fainberg. Representation of the propagator of a dirac particle in an external gravitational field as a sum over paths. JETP Lett., 47:565–568, 1988.
- [32] V. Ya. Fainberg and A. V. Marshakov. A propagator for fermionic string. Phys. Lett., B211:81, 1988.
- [33] V.Ya. Fainberg and A.V. Marshakov. Proc. Phian, 201, 1990.
- [34] Jan Ambjorn, Bergfinnur Durhuus, and Thordur Jonsson. A random walk representation of the dirac propagator. Nucl. Phys., B330:509, 1990.
- [35] J. Grundberg, T. H. Hansson, and A. Karlhede. Geometrical phases from spinning particles. *Phys. Rev.*, D41:2642, 1990.
- [36] G. P. Korchemsky. Quantum geometry of the dirac fermions. *Phys. Lett.*, B232:334, 1989.
- [37] J. Grundberg, T. H. Hansson, and A. Karlhede. On polyakov's spin factors. Nucl. Phys., B347:420-440, 1990.
- [38] E. S. Fradkin and D. M. Gitman. Path integral representation for the relativistic particle propagators and bfv quantization. *Phys. Rev.*, D44:3230-3236, 1991.

- [39] D. M. Gitman and A. V. Saa. Quantization of spinning particle with anomalous magnetic momentum. Class. Quant. Grav., 10:1447–1460, 1993.
- [40] D. M. Gitman and Sh. M. Shvartsman. Spinor and isospinor structure of relativistic particle propagators. *Phys. Lett.*, B318:122-126, 1993.
- [41] G. P. Korchemsky. Quantum geometry of dirac fermions. Int. J. Mod. Phys., A7:339–380, 1992.
- [42] T. M. Aliev, V. Ya. Fainberg, and N. K. Pak. Path integral for spin: A new approach. Nucl. Phys., B429:321-343, 1994.
- [43] J. W. van Holten. Propagators and path integrals. Nucl. Phys., B457:375-407, 1995.
- [44] A. I. Karanikas and C. N. Ktorides. Particle path integral approach to the study of dirac spin 1/2 field systems. *Phys. Rev.*, D52:5883–5897, 1995.
- [45] Dimitri M. Gitman. Path integrals and pseudoclassical description for spinning particles in arbitrary dimensions. Nucl. Phys., B488:490–512, 1997.
- [46] Bodo Geyer, Dmitry Gitman, and Ilya L. Shapiro. Path integral and pseudoclassical action for spinning particle in external electromagnetic and torsion fields. Int. J. Mod. Phys., A15:3861– 3876, 2000.
- [47] A. P. Balachandran, Per Salomonson, Bo-Sture Skagerstam, and Jan-Olof Winnberg. Classical description of particle interacting with nonabelian gauge field. *Phys. Rev.*, D15:2308, 1977.
- [48] A. Barducci, R. Casalbuoni, and L. Lusanna. Classical scalar and spinning particles interacting with external yang-mills fields. *Nucl. Phys.*, B124:93, 1977.
- [49] S. K. Wong. Field and particle equations for the classical yang-mills field and particles with isotopic spin. Nuovo Cim., A65S10:689-694, 1970.
- [50] M. Dresden and S. F. Chen. Solitons, gauge theories and the 'great einstein theorem'. *Physica*, 83A:1, 1975.
- [51] Rainer Dick. Vector and scalar confinement in gauge theory with a dilaton. *Phys. Lett.*, B409:321-324, 1997.
- [52] Rainer Dick. Confinement from a massive scalar in QCD. Eur. Phys. J., C6:701-703, 1999.
- [53] M. Chabab, R. Markazi, and E. H. Saidi. On the confining potential in 4d su(n(c)) gauge theory with dilaton. *Eur. Phys. J.*, C13:543-549, 2000.
- [54] Michael B. Green, J. H. Schwarz, and Edward Witten. SUPERSTRING THEORY. Cambridge University Press, 1987. Cambridge, Uk: Univ. Pr. (1987) (Cambridge Monographs On Mathematical Physics).
- [55] Y. Ohnuki and T. Kashiwa. Coherent states of fermi operators and the path integral. Prog. Theor. Phys., 60:548, 1978.
- [56] D. M. Gitman and I. V. Tyutin. Quantization of fields with constraints. Springer, 1990. Berlin, Germany: Springer (1990) 291 p. (Springer series in nuclear and particle physics).

- [57] D. M. Gitman and I. V. Tyutin. Hamiltonization of theories with degenerate coordinates. Nucl. Phys., B630:509-527, 2002.
- [58] A. N. Vasiliev. Functional methods in quantum field theory and statistical physics. Gordon and Breach Science Publishers, 1986. Amsterdam, Netherlands: Gordon and Breach (1998) 312 p.
- [59] Giuliano M. Gavazzi. Weyl-ordered fermions and path integrals. J. Math. Phys., 30, December 1989.