

Double Parton Scattering in Associate Higgs Boson Production with Bottom Quarks at Hadron Colliders

M. Y. Hussein^{1 a}

Department of Physics, College of Science, University of Bahrain, P.O. Box 32038, Kingdom of Bahrain

Abstract. Higgs boson production in association with bottom quarks, is one of the most important discovery channels for Higgs particles in the Standard Model (SM) and its supersymmetric extension at the LHC pp collider. The theoretical prediction of the corresponding cross section has been improved by including the complete next-to-leading order QCD corrections. We review the status results for the leading order to the partonic process and present calculations for the integrated cross-section in both single and double parton scattering collisions.

PACS. 14.80.Bn Standard-model Higgs bosons { 14.80.Cp Non-standard-model Higgs bosons { 11.80.La Multiple scattering { 13.85.Hd Inelastic scattering: many-particle final states

1 Introduction

The discovery of the SM Higgs boson is one of the most important goals and pressing issues of present and future colliders. An important prerequisite for identifying the most convenient signatures for detecting Higgs boson needs a precise knowledge of the various production cross-sections, decay branching ratios, decay width, and their couplings to other particles.

Recently, much progress has been made in the detection of a Higgs boson, next-leading order corrections are now known for most sub-process [1,2,3,4,5,6], knowledge of parton distribution functions has improved as more deep inelastic data become available and the range of possible input parameter values decreased.

The dominant production of a SM Higgs boson in hadronic interactions is through gluon-gluon fusion [7,8,9]. Various channels can be explored to search for Higgs boson at hadron colliders. There are only a few Higgs production mechanism which lead to detectable cross-section. Each of them use the preference of coupling of the SM Higgs to heavy particles either massive vector bosons or massive quarks.

The associated production of a Higgs boson with a pair of bb quarks has a small cross-section which is due to small size of Yukawa coupling $g_{bbH} = m_b/v \approx 0.02$ in the SM [10]. In some extensions of the SM, such as the MSSM, the Yukawa coupling of b-quarks can become strongly enhanced, the associate production of a Higgs boson with a pair of bb quarks can dominate over other production channels and this production mechanism can be a significant source of Higgs bosons [11].

Detecting two bottom quarks in the final state identifies uniquely the Higgs coupling responsible for the

enhanced cross-section and drastically reduces the background. This corresponds to an experiment measuring the Higgs decay along two high pt bottom quark jets.

In a four-flavor-number scheme with no b-quarks in the initial state, the lowest order processes are the tree level contributions from $gg \rightarrow b\bar{b}H$ and $qq \rightarrow b\bar{b}H$.

At high energies and due to large flux in particular at the LHC, another type of scattering mechanism contributes to the cross section besides to single scattering. Thus, for $qq \rightarrow b\bar{b}H$ production there would be two competing mechanisms: single parton scattering and double parton scattering featuring two Drell-Yan processes happening simultaneously.

The purpose of the present work is to point out that the same $b\bar{b}H$ final state can be produced also by double parton scattering collision process. The large rate of production of bb pairs expected at the LHC gives rise to a relatively large probability of production of a $b\bar{b}H$ in the process underlying the H production. In fact as a result of the present analysis it is found that double parton scattering may represent a rather sizable source of background.

2 Double Parton Scattering Mechanism

The multiple parton scattering occurs when two or more different pairs of parton scatter independently in the same hadronic collision [12,13,14].

With the only assumption of factorization of the two hard parton processes A and B, the inclusive cross section of a double parton-scattering in a hadronic collision is expressed by:

$$D_{(A;B)} = \frac{m}{2} \sum_{i,j;k,l} X_{ij}(x_1; x_2; b) \hat{A}_{ik}^A(x_1; x_1^0)$$

^a E-mail: m.hussein@sci.uob.bh

$$\hat{\sigma}_{j1}^B(x_2; x_2^0)_{k1}(x_1^0; x_2^0; b) dx_1 dx_1^0 dx_2 dx_2^0 d^2b \quad (1)$$

Where $\hat{\sigma}_{ij}(x_1; x_2; b)$ are the double parton distribution function, depending on the fractional momenta x_1, x_2 and the relative transverse distance b of the two parton undergoing the hard processes A and B , the indices i and j refer to the different parton species and $\hat{\sigma}_{ik}^A$ and $\hat{\sigma}_{jl}^B$ are the partonic cross section. The factor $m=2$ is for symmetry, specially $m=1$ for indistinguishable parton processes and $m=2$ for distinguishable processes.

The double distributions $\hat{\sigma}_{ij}(x_1; x_2; b)$ are the main reason of interest in multiparton collisions. This distribution contains in fact all the information of probing the hadron in two different points contemporarily through the hard processes A and B . The cross section for multiparton process is sizable when the flux of partons is large, namely at small x . Given the large flux one may hence expect that correlations in momentum fraction will not be a major effect and partons to be rather correlated in transverse space. Neglecting the effect of parton correlations in x one writes:

$$\hat{\sigma}_{ij}(x_1; x_2; b) = \hat{\sigma}_i(x_1) \hat{\sigma}_j(x_2) F_{ij}^i(b) \quad (2)$$

where $\hat{\sigma}_i(x)$ the usual one body parton distribution $F_{ij}^i(b)$ is function and is a function normalized to one and representing the pair density in transverse space. The inclusive cross section hence simplifies to:

$$\sigma_{(A;B)}^D = \frac{m}{2} \sum_{ijkl} \hat{\sigma}_{k1}^{ij}(A) \hat{\sigma}_{l1}(B) \quad (3)$$

where $\hat{\sigma}_{j1}(A)$ and $\hat{\sigma}_{k1}(B)$ are the hadronic inclusive cross section for the two partons labeled i and j undergoes the hard interaction labeled A and for two partons k and l to undergo the hard interaction labeled B ;

$$\hat{\sigma}_{k1}^{ij} = \int d^2b F_k^i(b) F_l^j(b) \quad (4)$$

are the geometrical coefficients with dimension an inverse cross section and depending on various parton processes. These coefficients are the experimentally accessible quantities carrying the information of the parton correlation in transverse momentum. The cross section for multiple parton collisions has been further simplified as:

$$\sigma_{(A;B)}^D = \frac{m}{2} \frac{\hat{\sigma}(A) \hat{\sigma}(B)}{\text{eff}} \quad (5)$$

Where all the information on the structure of the hadron in transverse space is summarized in the value of the scale factor, eff .

The experimental value measured by CDF yields [15]

$$\text{eff} = 1.45 \cdot 1.7^{+1.7}_{-2.3} \text{ mb}$$

It is believed that is largely independent of the center-of-mass energy of the collision and on the nature of the partonic interactions. The experimental evidence is not inconsistent with the simplest hypothesis of neglecting correlations in momentum fractions.

3 Cross-Section Results

Higgs search in the $b\bar{b}h$ channel, however employ cuts on the b -quark transverse momenta and thus requires theoretical predictions for exclusive final states. The solution is provided by calculations based on the parton processes $q\bar{q}; g\bar{g} \rightarrow b\bar{b}h$.

In this article we evaluate the fully cross section for $b\bar{b}h$ production by requiring that the transverse momentum of both final state bottom and antibottom quarks be larger than 20 GeV. This corresponds to an experiment measuring the Higgs decay products along with two high pt bottom quark jets. These cuts reduce the cross section by one or two orders of magnitude, but also greatly reduce the background and ensure that the Higgs boson was emitted from a bottom quark and is therefore proportional to the square of the b -quark Yukawa coupling.

The cross section for leading order sub-process for Higgs-boson production in association with bottom quarks obtained using MRST parton distribution [16], the packages MadGraph [17] and HELAS [18] and the integration was performed by VEGAS [19] as function of Higgs mass for the LHC with $\sqrt{s} = 14 \text{ TeV}$ are displayed in Fig. 1.




Fig. 1. Leading order cross section (pb) for Higgs boson production in association with bottom quarks as a function of the Higgs boson mass at the LHC.

The cross-section for Higgs production in association with bottom quarks are not large but may be useful if high luminosity is available, since the Higgs boson can be "tagged" by triggering on the bottom quarks.

Sizable rates of events where two bottom quarks associate with Higgs boson are produced contemporarily at the LHC, as a consequence of the large parton luminosity. The corresponding integrated rate is evaluated by combining the integrated cross section for Higgs boson and production at LHC energy.

If one use the cross section for Higgs boson production from $pp \rightarrow H + X, (b\bar{b})' \approx 5 \cdot 10^2 \text{ b}$ as a value for the scale factor $\text{eff} = 1.45 \text{ mb}$ (the observed

value is $\sigma_{eff} = 1.45 \cdot 1.7^{+1.7}_{-2.3}$ mb) one obtain the cross section for a double parton collision producing a Higgs boson and a $b\bar{b}$ pair.

The large rate of $b\bar{b}$ pair at the LHC gives rise to a relatively sizable production of Higgs boson associated with $b\bar{b}$ quarks.

Figure 2 shows the double parton scattering to the Higgs boson associated with $b\bar{b}$ quarks.




Fig. 2. Total cross section for Higgs boson associated with bottom quarks in the SM as a function of the Higgs boson mass at LHC energy.

4 Summary

In this work we have investigated $b\bar{b}h$ production at the LHC, which is an important discovery channel for Higgs boson in the SM and its extension in the MSSM at large values of $\tan\beta$, where the bottom Yukawa coupling is strongly enhanced.

Our calculations correspond to the cross section for Higgs boson in association with two tagged b jets in single and double parton scattering mechanism.

Although the double parton collision cross section is not large, but it should be taken in consideration because a sizable rate of events where pairs of quarks are produced at the LHC, as a consequence of the large parton luminosity.

However, in the hadron collider environment, the large QCD backgrounds may cause the observation impossible for those events if Higgs bosons decay hadronically. Individual channels with hadronic decays should be studied on case by case.

References

1. T. Han and S. Willenbrock, Phys. Lett. B 273, (1991) 167; T. Han, G. Valencia and S. Willenbrock, Phys. Rev. Lett. 69, (1992) 3274.
2. S. Dittmaier, M. Kramer and M. Spira, Phys. Rev. D 70, (2004) 074010; S. Dawson, C. B. Jackson, L. Reina and D. Wackerth, Phys. Rev. D 69, (2004) 074027;
3. S. Dawson, Nucl. Phys. B 359, (1991) 283.
4. A. Djouadi, M. Spira, and P. M. Zerwas, Phys. Lett. B 264, (1991) 440.
5. D. G. Randenz, M. Spira, and P. M. Zerwas, Phys. Rev. Lett. 70, (1993) 1472.
6. M. Spira, A. Djouadi, D. G. Randenz and P. M. Zerwas, Phys. Lett. B 264, (1991) 440.
7. S. Moch and M. Vogt, Phys. Lett. B 631, (2005) 48; E. Laenen and L. M. Uggone, Phys. Lett. B 632, (2005) 270.
8. T. Han, G. Valencia and S. Willenbrock, Phys. Rev. Lett. 69, (1992) 3274; E. Berger and J. Campbell, Phys. Rev. D 70, (2004) 073011.
9. J. Campbell, R. Ellis, F. Maltoni and S. Willenbrock, Phys. Rev. D 67, (2003) 095002.
10. S. Dawson, C. B. Jackson, L. Reina and D. Wackerth, Int. J. Mod. Phys. A 20, (2005) 3353.
11. Dawson, C. B. Jackson, L. H. Orr, L. Reina and D. Wackerth, Eur. Phys. J. C 33, (2004) 5451.
12. J. Campbell, S. Dawson, S. Dittmaier, C. Jackson, M. Kramer, F. Maltoni, L. Reina, M. Spira, D. Wackerth and S. Willenbrock, hep-ph/0405302.
13. P. V. Landshof and J. C. Polkinghorne, Phys. Rev. D 18, 3344 (1987); C. G. Oebler, F. Halzen and D. H. Scott, Phys. Rev. D 22, (1980) 2789; F. Halzen, P. Hofer and W. J. Stirling Phys. Lett. B 188, (1987) 375; N. Paver and D. Treleani, Nuovo Cim. A 70, (1982) 215; M. Broun and D. Treleani, Eur. Phys. J. C 18, (2001) 511.
14. A. DelFabbro and D. Treleani, Phys. Rev. D 61 (2000) 077501; Phys. Rev. D 63 (2001) 057901.
15. F. Abe et al. (CDF Collaboration), Phys. Rev. D 57, (1997) 503; Phys. Rev. D 56, (1997) 3811.
16. A. D. Martin, R. G. Roberts, W. J. Stirling and R. S. Thorne, Eur. Phys. J. C 14 (2000) 133.
17. T. Stelzer and W. F. Long, Comp. Phys. Comm. 81, (1994) 337.
18. E. Murayama, I. Watanabe and K. Hagiwara, HELAS#ELicity Amplitude Subroutine for Feynman Diagram Evaluations, KEK report 91-11, January 1992.
19. G. P. Lepage, J. Comp. Phys. 27, (1978) 192.