

Di-photon Higgs decay in the MSSM with explicit CP violation

S Hesselbach^{1,*}, S Moretti^{1,2}, S Munir¹ and P Poulose^{1,3}

¹ School of Physics & Astronomy, University of Southampton, Highfield, Southampton SO17 1BJ, UK

² Laboratoire de Physique Théorique, Université Paris-Sud, F-91405 Orsay Cedex, France

³ Physics Department, IIT Guwahati, Assam, INDIA - 781039

E-mail: s.hesselbach@phys.soton.ac.uk

Abstract. The Minimal Supersymmetric Standard Model (MSSM) with explicit CP violation is studied with the help of the di-photon decay channel of the lightest neutral Higgs boson. Effects of CP violation, entering via the scalar/pseudo-scalar mixing at higher order as well as through the Higgs-sfermion-sfermion couplings at tree-level, are analyzed in the MSSM with and without light sparticles. A light stop may have a strong impact on the decay width and Branching Ratio (BR) of the decay process $H_1 \rightarrow \gamma\gamma$, whereas other light sparticles have only little influence. In some regions of the MSSM parameter space with large CP-violating phase $\phi_\mu \sim 90^\circ$ a light stop can change the BR by more than 50%.

1. Introduction

Supersymmetry (SUSY) is one of the most favoured scenarios of new physics and will be searched for in all possible ways at the upcoming Large Hadron Collider (LHC) at CERN. In the Minimal Supersymmetric Standard Model (MSSM) many parameters can well be complex and thus explicitly break CP invariance inducing CP violation also in the Higgs sector beyond Born approximation [1]. After elimination of unphysical phases and imposing universality conditions at the unification scale two independent phases remain, the phase ϕ_μ of the higgsino mass term μ and a common phase ϕ_{A_f} of the soft trilinear Yukawa couplings A_f in the sfermion sector [2]. Despite the fact that the SUSY phases may be severely constrained by bounds on the Electric Dipole Moments (EDMs), these constraints are rather model dependent and may be evaded in scenarios with heavy first and second generation sfermions, due to cancellations among various contributions to the EDMs or due to additional contributions from lepton flavour violating terms in the Lagrangian, for a review see e.g. [3].

In this contribution we study the di-photon decay mode, $H_1 \rightarrow \gamma\gamma$, of the lightest neutral Higgs boson H_1 , which involves direct, i.e. leading, effects of the SUSY phases through couplings of the H_1 to SUSY particles in the loops as well as indirect, i.e. sub-leading, effects through the scalar/pseudo-scalar mixing yielding the Higgs mass-eigenstate H_1 . In scenarios with heavy SUSY particles, where the CP violation enters solely through the scalar/pseudo-scalar mixing, the SUSY CP phases can result in a strong suppression of the branching ratio (BR) of the decay $H_1 \rightarrow \gamma\gamma$ as well as of the rate of the combined production and decay process $gg \rightarrow H_1 \rightarrow \gamma\gamma$

* Speaker




Figure 1. Mass of the lightest neutral Higgs boson H_1 for $\phi_\mu = 0^\circ$ (solid, red line) and $\phi_\mu = 90^\circ$ (dashed, green line) with $|A_f| = 1.5$ TeV, $|\mu| = 1$ TeV and different values of $\tan\beta$. In the left plot all sparticles are heavy (~ 1 TeV) for $M_{\tilde{U}_3} = M_{\tilde{Q}_3} = M_{\text{SUSY}} = 1$ TeV, whereas in the right plot a light stop (~ 200 GeV) is present for $M_{\tilde{U}_3} = 250$ GeV and $M_{\tilde{Q}_3} = M_{\text{SUSY}} = 1$ TeV.

[4]. Here, we summarize the results of [5, 6] focusing especially on the effects of light SUSY particles on the decay $H_1 \rightarrow \gamma\gamma$. The analysis of the full production and decay process at the LHC is postponed to a forthcoming publication [7].

2. Di-photon Higgs decay in CP-violating MSSM

In order to analyze the Higgs decays in the CP-violating MSSM we have used the publicly available FORTRAN code CPSUPERH [8], version 2, which calculates the mass spectrum and decay widths of all Higgs bosons along with their couplings to SM and SUSY particles. The leading terms in the CP-violating scalar/pseudo-scalar mixing in the Higgs sector are proportional to $\text{Im}(\mu A_f)$, hence we assume $\phi_{A_f} = 0$ and analyze the effects of nonzero ϕ_μ in the following.

A random parameter space scan to study the general behaviour of the $\text{BR}(H_1 \rightarrow \gamma\gamma)$ for non-zero ϕ_μ has revealed that about 50% deviations are possible for M_{H_1} around 104 GeV for $\phi_\mu = 100^\circ$, and an average of 30% deviation occurs over the mass range 90–130 GeV. Furthermore, a strong impact of a light stop \tilde{t}_1 on the BR has been established [5]. In [6] these results have been consolidated by analyzing the details at the matrix element level including the phase dependence of the Higgs mixing matrix elements and of the respective couplings and by performing a more thorough study of the dependence on the SUSY parameters.

In figure 1 the mass of H_1 is shown as a function of the mass of the charged Higgs boson (M_{H^+}) for $\phi_\mu = 0^\circ, 90^\circ$, $\tan\beta = 2, 5, 20, 50$ and the two cases $M_{\tilde{U}_3} = M_{\tilde{Q}_3} = M_{\text{SUSY}} = 1$ TeV, where all SUSY particles are heavy, and $M_{\tilde{U}_3} = 250$ GeV and $M_{\tilde{Q}_3} = M_{\text{SUSY}} = 1$ TeV, where a light \tilde{t}_1 with mass ~ 200 GeV is present. While in the low $\tan\beta$ case the mass shift induced by the change in ϕ_μ from 0° to 90° is about 10%, in the case of $\tan\beta = 20$ or above it is about 1% or less. The sudden shift in the dependence of M_{H_1} on M_{H^+} around $M_{H^+} = 150$ GeV is due to a cross over in the Higgs mass eigenstates at that point, where all three neutral Higgs states are approximately degenerate in mass around 120 GeV.

Figure 2 shows the $\text{BR}(H_1 \rightarrow \gamma\gamma)$ for five representative ϕ_μ values between 0° and 180° as a function of M_{H^+} for the two cases $M_{\tilde{U}_3} = 1$ TeV (all SUSY particles heavy) and $M_{\tilde{U}_3} = 250$ GeV (light \tilde{t}_1). The respective values of M_{H_1} are indicated separately on the horizontal lines above for each ϕ_μ value. Again the cross over point in the Higgs mass eigenstates at $M_{H^+} \sim 150$ GeV is clearly visible. Below this point the BRs are very small and there is a strong ϕ_μ dependence of




Figure 2. BR of $H_1 \rightarrow \gamma\gamma$ for $|A_f| = 1.5$ TeV, $|\mu| = 1$ TeV and $\tan\beta = 20$. Values of M_{H_1} corresponding to representative points on the M_{H^+} axis are indicated on the horizontal lines above separately for the values of ϕ_μ used. The left plot corresponds to the case with $M_{\tilde{U}_3} = 1$ TeV (no light SUSY particles), while the right plot corresponds to the case with $M_{\tilde{U}_3} = 250$ GeV (a light stop is present).

M_{H_1} , hence our analysis is not relevant in this parameter region. Above $M_{H^+} \sim 150$ GeV with $M_{H_1} \gtrsim 115$ GeV the ϕ_μ dependence of M_{H_1} is within the expected experimental uncertainty and the BR is large enough to be important for the LHC Higgs search. In scenarios with heavy SUSY particles (left plot) the BR increases with increasing ϕ_μ leading to a 50% increase for $\phi_\mu = 90^\circ$ at $M_{H^+} \sim 200$ GeV. This ϕ_μ dependence is caused mainly by the ϕ_μ dependence of the H_1 couplings to W^\pm bosons and t and b quarks which appear in the loop-induced decay $H_1 \rightarrow \gamma\gamma$. When a light \tilde{t}_1 is present the additional ϕ_μ dependence in the stop sector causes a considerable change of the ϕ_μ dependence of the BR. First the BR increases again with increasing ϕ_μ up to a maximum for some value of ϕ_μ around 40° , beyond which, however, the BR decreases to about 50% at $\phi_\mu = 180^\circ$.

Concerning the dependence on other SUSY parameters we have found that a smaller value of $|A_f|$ considerably changes the ϕ_μ dependence of the BR in scenarios with light \tilde{t}_1 , whereas a smaller $|\mu|$ value leads generally to a smaller ϕ_μ dependence. Other light SUSY particles have only a negligible effect on the BR.

3. Summary

We have analyzed the BR of the di-photon decay of the lightest Higgs boson in the CP-violating MSSM with complex μ parameter. We have found that the strong ϕ_μ dependence of the BR considerably changes in the presence of a light scalar top \tilde{t}_1 . In general, the BR may be increased or decreased for a non-zero ϕ_μ depending on the SUSY parameter point.

References

- [1] Pilaftsis A 1998 *Phys. Lett.* **B435** 88–100 (*Preprint hep-ph/9805373*)
- [2] Pilaftsis A and Wagner C E M 1999 *Nucl. Phys.* **B553** 3–42 (*Preprint hep-ph/9902371*)
- [3] Olive K A, Pospelov M, Ritz A and Santoso Y 2005 *Phys. Rev.* **D72** 075001 (*Preprint hep-ph/0506106*)
- [4] Choi S Y, Hagiwara K and Lee J S 2002 *Phys. Lett.* **B529** 212–221 (*Preprint hep-ph/0110138*)
- [5] Moretti S, Munir S and Poulou P 2007 *Phys. Lett.* **B649** 206–211 (*Preprint hep-ph/0702242*)
- [6] Hesselbach S, Moretti S, Munir S and Poulou P 2007 *Preprint arXiv:0706.4269* [hep-ph]
- [7] Hesselbach S, Moretti S, Munir S and Poulou P In preparation
- [8] Lee J S *et al.* 2004 *Comput. Phys. Commun.* **156** 283–317 (*Preprint hep-ph/0307377*)