

---

# Newly observed charmed states: the case of $X(3872)$

Stefano Nicotri

Università degli studi di Bari and INFN sezione di Bari,  
via Orabona 4, I-70126, Bari, Italy  
stefano.nicotri@ba.infn.it

**Summary.** I briefly review the properties of the hidden charm meson  $X(3872)$ , discussing some puzzling aspects concerning its theoretical interpretation.

## Introduction

Recently many new mesons have been observed in the open and hidden charm sectors [1, 2]. From the need of theoretical interpretation of those states, the interesting possibility of identifying new “exotic” structures has been pointed out. In the following I discuss the case of  $X(3872)$ , which represents one of the most interesting examples in this sense. In particular I focus the attention on the properties of the radiative decays  $X \rightarrow D\bar{D}\gamma$  [3].

## $X(3872)$

The experimental observations can be summarized as follows:


- the  $X(3872)$  resonance has been found in  $J/\psi\pi^+\pi^-$  distribution by four experiments, both in  $B$  decays ( $B^{-(0)} \rightarrow K^{-(0)}X$ ), both in  $p\bar{p}$  annihilation [4]. The mass is  $M = 3871.9 \pm 0.6$  MeV while the width remains unresolved:  $\Gamma < 2.3$  MeV (90 % CL);
- there is no evidence of resonances in the charged mode  $J/\psi\pi^\pm\pi^0$  or in  $J/\psi\eta$  [5];
- the state is not observed in  $e^+e^-$  annihilation;
- for  $X$  produced in  $B$  decays the ratio  $\frac{B(B^0 \rightarrow K^0 X)}{B(B^+ \rightarrow K^+ X)} = 0.61 \pm 0.36 \pm 0.06$  is obtained [5];
- the dipion spectrum in  $J/\psi\pi^+\pi^-$  is peaked at large mass [6];
- the decay in  $J/\psi\pi^+\pi^-\pi^0$  is observed [7] with  $\frac{B(X \rightarrow J/\psi\pi^+\pi^-\pi^0)}{B(X \rightarrow J/\psi\pi^+\pi^-)} = 1.0 \pm 0.4 \pm 0.3$ : this implies G-parity violation;

- the radiative mode  $X \rightarrow J/\psi\gamma$  is found [7, 8] with  $\frac{B(X \rightarrow J/\psi\gamma)}{B(X \rightarrow J/\psi\pi^+\pi^-)} = 0.19 \pm 0.07$ , therefore charge conjugation of the state is  $C=+1$ ;
- the angular distribution of the final state is compatible with the spin-parity assignment  $J^P = 1^+$  [9];
- there is a signal in  $D^0\bar{D}^0\pi^0$  with  $\frac{B(X \rightarrow D^0\bar{D}^0\pi^0)}{B(X \rightarrow J/\psi\pi^+\pi^-)} = 9 \pm 4$  [10]; the near-threshold  $D^0\bar{D}^0\pi^0$  enhancement in  $B \rightarrow D^0\bar{D}^0\pi^0 K$  has a the peak at  $M = 3875.4 \pm 0.7_{-2.0}^{+1.2}$  MeV with  $B(B \rightarrow KX \rightarrow KD^0\bar{D}^0\pi^0) = (1.27 \pm 0.31_{-0.39}^{+0.22}) \times 10^{-4}$  [11].


All the measurements are compatible with the assignment  $J^{PC} = 1^{++}$ , but if the  $2^3P_1$  is identified with  $Y(3940)$ , another recently observed meson, there is an overpopulation of  $1^{++} c\bar{c}$  mesons. This, together with the observation of the G-parity violating  $X$  transitions suggested the conjecture that  $X(3872)$  is not a charmonium  $\bar{c}c$  state. Indeed, the coincidence between the  $X$  mass as averaged by PDG and the  $D^{*0}\bar{D}^0$  mass inspired the proposal that  $X(3872)$  could be a molecular quarkonium [12], a  $D^{*0}$  and  $\bar{D}^0$  bound state with small binding energy due to a single pion exchange [13]. In this scheme the wave function of  $X(3872)$  is given by [14]:

$$|X(3872)\rangle = a|D^{*0}\bar{D}^0 + \bar{D}^{*0}D^0\rangle + b|D^{*+}D^- + D^{*-}D^+\rangle + \dots \quad (1)$$

(with  $|b| \ll |a|$ ) and so one could explain why this state seems not to have definite isospin, why the mode  $X \rightarrow J/\psi\pi^0\pi^0$  was not found, and why, if the binding mechanism is a single pion exchange (which is in fact a controversial point), there are no  $DD$  molecular states. The opposing remark concerning the large value of the ratio  $\frac{B(X \rightarrow J/\psi\pi^+\pi^-\pi^0)}{B(X \rightarrow J/\psi\pi^+\pi^-)}$  is that the ratio of the amplitudes is small:  $\frac{A(X \rightarrow J/\psi\pi^0\pi^0)}{A(X \rightarrow J/\psi\omega)} \simeq 0.2$ , so that the isospin violating amplitude is 20% of the isospin conserving one, an effect that could be related to the mass difference between neutral and charged  $D$  mesons, considering the contribution of  $DD^*$  intermediate states to  $X$  decays [15, 16]. Moreover, the central value of the mass measured in  $D^0\bar{D}^0\pi^0$  is 4 MeV higher than the PDG value (with a large systematic error), a result difficult to explain in the molecular picture [17]. It has also been suggested that the molecular interpretation would imply that the radiative decay in neutral  $D$  mesons:  $X \rightarrow D^0\bar{D}^0\gamma$  should be dominant with respect to  $X \rightarrow D^+D^-\gamma$  [14]. However, assuming that  $X(3872)$  is an ordinary  $J^{PC} = 1^{++}$  charmonium and describing the  $X(3872) \rightarrow D\bar{D}\gamma$  amplitude by polar diagrams with  $D^*$  and  $\psi(3770)$  as intermediate particles the ratio  $R = \frac{\Gamma(X \rightarrow D^+D^-\gamma)}{\Gamma(X \rightarrow D^0\bar{D}^0\gamma)}$  depends only on the ratio  $c/\hat{g}_1$ ,  $c$  being the parameter governing the radiative  $X\psi(3770)\gamma$  matrix element, and  $\hat{g}_1$  representing the coupling  $XD\bar{D}^*$  (or  $XD^*\bar{D}$ ). The ratio  $R$  is plotted in fig.1 versus  $c/\hat{g}_1$  and it is shown that in any case  $R < 0.7$ . Thus, there is a suppression of the radiative  $X$  decay mode into charged  $D$  mesons with respect to the mode with neutral  $D$ . Moreover, for small values of  $c/\hat{g}_1$  the ratio  $R$  is tiny, so that this seems not a peculiar feature of a molecular structure of  $X(3872)$ .



**Fig. 1.** Ratio of charged  $X \rightarrow D^+D^-\gamma$  to neutral  $X \rightarrow D^0\bar{D}^0\gamma$  decay widths versus the ratio of hadronic parameters  $c/\hat{g}_1$ .



**Fig. 2.** Photon spectrum (in arbitrary units) in  $X \rightarrow D^0\bar{D}^0\gamma$  (top) and  $X \rightarrow D^+D^-\gamma$  (bottom) decays for values of the hadronic parameter  $c/\hat{g}_1 = 1$  (left) and  $c/\hat{g}_1 = 300$  (right).

The photon spectrum in radiative  $X$  decays to both neutral and charged  $D$  meson pairs for two representative values of  $c/\hat{g}_1$ , namely  $c/\hat{g}_1 = 1$  and  $c/\hat{g}_1 = 300$ , is depicted in fig.2. For small values of the parameter  $c/\hat{g}_1$ , i.e. in the condition where the intermediate  $D^*$  dominates the decay amplitude,

the photon spectrum in the  $D^0\bar{D}^0\gamma$  mode essentially coincides with the line corresponding to the  $D^*$  decay at  $E_\gamma \simeq 139$  MeV and width determined by the  $D^*$  width. The narrow peak is different from the line shape expected in a molecular description, which is related to the wave function of the two heavy mesons bound in the  $X(3872)$ , in particular to the binding energy of the system, being broader for larger binding energy. On the other hand, the photon spectrum in the charged  $D^+D^-\gamma$  mode is broader, with a peak at  $E_\gamma \simeq 125$  MeV, the total  $X \rightarrow D^+D^-\gamma$  rate being severely suppressed with respect to the  $X \rightarrow D^0\bar{D}^0\gamma$  one.

At the opposite side of the  $c/\hat{g}_1$  range, where  $\psi(3770)$  gives a large contribution to the radiative amplitude, a peak at  $E_\gamma \simeq 100$  MeV appears both in neutral and charged  $D$  meson modes, in the first case together with the structure at  $E_\gamma \simeq 139$  MeV. This spectrum was described also in [14], where in this case the radiative decay was interpreted as deriving from the  $\bar{c}c$  core of  $X(3872)$ . In this range of parameters the ratio of the  $X \rightarrow D^+D^-\gamma$  to  $X \rightarrow D^0\bar{D}^0\gamma$  rates reaches the largest value.

The experimental determination of the photon spectrum of the type depicted in fig.2, together with the measurement of the  $X \rightarrow D\bar{D}\gamma$  widths is a challenging task. Nevertheless, this measurement is important to shed light on the structure of  $X(3872)$ . Other methods to distinguish between a molecular and a quarkonium description have been proposed, and concern the pionic  $X$  decays to  $\chi_{cJ}$  [18].

## Conclusions

I have summarized the features of  $X(3872)$ , discussing in particular possible tests of the molecular interpretation of this state. I have focused the attention on the radiative decays  $X \rightarrow D\bar{D}\gamma$ , showing that the suppression of the charged channel with respect to the neutral one is not a distinctive feature of the molecular picture. However, in this decay mode the photon spectrum is expected to be different in the  $c\bar{c}$  and molecular description, so that its experimental investigation could be useful to shed more light on this puzzling hadron.

## Acknowledgements

I thank the conference organizers and convenors. I also acknowledge P. Colangelo and F. De Fazio for collaboration. This work has been supported in part by the EU Contract No. MRTN-CT-2006-035482, "FLAVIANet".

## References

- [1] W. M. Yao *et al.* [Particle Data Group], J. Phys. G **33** (2006) 1.
- [2] For reviews see: P. Colangelo *et al.* Mod. Phys. Lett. A **19** (2004) 2083; E. S. Swanson, Phys. Rept. **429** (2006) 243. P. Colangelo *et al.* arXiv:hep-ph/0609240; J. L. Rosner, J. Phys. Conf. Ser. **69** (2007) 012002 and P. Colangelo *et al.* arXiv:0706.0053 [hep-ph].
- [3] P. Colangelo, F. De Fazio and S. Nicotri, Phys. Lett. B **650** (2007) 166.
- [4] S. K. Choi *et al.* [Belle Collaboration], Phys. Rev. Lett. **91** (2003) 262001; D. Acosta *et al.* [CDF II Collaboration], Phys. Rev. Lett. **93** (2004) 072001; V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **93** (2004) 162002; B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **71** (2005) 071103.
- [5] D. Bernard, talk at the International Conference on QCD and Hadronic Physics, June 16-20 2005, Beijing, China.
- [6] A. Abulencia *et al.* [CDF Collaboration], Phys. Rev. Lett. **96** (2006) 102002.
- [7] K. Abe *et al.* [Belle Collaboration], arXiv:hep-ex/0505037.
- [8] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **74** (2006) 071101
- [9] K. Abe *et al.* [Belle Collaboration], arXiv:hep-ex/0505038; A. Abulencia *et al.* [CDF Collaboration]. arXiv:hep-ex/0612053.
- [10] G. Gokhroo *et al.*, [Belle Collaboration], BELLE-CONF-0568 (2005); P. Pakhlov, talk at ICHEP 2006, 26/7-2/8 2006, Moskow, Russia.
- [11] G. Gokhroo *et al.*, Phys. Rev. Lett. **97** (2006) 162002, The signal has been confirmed by BaBar Collaboration, see P. Grenier, arXiv: 0705.2432.
- [12] M. B. Voloshin and L. B. Okun, JETP Lett. **23** (1976) 333.
- [13] F. E. Close and P. R. Page, Phys. Lett. B **578** (2004) 119; M. B. Voloshin, Phys. Lett. B **579** (2004) 316; N. A. Tornqvist, Phys. Lett. B **590** (2004) 209; C. Y. Wong, Phys. Rev. C **69** (2004) 055202; E. Braaten and M. Kusunoki, Phys. Rev. D **69** (2004) 074005; E. S. Swanson, Phys. Lett. B **588** (2004) 189 and **598** (2004) 197; M. T. AlFiky *et al.* Phys. Lett. B **640** (2006) 238; E. Braaten *et al.* arXiv:hep-ph/0702128; S. Fleming *et al.* Phys. Rev. D **76** (2007) 034006.
- [14] M. B. Voloshin, Int. J. Mod. Phys. A **21** (2006) 1239.
- [15] M. Suzuki, Phys. Rev. D **72** (2005) 114013.
- [16] C. Meng and K. T. Chao, Phys. Rev. D **75** (2007) 114002.
- [17] An argument in favour of the possible coexistence of a  $D^0\bar{D}^0\pi^0$  signal with the other features of the molecular description has been provided by C. Hanhart *et al.* Phys. Rev. D **76** (2007) 034007.
- [18] S. Dubynskiy and M. B. Voloshin, arXiv:0709.4474 [hep-ph].