

Ratio of charm to bottom R_{AA} as a test of pQCD vs. AdS/CFT energy loss

W. A. Horowitz

Department of Physics, Columbia University, 538 West 120th Street, New York, NY 10027, USA

Frankfurt Institute for Advanced Studies (FIAS), 60438 Frankfurt am Main, Germany

Abstract. LHC predictions for the charm and bottom nuclear modification factors, $R_{AA}^c(p_T)$ and $R_{AA}^b(p_T)$, using pQCD and AdS/CFT drag energy loss models are given. We show that a new observable, the double ratio $R^{cb}(p_T) = R_{AA}^c(p_T)/R_{AA}^b(p_T)$, allows for easy experimental distinction between the two classes of energy loss models.

The theoretical framework of a weakly-coupled QGP used in pQCD models that quantitatively describe the high- p_T π^0 , η suppression at RHIC is challenged by several experimental observables, not limited to high- p_T only, suggesting the possibility that a strongly-coupled picture might be more accurate. One seeks a measurement that may clearly falsify one or both approaches; heavy quark jet suppression is one possibility. Strongly-coupled calculations, utilizing the AdS/CFT correspondence, have been applied to high- p_T jets in three ways [1, 2, 3]. We will focus on predictions from the AdS/CFT heavy quark drag model and compare them to pQCD predictions from WHDG convolved radiative and elastic energy loss and radiative only energy loss [4]. Comparisons between AdS/CFT calculations and data are difficult. First, one must accept the double conjecture of $QCD \leftrightarrow SYM \leftrightarrow AdS/CFT$. Second, to make contact with experiment, one must make further assumptions to map quantities such as the coupling and temperature in QCD into the SUGRA dual. For example, the AdS/CFT prediction for the heavy quark diffusion coefficient is $D = 4/\sqrt{\lambda}/(2\pi T)$ [2], where $\lambda = g_{SYM}^2 N_c$ is the 't Hooft coupling. The “obvious” first such mapping [5] simply equates constant couplings, $g_s = g_{SYM}$, and temperatures, $T_{SYM} = T_{QCD}$. Using this prescription with the canonical $N_c = 3$ and $\alpha_s = .3$ yields $D \approx 1.2/(2\pi T)$. It was claimed in [2] that $D = 3/(2\pi T)$ agrees better with data; this requires $\alpha_s \approx .05$. An “alternative” mapping [5] equates the quark-antiquark force found on the lattice to that computed using AdS/CFT, giving $\lambda \approx 5.5$, and the QCD and SYM energy densities, yielding $T_{SYM} = T_{QCD}/3^{1/4}$. The medium density to be created at LHC is unknown; we will take the PHOBOS extrapolation of $dN_g/dy = 1750$ and the KLN model of the CGC, $dN_g/dy = 2900$, as two sample values. We will search for general trends associated with AdS/CFT drag (denoted hereafter simply as AdS/CFT) or pQCD as the aforementioned uncertainties mean little constrains the possible normalizations of AdS/CFT R_{AA}^Q predictions for LHC.

The AdS/CFT derivation of the drag on a heavy quark yields $dp_T/dt = -\mu_Q p_T = -(\pi\sqrt{\lambda}T_{SYM}^2/2m_Q)p_T$ [3], giving an average fractional energy loss of $\bar{\epsilon} = 1 - \exp(-\int dt \mu_Q)$. Asymptotic pQCD energy loss for heavy quarks in a static medium goes as $\bar{\epsilon} \approx \kappa \alpha_s L^2 \hat{q} \log(p_T/m_Q)/p_T$, where κ is a proportionality constant and L is the pathlength traversed by the heavy quark. Note that AdS/CFT fractional momentum loss is independent of momentum while pQCD loss decreases with jet




Figure 1. (Color Online) (a) Charm and bottom $R_{AA}(p_T)$ predictions for LHC. The generic trend of pQCD curves increasing with p_T while AdS/CFT curves decrease is seen for representative input parameters; similar trends occurred for the other input possibilities considered. (b) Double ratio of charm to bottom $R_{AA}(p_T)$. pQCD and AdS/CFT drag curves fall into two distinct groups; the LHC should easily distinguish between the two trends.

energy. The heavy quark production spectrum may be approximated by a slowly varying power law of index $n_Q(p_T) + 1$, then $R_{AA}^Q \approx (1 - \bar{\epsilon})^{n_Q(p_T)}$. Since $n_Q(p_T)$ is a slowly increasing function of momentum, we expect $R_{AdS}^Q(p_T)$ to decrease while $R_{pQCD}^Q(p_T)$ to increase as momentum increases. This behavior is reflected in the full numerical calculations shown in Fig. 1 (a); details of the model can be found in [6].

For large opacity pQCD predicts nearly flat R_{AA}^Q , masking the difference discussed above. One can see in Fig. 1 (b) that the separation of AdS/CFT and pQCD predictions is enhanced when the double ratio of charm to bottom nuclear modification, $R^{cb}(p_T) = R_{AA}^c(p_T)/R_{AA}^b(p_T)$, is considered. Asymptotic pQCD energy loss goes as $\log(m_Q/p_T)/p_T$, becoming insensitive to quark mass for $p_T \gg m_Q$; hence $R_{pQCD}^{cb} \rightarrow 1$. Expanding the R_{AA} formula for small ϵ yields $R_{pQCD}^{cb}(p_T) \approx 1 - p_{cb}/p_T$, where $p_{cb} = \kappa \alpha_s n(p_T) L^2 \log(m_b/m_c) \hat{q}$ and $n_c \approx n_b = n$. Therefore the ratio approaches unity more slowly for larger suppression. This behavior is reflected in the full numerical results for the moderately quenched pQCD curves, but is violated by the highly oversuppressed $\hat{q} = 100$ curve. The AdS/CFT drag, however, is independent of p_T . Approximating the medium with a static plasma of thickness L gives $R_{AA}^Q \approx \int_0^L d\ell \exp(-n_Q \mu_Q \ell) \approx 1/n_Q \mu_Q L$ which yields $R^{cb}(p_T) \approx n_b(p_T) m_c / n_c(p_T) m_b \approx m_c/m_b \approx .27$. This behavior is also reflected in the full numerical results shown in Fig. 1 (b), and so, remarkably, the pQCD and AdS/CFT curves fall into easily distinguishable groups, robust to changes in input parameters. An estimate for the momentum after which corrections to the above AdS/CFT drag formula are needed, $\gamma > \gamma_c$, found in the static string geometry is $\gamma_c = 1/1 + (2m_Q/T\sqrt{\lambda})$ [7]. Since temperature is not constant we show the smallest speed limit, using $T(\tau_0, \vec{x} = \vec{0})$, and largest, from T_c , represented by “O” and “|,” respectively. A deviation of R^{cb} away from unity at LHC in year 1 would pose a serious challenge to the usual pQCD paradigm. An observation of a significant increase in R^{cb} with jet momenta would imply that the current AdS/CFT picture is only applicable at low momenta, if at all. For a definitive statement to be made a $p + Pb$ control run will be crucial.

- [1] H. Liu, K. Rajagopal and U. A. Wiedemann, Phys. Rev. Lett. **97**, 182301 (2006).
- [2] J. Casalderrey-Solana and D. Teaney, Phys. Rev. D **74**, 085012 (2006).
- [3] C. P. Herzog, A. Karch, P. Kovtun, C. Kozcaz and L. G. Yaffe, JHEP **0607**, 013 (2006); S. S. Gubser, Phys. Rev. D **74**, 126005 (2006).
- [4] S. Wicks, W. Horowitz, M. Djordjevic and M. Gyulassy, Nucl.Phys. **A784** 426 (2007).
- [5] S. S. Gubser, arXiv:hep-th/0611272.
- [6] W. A. Horowitz and M. Gyulassy, arXiv:0706.2336 [nucl-th].
- [7] S. S. Gubser, arXiv:hep-th/0612143.