

# NON-STATISTICAL $\gamma$ RAYS FROM FRAGMENTS

F.F.Karpeshin

*St. Petersburg University, Institute of Physics*  
SU-198504 St. Petersburg, Russia

February 9, 2022

## 1 Introduction

Observation of the shake effects brought about by the neck rupture in fission is of great interest. From mathematical view point, the rupture means break-down of the analyticity of the Hamiltonian with respect to time. Thus, muon shake in muon-induced prompt fission manifests itself in muonic conversion. The calculated probability agrees with the experiment [1, 2]. Herein we calculate the probability of emission of  $\gamma$  quanta, the results are also of interest in connection with experiments [3, 4], in which works non-statistical  $\gamma$  rays from  $^{252}\text{Cf}$  spontaneous fission are under investigation.

The process of snapping-back of the nuclear surface gives rise to the oscillations of the surface, whose lifetime is determined by the dissipation. It can be evaluated as  $\tau_{diss} \approx 10^{-19}$  s [5]. The oscillations generate nonstationary electromagnetic field in space. That causes electromagnetic processes of internal conversion and  $\gamma$  radiation.

The nuclear vibrations can be considered like the motion of a classical droplet. Write down the conventional expansion of the nuclear form in spherical harmonics

$$R(\theta, \phi) = R_0 \left( 1 + \beta_0 + \sum_{\lambda, \mu} \beta_{\lambda, \mu} Y_{\lambda, \mu}(\theta, \phi) \right). \quad (1)$$

Main properties of the fragments can be described by a lowering for the quadrupole and octupole terms. Such superposition leads to a pear-like form of the nucleus. It is essential that the electric dipole term in this case must be included in eq. (1) to keep the centre of mass fixed [6], the relation  $\beta_1 = -0.743\beta_2\beta_3$  following from the latter condition [1, 7]. The other consequence is the appearance of the polarization electric dipole moment in the nucleus [6]:

$$d \equiv D/e = -\kappa\beta_2\beta_3. \quad (2)$$

Polarizability  $\lambda$  (or  $\kappa$ ) in eq. (2) can be evaluated e.g. from formulae [6, 7], which agree with experiment (see also other refs. in [7]).

Considering the oscillations quasiclassically and taking into account the relaxation, put down

$$\beta_i(t) = \beta_i^{(0)} \sin \omega_i t \exp(-\gamma_i t/2), \quad i = 2, 3. \quad (3)$$

Then the spectral density of the radiated energy is given in the classical limit [8] by the following expression:

$$d\mathcal{E}_\omega = \frac{4}{3} |\ddot{D}_\omega|^2, \quad (4)$$

where  $D_\omega$  is the Fourier transform of the second derivative of  $D(t)$  with respect to time. Using (2) and (3) in eq. (4), we find

$$\begin{aligned} \ddot{D}_\omega &= -\kappa \beta_2^{(0)} \beta_3^{(0)} \int_0^\infty \exp(i\omega t) \frac{d^2}{dt^2} \sin \omega_2 t \sin \omega_3 t \exp(-\gamma t/2) dt = \\ &= i D_0 \frac{(\omega_2 + \omega_3)^2}{\omega - \omega_2 - \omega_3 + i\frac{\gamma}{2}}, \end{aligned} \quad (5)$$

where  $\gamma = \gamma_2 + \gamma_3$  is the total quenching, and  $D_0 = -\kappa \beta_2^{(0)} \beta_3^{(0)}$ .

Supposing  $\beta_2^{(0)} \approx \beta_3^{(0)} \approx 0.7$  [5], we calculate by means of formulae [7]  $d_0 \approx 5$  Fm. Using then the LDM values for a representative heavy fragment  $^{140}\text{Xe}$ , which are  $\hbar\omega_2 = 2.2$  MeV,  $\hbar\omega_3 = 2.8$  MeV, and evaluating  $\gamma$  from the lifetime  $\tau_{diss} = \gamma^{-1} = 10^{-19}$  s, as it is stated previously, one immediately finds by means of eq. (4)

$$N_\gamma = \int_0^\infty \frac{d\mathcal{E}_\omega}{\hbar\omega} \approx 8 \cdot 10^{-3} \text{ fission}^{-1}. \quad (6)$$

We conclude that this value is in qualitative agreement with experiment [3, 4], taking into account the uncertainties connected with the value of  $\tau_{diss}$ . For the value supposed,  $\tau_{diss} \approx 10^{-19}$  s, the contribution of the proposed mechanism is enough to explain the experimental value.

On the other hand, we see that this contribution is proportional to  $\tau_{diss}$ . Therefore, study of non-statistical  $\gamma$  rays from fission gives direct information about dissipation in large-amplitude collective motion represented by postrupture oscillations in the fragments.

## References

- [1] F.F.Karpeshin. Z. Phys. **A344**, 55 (1992); Yad. Fiz., 1992, 55, 2893.
- [2] G. Ye.Belovitsky et al. In: "Fiftirth Anniversary or Nuclear Fission", Proc. Intern. Conf., St. Petersburg, 1989. V. 1, P. 313.
- [3] H. van der Ploeg et al. Phys. Rev. Lett., 1992, **68**, 3145; KVI annual report, 1991, p. 17.
- [4] A.Wiswesser e.a. GSI annual report, 1991, P. 79.
- [5] Yu. P. Gangrsky, B. N.Markov, V. P. Perelygin. Registratsia i spektrometria oskolkov delenia. Moscow: Energoizdat, 1992 (in Russian) (Registration and Spectrometry of the fission fragments)
- [6] V.M.Strutinsky. At. Energ., 1956, No. 4, p. 150.
- [7] V.Yu.Denisov. Yad. Fiz. 1992, **55**, 2647; 1989, **49**, 644.
- [8] L.D.Landau, E.M. Lifshitz. Teoria Polya, Moscow: Nauka, 1973. (*In Russian*) (Theory of Field)