

# CP violation and final state interactions in $B \rightarrow K\pi\pi$ decays

L. Leśniak<sup>1</sup>, B. El-Bennich<sup>2</sup>, A. Furman<sup>3</sup>, R. Kamiński<sup>1</sup>, B. Loiseau<sup>2</sup>  
and B. Moussallam<sup>4</sup>

<sup>1</sup> The Henryk Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences, 31-342 Kraków, Poland

<sup>2</sup> Laboratoire de Physique Nucléaire et de Hautes Énergies (IN2P3-CNRS-Universités Paris 6 et Paris 7), Groupe Théorie, Université Pierre et Marie Curie, 4 Pl. Jussieu, F-75252 Paris, France

<sup>3</sup> ul. Bronowicka 85/26, 30-091 Kraków, Poland

<sup>4</sup> Institut de Physique Nucléaire, Université Paris-Sud, F-91406 Orsay, France

E-mail: Leonard.Lesniak@ifj.edu.pl

## Abstract.

Effects of CP violation and of final state interactions between pairs of pseudoscalar mesons are studied in three-body  $B^+$ ,  $B^-$ ,  $B^0$  and  $\bar{B}^0$  decays into  $K\pi\pi$ . An alternative approach to the isobar model for three-body B decays is proposed. It is based on the QCD factorization approximation and the knowledge of the meson-meson form factors. Some phenomenological charming penguin amplitudes are needed to describe the branching fractions, direct CP asymmetries of the quasi-two-body  $B \rightarrow K^*(892)\pi$  and  $B \rightarrow K_0^*(1430)\pi$  decays as well as the  $K\pi$  effective mass and the helicity angle distributions. The experimental branching fractions for the  $B \rightarrow K_0^*(1430)\pi$  decay, obtained by the Belle and BaBar collaborations using the isobar model, are larger than our predictions by about 52 per cent.

## 1. Introduction

CP violation in the charged B meson decays has been discovered for the first time in the three body charmless hadronic reaction  $B \rightarrow \rho(770)^0 K$ ,  $\rho(770)^0 \rightarrow \pi^+\pi^-$  [1, 2]. Data on three body  $B$  decays are commonly analyzed using the isobar model which violates unitarity and requires many free parameters. In [3] we have developed a model with a unitarized description of final state interactions between the two pions. We found interesting interferences between the  $\rho(770)^0$  and  $f_0(980)$  resonances. Within a similar approach and for the  $K\pi$  effective masses smaller than about 1.8 GeV, we study another part of the Dalitz plots corresponding to the  $B^\pm$ ,  $B^0$  and  $\bar{B}^0$  decays where the strange resonances  $K^*$  are observed. Among them, one encounters two resonances  $K^*(892)$  and  $K_0^*(1430)$  formed in the  $K\pi$   $P$ - and  $S$ - waves, respectively. These two resonances dominate the pion-kaon vector and scalar form factors which play an essential role in the description of the decay amplitudes. The strange form factors are determined from unitary coupled-channel equations using the experimental data on the  $\pi K$  phase shifts and inelasticities and some constraints from chiral perturbation theory.

## 2. Decay amplitudes for $B \rightarrow K\pi\pi$ reactions

The amplitudes for the  $B \rightarrow K\pi\pi$  decays consist of two terms. The first part corresponds to the quark weak transitions  $b \rightarrow s\bar{d}d$  or  $b \rightarrow s\bar{u}u$  in the QCD factorization approximations. The second one is a phenomenological long-distance amplitude with  $c$ - or  $u$ -quarks in loop. Such an amplitude with the  $c$ -quark in loop, called the charming penguin term [4], could be related to  $B$  decay processes with intermediate  $D_s^{(*)}D^{(*)}$  states. For example, the  $S$ -wave part of the  $B^- \rightarrow K^- \pi^+ \pi^-$  decay amplitude reads:

$$A_S = \frac{G_F}{\sqrt{2}} (M_B^2 - m_\pi^2) \frac{m_K^2 - m_\pi^2}{q^2} F_0^{B \rightarrow \pi}(q^2) f_0^{K^- \pi^+}(q^2) \times \left\{ \lambda_u(a_4^u + P_u - a_{10}^u/2) + \lambda_c(a_4^c + P_c - a_{10}^c/2) - \frac{2q^2}{(m_b - m_d)(m_s - m_d)} \times [\lambda_u(a_6^u + S_u - a_8^u/2) + \lambda_c(a_6^c + S_c - a_8^c/2)] \right\}. \quad (1)$$

Here  $q^2$  denotes the  $K^- \pi^+$  effective mass squared,  $F_0^{B \rightarrow \pi}(q^2)$  is the scalar  $B$  to  $\pi$  form factor,  $f_0^{K^- \pi^+}(q^2)$  is the scalar  $K^- \pi^+$  form factor and the remaining symbols are fully explained in ref. [3]. The  $P$ -wave amplitude, proportional to the vector form factors  $F_1$  and  $f_1$ , is given by

$$A_P = 2\sqrt{2}G_F \mathbf{p}_{\pi^-} \cdot \mathbf{p}_{\pi^+} F_1^{B \rightarrow \pi}(q^2) f_1^{K^- \pi^+}(q^2) [\lambda_u(a_4^u + P_u - a_{10}^u/2) + \lambda_c(a_4^c + P_c - a_{10}^c/2)]. \quad (2)$$


The long-distance penguin amplitudes are parametrized by four complex parameters  $S_u, S_c, P_u$  and  $P_c$ .

## 3. Comparison with experimental results

We analyse the experimental data of the Belle and BaBar collaborations [1, 2, 5, 6] for the four reactions:  $B^- \rightarrow (K^- \pi^+) \pi^-$ ,  $B^+ \rightarrow (K^+ \pi^-) \pi^+$ ,  $\bar{B}^0 \rightarrow (\bar{K}^0 \pi^-) \pi^+$  and  $B^0 \rightarrow (K^0 \pi^+) \pi^-$ . The physical observables fitted in our model include 12 values of the branching fractions and direct CP-violating asymmetries for the  $B \rightarrow K^*(892)^0 \pi$  and  $B \rightarrow K_0^*(1430)^0 \pi$  decays in different charge combinations and 249 data points for the  $K\pi$  effective mass and helicity angle distributions. If the charming penguin parameters are not included in the fit then the theoretical values of the branching fractions for these decays are too small by factors ranging between 2.3 and 3.6. The results without the charming penguin terms are in agreement with those recently calculated by Cheng, Chua and Soni [7]. We have then performed two separate fits with 4 penguin parameters: in the first fit the experimental branching fractions for the  $B \rightarrow K_0^*(1430)^0 \pi$  decays have been included while in the second fit they have been omitted. The fit was much better in the second case.

Typical results of the  $K\pi$  effective mass distributions for these two cases are shown in Fig. 1(a) and 1(b), respectively. In Fig. 1(a) the theoretical curve underestimates the narrow maximum of the  $K^*(892)$  resonance and overestimates the wide maximum due to the  $K_0^*(1430)$  resonance. On the contrary, in Fig. 1(b) both maxima are well reproduced. The average branching fractions and asymmetries for the second fit are given in Table 1. The CP asymmetries are in a general agreement with data. The theoretical value of the  $B^\pm \rightarrow K^*(892)^0 \pi^\pm$  branching fraction is close to the Belle and BaBar results, however the corresponding  $B^\pm \rightarrow K_0^*(1430)^0 \pi^\pm$  value, equal to  $12.9 \pm 0.6$ , is substantially smaller than the experimental values. One should stress the fact that these experimental numbers are *not* directly measured. They are calculated by both collaborations using the isobar model. We calculate our branching fractions simply by integrating the  $K\pi$  effective mass distributions over the energy range specified in Table 1. The published experimental branching fractions correspond to the full range of the  $K\pi$  effective masses. In Table 1, their values are renormalized for the  $K^*(892)$  and  $K_0^*(1430)$  by 0.83 and 0.81, respectively. The renormalized average branching fractions for the  $B^\pm \rightarrow K_0^*(1430)^0 \pi^\pm$  decays

are about 52 % larger than the result of our model fitted to the same  $K\pi$  effective mass and helicity angle data. The published experimental branching fractions are then too large by the same amount. A partial explanation of this fact can rely on the presence of a phenomenological nonresonant part of the decay amplitudes used in the framework of the isobar model. In our approach both resonant and nonresonant parts are included in the strange  $K\pi$  scalar form factor.



**Figure 1.** Averaged  $K\pi$  effective mass distributions for the  $B^\pm \rightarrow K^\pm \pi^\mp \pi^\pm$  decays. The dotted line corresponds to the  $P$ -wave, the dashed line to the  $S$ -wave and the solid one to the full result. The experimental  $B \rightarrow K_0^*(1430)^0 \pi$  values of the branching fractions are included in the fit shown in (a) but not included in that shown in (b). Data are taken from [2].

**Table 1.** Average branching fractions  $\text{Br} \cdot 10^6$  and direct CP asymmetries  $A_{CP} \cdot 10^2$  corresponding to the fit shown in Fig. 1(b).

obs.	channel	$K\pi$ mass range [GeV]	our model	Belle	BaBar
Br	$K^*(892)^0 \pi^\pm$	0.82 – 0.97	$6.14 \pm 0.15$	$5.35 \pm 0.59$	$7.46 \pm 0.81$
$A_{CP}$	$K^*(892)^0 \pi^\pm$	0.82 – 0.97	$-8.0 \pm 2.4$	$-14.9 \pm 6.8$	$6.8 \pm 10.4$
Br	$K_0^*(1430)^0 \pi^\pm$	1.00 – 1.76	$12.9 \pm 0.6$	$25.9 \pm 2.5$	$27.5 \pm 2.2$
$A_{CP}$	$K_0^*(1430)^0 \pi^\pm$	1.00 – 1.76	$-0.4 \pm 1.3$	$7.6 \pm 4.5$	$-6.4 \pm 4.0$

## Acknowledgments

This work has been performed within the framework of the IN2P3-Polish Laboratories Convention (Project No. CSI-12) and within an agreement between the CNRS (France) and the Polish Academy of Sciences (Project No. 19481).

## 4. References

- [1] Aubert B et al. (BaBar Coll.) 2005 *Phys. Rev. D* **72** 072003
- [2] Garmash A et al. (Belle Coll.) 2006 *Phys. Rev. Lett.* **96** 251803
- [3] El-Bennich B, Furman A, Kamiński R, Leśniak L and Loiseau B 2006 *Phys. Rev. D* **74** 114009
- [4] Ciuchini M, Franco E, Martinelli G and Silvestrini L 1997 *Nucl. Phys. B* **501** 271
- [5] Garmash A et al. (Belle Coll.) 2007 *Phys. Rev. D* **75** 012006
- [6] Aubert B et al. (BaBar Coll.) 2006 *Phys. Rev. D* **73** 031101(R)

- [7] Cheng H-Y, Chua C-K and Soni A 2007 Charmless three-body decays of  $B$  mesons *Preprint* arXiv:0704.1049 [hep-ph]