

The scalar unparticle effect on the charged lepton electric dipole moment

E. O. Iltan *

Physics Department, Middle East Technical University
Ankara, Turkey

Abstract

We study the charged lepton electric dipole moment which is induced by the scalar unparticle mediation and we predict the appropriate range for the free parameters appearing in the effective lagrangian which drives the unparticle-standard model lepton interactions. We observe that the charged lepton electric dipole moment is strongly sensitive to the scaling dimension d_u of the unparticle and the new couplings in the effective interaction. Furthermore, we see that the current experimental limits of charged lepton electric dipole moments can ensure an appropriate range for these free parameters.

*E-mail address: eiltan@newton.physics.metu.edu.tr

1 Introduction

Scale invariance has been studied in various disciplines of physics and the scale invariant field theories has been investigated extensively. However, it is not clear that how such possible scale invariant sector comes out in the experiments and how it is determined. Recently, Georgi [1] proposed a hypothetical non-trivial scale invariant sector, having a non-trivial infrared fixed point. It is a new one beyond the standard model (SM) and has the scaling dimension d_u . Georgi suggest that at low energy, around $\Lambda_U \sim 1 TeV$, this sector appears as so called unparticle stuff with non-integer scaling dimension d_u and it looks like a number of d_u massless particles which are invisible. The properties of the suggested matter, the unparticle, are different from the properties of the particle [1, 2]. The unparticles couple to the SM fields and the interactions are described by the effective lagrangian in the low energy level. This approach makes it possible to test the effects of the possible scale invariant sector experimentally with the help of the unparticle idea. The missing energies at various processes which can be measured at LHC or e^+e^- colliders, the various decays and the dipole moments of fundamental particles (that can occur with the unparticle mediation theoretically) are among the possible candidates to search the unparticles and their properties. There is an extensive phenomenological work done in the literature to search the possible effects of the unparticles [2]-[65]: the effects on the missing energy of many processes, the anomalous magnetic moments, $D^0 - \bar{D}^0$ and $B^0 - \bar{B}^0$ mixing, lepton flavor violating interactions, direct CP violation in particle physics; the phenomenological implications in cosmology and in astrophysics.

In this work we study the contribution of the scalar unparticle to the electric dipole moments (EDMs) of the charged leptons. The fermion EDMs are worthwhile to examine since their existence is the sign for the CP violation which is among the most interesting physical phenomena. There are number of experimental and theoretical studies on fermion EDMs in the literature. The current experimental results on the lepton EDMs are, $d_e = (1.8 \pm 1.2 \pm 1.0) \times 10^{-27} e cm$ [66], $d_\mu = (3.7 \pm 3.4) \times 10^{-19} e cm$ [67] and $d_\tau = (3.1) \times 10^{-16} e cm$ [68], respectively. From the theoretical point of view, for non-zero fermion EDMs, one needs a CP violating phase, which is carried by the complex Cabibo Kobayashi Maskawa (CKM) (the lepton mixing matrix) in the quark (lepton) sector, in the SM. Since the theoretical values of the fermion EDMs in the SM are negligibly small (see for example [69]-[72]), one investigates new models beyond, in order to obtain the numerical values not far from the experimental limits. The lepton EDMs have been studied theoretically in various models [73]-[79]; in the framework of the seesaw model [73], in the two Higgs doublet model (2HDM) [74], in the framework of the SM with the inclusion of

non-commutative geometry [75], with the inclusion of non-universal extra dimensions [76], in the split fermion scenario [77, 78], in the Randall Sundrum background [79].

In the present work, we consider that the scalar unparticle mediation is responsible for the CP violating interaction which leads to the charged lepton EDM and we predict the appropriate range for the free parameters appearing in the effective lagrangian which drive the unparticle-SM lepton interactions. We observe that the charged lepton EDMs are strongly sensitive to the scaling dimension d_u and the new unparticle-SM lepton couplings. Furthermore, the current experimental limits of charged lepton EDMs can ensure an appropriate range for these free parameters.

The paper is organized as follows: In Section 2, we present the contribution of the scalar unparticle to the charged lepton EDMs. Section 3 is devoted to discussion and our conclusions.

2 Electric dipole moments of charged leptons induced by the scalar unparticle

The fermion EDM is among the most powerful tools to search the CP violation since it is driven by the CP violating fermion-fermion-photon effective interaction. The CP violation is carried by the complex phase of the CKM matrix (a possible lepton mixing matrix) in the quark (lepton) sector, in the SM. However, their theoretical values are the extremely small in the SM since they exist at beyond two loop orders and the predictions in this model are far from the current experimental limits (see for example [69]-[72]). In the present work, we consider that the unparticle stuff is also responsible for the CP violating interaction which stimulates the lepton EDMs. The unparticle idea has been created by Georgi [1, 2] and this idea is based on the very high energy level interaction of two sectors, the SM and the ultraviolet sector with non-trivial infrared fixed point. Around $\Lambda_U \sim 1 TeV$ this sector manifest itself as new degrees of freedom, called unparticles, being massless and having non integral scaling dimension d_u . Finally, at low energies the related interactions might be driven by the effective field theory and the corresponding effective Lagrangian might be

$$\mathcal{L}_{eff} \sim \frac{\eta}{\Lambda_U^{d_{tot}}} O_{SM} O_U, \quad (1)$$

where $d_{tot} = d_u + d_{SM} - n$ with the space-time dimension n . Here the parameter η is related to the energy scale of ultraviolet sector, the low energy one and the matching coefficient (see [1, 2, 6] for details). Since the operators with lowest possible dimension have the most powerful effect in the low energy effective theory, one chooses the appropriate operators O_U which have

all possible Lorentz structures (see for example [12]).

Our aim is to study the charged lepton EDMs and we need the effective interaction among the unparticle and the SM leptons in order to construct the dipole type transition. The lowest order effective interaction we are interested in is:

$$\mathcal{L} = \frac{\eta_{ij}^S}{\Lambda^{d_u-1}} \bar{l}_{iL} l_{jR} O_U + h.c. , \quad (2)$$

where L and R denote chiral projections $L(R) = 1/2(1 \mp \gamma_5)$, l is the lepton field, O_U is the scalar unparticle operator. The couplings η_{ij}^S are free parameters, complex in general and the scalar (pseudoscalar) λ_{ij}^S (λ_{ij}^P) coupling reads

$$\lambda_{ij}^{S(P)} = \frac{1}{2} (\eta_{ij}^S \pm (\eta_{ij}^S)^*) . \quad (3)$$

This is the case that a possible new source for the CP violating interaction emerges (see Discussion section) and the EDM of the charged leptons due to the unparticle stuff, here the scalar unparticles, appear.

The effective EDM interaction for a charged lepton l is

$$\mathcal{L}_{EDM} = i d_l \bar{l} \gamma_5 \sigma^{\mu\nu} l F_{\mu\nu} , \quad (4)$$

where $F_{\mu\nu}$ is the electromagnetic field tensor, ' d_l ' is the EDM of the charged lepton which is a real number by hermiticity. The possible CKM type lepton mixing matrix causes the non-zero EDMs for the charged leptons, however, here, we consider only the effective interaction of the SM leptons and the unparticle operator to create the CP violation, in order to make predictions of the acceptable range for the free parameters of the unparticle stuff, by using the experimental current limits. From the theoretical point of view, the lepton EDM arises at least in the one loop level with the mediation of the scalar unparticle, in the present case (see Fig.1). The scalar unparticle propagator is obtained with the help of the scale invariance and using the two point function of the unparticle, it is obtained as [2, 3]

$$\int d^4x e^{ipx} \langle 0|T(O_U(x) O_U(0))0 \rangle = i \frac{A_{d_u}}{2\pi} \int_0^\infty ds \frac{s^{d_u-2}}{p^2 - s + i\epsilon} = i \frac{A_{d_u}}{2 \sin(d_u \pi)} (-p^2 - i\epsilon)^{d_u-2} , \quad (5)$$

with

$$A_{d_u} = \frac{16 \pi^{5/2}}{(2\pi)^{2d_u}} \frac{\Gamma(d_u + \frac{1}{2})}{\Gamma(d_u - 1) \Gamma(2d_u)} . \quad (6)$$

For $p^2 > 0$, the function $\frac{1}{(-p^2 - i\epsilon)^{2-d_u}}$ in eq. (5) becomes

$$\frac{1}{(-p^2 - i\epsilon)^{2-d_u}} \rightarrow \frac{e^{-i d_u \pi}}{(p^2)^{2-d_u}} , \quad (7)$$

with a non-trivial phase as a result of non-integral scaling dimension.

Now, we would like to present the EDMs of charged leptons l , driven by the scalar unparticle mediation:

$$d_l = -i \frac{e}{32 \pi^2} \frac{A_{d_u}}{\Lambda^{2d_u-2} \sin(d_u \pi)} \sum_{i=1}^3 \left\{ \lambda_{il}^S ((\lambda_{il}^P)^* - \lambda_{il}^P) Q_i m_i \int_0^1 dx \int_0^{1-x} dy f(x, y, m_i, m_l) \right\}, \quad (8)$$

where subscript i denotes the internal charged leptons, $i = e, \mu, \tau$, Q_i is the charge of i^{th} lepton and $f_1(x, y, m_i, m_l)$ is the function given by

$$f(x, y, m_i, m_l) = \frac{(1-x-y)^{1-d_u} (x+y)}{\left((x+y) (m_l^2 (1-x-y) - m_i^2) \right)^{2-d_u}}. \quad (9)$$

Notice that both scalar and pseudoscalar couplings of unparticles to leptons should be considered to obtain non-zero EDM.

3 Discussion

This section is devoted to the analysis of the possible effects of the scalar unparticle on the charged lepton EDMs and the prediction of the scale dimension of the unparticle by considering the current experimental limits of the EDMs. The non-zero EDM of a fermion is the evidence for the existence of the CP violation since it is driven by the CP violating fermion-fermion-photon effective interaction. The source of the CP violating phase in the SM is the complexity of the CKM matrix in the quark sector and the possible lepton mixing matrix in the lepton sector with non-zero neutrino masses. In the present work, we assume that there exist a new CP phase that arises from the complex couplings, appearing in the tree level fermion-fermion-unparticle interaction, in the effective theory and we investigate its contribution to the charged lepton EDMs. Here the couplings, the CP phase and the scale dimension of the unparticle(s) are the free parameters which should be restricted.

- The scale dimension d_u is restricted in the range $1 < d_u < 2$. Here, $d_u > 1$ is due to the non-integrable singularities in the decay rate [2] and $d_u < 2$ is due to the convergence of the integrals [20].
- Here, we take the coupling η_{ij}^S in eq.(2) as $\eta_{ij}^S = e^{i\theta} |\eta_{ij}^S|$ where θ is the CP violating parameter. In terms of η_{ij}^S the scalar and pseudoscalar couplings read $\lambda_{ij}^S = \frac{1}{2}(\eta_{ij}^S + (\eta^S)_{ij}^*)$ and $\lambda_{ij}^P = \frac{1}{2}(\eta_{ij}^S - (\eta^S)_{ij}^*)$ and, finally, we take $\lambda_{ij}^S = \lambda_{ij} \cos \theta$ and $\lambda_{ij}^P = i \lambda_{ij} \sin \theta$. For the CP parameter θ we take an intermediate value in our numerical calculations.

- We assume that the diagonal couplings λ_{ii} are aware of the lepton family and obey the hierarchy, $\lambda_{\tau\tau} > \lambda_{\mu\mu} > \lambda_{ee}$, and the off-diagonal couplings, $\lambda_{ij}, i \neq j$ are family blind and universal, obeying the equality $\lambda_{ij} = \kappa\lambda_{ee}$ where $\kappa < 1$. Here we choose $\kappa = 0.5$ and take the numerical values of diagonal couplings so that the experimental current limits of charged lepton EDMs can be reached.

Now, we start to predict the magnitudes of the charged lepton EDMs and to estimate the appropriate range for the numerical value of the scaling dimension d_u by using the current experimental limits of the EDMs. Here, we consider that the CP violating phase is carried by the unparticle-charged lepton couplings so that the charged lepton EDM arises with the unparticle mediation.

In Fig.2, we present the contribution of the scalar unparticle to the EDM d_e with respect to the scale parameter d_u for the energy scale $\Lambda_u = 10 TeV$, the coupling $\lambda_{ee} = 0.01$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for total (diagonal, off-diagonal) part. It is observed that there is a strong sensitivity to the scale d_u and the decrease of d_u causes a sharp enhancement in the EDM d_e . The diagonal (off-diagonal) part becomes strong for the small (large) values of d_u and the total contribution reaches the experimental current value for the range of the scale d_u , $1.6 \leq d_u \leq 1.8$.

Fig.3 is devoted to the EDM d_e with respect to the energy scale Λ_u , for the coupling $\lambda_{ee} = 0.01$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.4, d_u = 1.6, d_u = 1.8$, including diagonal and off diagonal terms. With the increasing value of the energy scale Λ_u the EDM decreases and its sensitivity becomes weak, as expected.

Fig.4 represents the EDM d_e with respect to the coupling $\lambda = \lambda_{ee}$ for the energy scale $\Lambda_u = 10 TeV$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.4, d_u = 1.6, d_u = 1.8$, including diagonal and off diagonal terms. This figure shows that the EDM is strongly sensitive to the coupling λ as expected and, even for its small values of the order of 10^{-4} , it is possible to reach the experimental current limit, for the scale dimension $d_u \sim 1.4$.

Fig.5 is devoted to the contribution of the scalar unparticle to the EDM d_μ with respect to the scale parameter d_u for the energy scale $\Lambda_u = 10 TeV$, the coupling $\lambda_{\mu\mu} = 0.1$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for total (diagonal, off-diagonal) part. The EDM d_μ lies in the broad range

$10^{-26} e - cm \leq d_\mu \leq 10^{-19} e - cm$ for the interval of the scale parameter $1.0 < d_u < 2.0$ and the experimental current limit is reached for the small values of the scale d_u , $d_u \leq 1.1$.

Fig. 6 represents the EDM d_μ with respect to the energy scale Λ_u , for the coupling $\lambda_{\mu\mu} = 0.1$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.1$, $d_u = 1.4$, $d_u = 1.6$, including diagonal and off diagonal terms. The increasing value of the energy scale Λ_u causes that the EDM decreases with the suppressed sensitivity, similar to the electron EDM case.

In Fig. 7 we show the EDM d_μ with respect to the coupling $\lambda = \lambda_{\mu\mu}$ for the energy scale $\Lambda_u = 10 TeV$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.1$, $d_u = 1.4$, $d_u = 1.6$, including diagonal and off diagonal terms. It is observed that the experimental current limit can be reached if the Yukawa coupling λ is of the order of 0.1 (1.0), for the scale dimension $d_u < 1.1$ ($d_u \sim 1.4$).

Figs. 8 is devoted to the contribution of the scalar unparticle to the EDM d_τ with respect to the scale parameter d_u for the energy scale $\Lambda_u = 10 TeV$, the coupling $\lambda_{\tau\tau} = 1.0$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed) line represents the EDM for total (off-diagonal) contribution. For the interval of the scale parameter $1.0 < d_u < 2.0$ the EDM d_τ takes the values in the range $10^{-23} e - cm \leq d_\mu \leq 10^{-17} e - cm$. The off diagonal contribution is almost five order smaller than the diagonal contribution which coincides with the total one. Here, the experimental current value can be reached for the small values of the scale d_u , $d_u < 1.1$.

Fig. 9; 10 represents the EDM d_τ with respect to the energy scale Λ_u ; the coupling $\lambda = \lambda_{\tau\tau}$, for the coupling $\lambda_{\tau\tau} = 1.0$; for the energy scale $\Lambda_u = 10 TeV$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.1$, $d_u = 1.4$, $d_u = 1.6$, including diagonal and off diagonal terms. Here it is observed that the experimental value $\sim 10^{-16} GeV$ is reached for the energy scale Λ_u , $\Lambda_u < 10 TeV$, $d_u < 1.1$ and for the coupling λ at least of the order of 1.0. The contribution of the scalar unparticle to the EDM decreases with the increasing values of the energy scale Λ_u .

Finally, in Figs. 11; 12; 13 we present the EDMs d_e ; d_μ ; d_τ with respect to the CP parameter $\sin\theta$, for the energy scale $\Lambda_u = 10 TeV$ and the couplings $\lambda_{ee} = 0.01$; $\lambda_{\mu\mu} = 0.1$; $\lambda_{\tau\tau} = 1.0$, respectively. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.4$, $d_u = 1.6$, $d_u = 1.8$; $d_u = 1.1$, $d_u = 1.4$, $d_u = 1.6$; $d_u = 1.1$, $d_u = 1.4$, $d_u = 1.6$, including diagonal and off diagonal terms. These figures show that the EDM changes almost one order in the range

$0.1 < \sin\theta < 0.8$ and, even for the tiny CP parameter, for the appropriate values of the scaling dimension, it is possible to reach the current experimental limits.

Now we would like to summarize our results:

- The charged lepton EDMs are strongly sensitive to the scaling dimension d_u . The experimental current limit of d_e can be reached in the range $1.6 \leq d_u \leq 1.8$. For d_μ and d_τ the current limits are reached for the small values of the scale d_u , $d_u \leq 1.1$. The more reliable range for the the scaling dimension can be estimated with more accurate measurements of the charged lepton dipole moments.
- The contribution of the scalar unparticle to the EDM decreases with the increasing values of the energy scale Λ_u as expected and the e sensitivity is large for the numerical values $\Lambda_u < 10 TeV$.
- The EDMs are strongly sensitive to the coupling λ and with the choice of the appropriate numerical value of the scaling dimension, the experimental current limits of charged lepton EDMs can be reached, even with tiny couplings.
- The CP parameter θ is the source of the charged lepton EDM and it is another free parameter that should be restricted. The EDMs change almost one order in the range $0.1 < \sin\theta < 0.8$.

The forthcoming most accurate experimental measurement of charged lepton EDMs would be illuminating to search the effects of unparticles and to predict their scaling dimensions.

References

- [1] H. Georgi, *Phys. Rev. Lett.* **98**, 221601 (2007).
- [2] H. Georgi, *Phys. Lett.* **B650**, 275 (2007).
- [3] K. Cheung, W. Y. Keung and T. C. Yuan, *Phys. Rev. Lett.* **99**, 051803 (2007).
- [4] M. X. Luo and G. H. Zhu, hep-ph/0704.3532.
- [5] M. A. Stephanov, *Phys. Rev. D* **76**, 035008 (2007).
- [6] R. Zwicky, hep-ph/0707.0677.
- [7] K. Cheung, W. Y. Keung and T. C. Yuan, hep-ph/0706.3155.

- [8] M. X. Luo, W. Wu and G. H. Zhu, hep-ph/0708.0671.
- [9] C. H. Chen and C. Q. Geng, hep-ph/0705.0689.
- [10] C. H. Chen and C. Q. Geng, *Phys. Rev.* **D76**, 036007 (2007).
- [11] C. H. Chen and C. Q. Geng, hep-ph/0709.0235.
- [12] S. L. Chen and X. G. He, hep-ph/0705.3946.
- [13] T. M. Aliev, A. S. Cornell and N. Gaur, hep-ph/0705.1326.
- [14] T. M. Aliev, A. S. Cornell and N. Gaur, *JHEP* **07**, 072 (2007).
- [15] T.M. Aliev, M. Savci hep-ph/0710.1505
- [16] G. J. Ding and M. L. Yan, hep-ph/0705.0794.
- [17] G. J. Ding and M. L. Yan, hep-ph/0706.0325.
- [18] X. Q. Li and Z. T. Wei, *Phys. Lett.* **B651**, 380 (2007).
- [19] X. Q. Li and Z. T. Wei, hep-ph/0707.2285.
- [20] Y. Liao, hep-ph/0705.0837.
- [21] Y. Liao, hep-ph/0708.3327.
- [22] Y. Liao and J. Y. Liu, hep-ph/0706.1284.
- [23] P. J. Fox, A. Rajaraman and Y. Shirman, hep-ph/0705.3092.
- [24] S. Catterall and F. Sannino, *Phys. Rev.* **D76**, 034504 (2007).
- [25] C. D. Lu, W. Wang and Y. M. Wang, hep-ph/0705.2909.
- [26] N. Greiner, hep-ph/0705.3518.
- [27] D. Choudhury, D. K. Ghosh and Mamta, hep-ph/0705.3637
- [28] H. Davoudiasl, hep-ph/0705.3636.
- [29] S. L. Chen, X. G. He and H. C. Tsai, hep-ph/0707.0187.
- [30] P. Mathews and V. Ravindran, hep-ph/0705.4599.

- [31] S. Zhou, hep-ph/0706.0302.
- [32] R. Foadi, M. T. Frandsen, T. A. Ryttov and F. Sannino, hep-ph/0706.1696.
- [33] M. Bander, J. L. Feng, A. Rajaraman and Y. Shirman, hep-ph/0706.2677.
- [34] T. G. Rizzo, hep-ph/0706.3025.
- [35] H. Goldberg and P. Nath, hep-ph/0706.3898.
- [36] T. Kikuchi and N. Okada, hep-ph/0707.0893.
- [37] R. Mohanta and A. K. Giri, hep-ph/0707.1234.
- [38] R. Mohanta and A. K. Giri, hep-ph/0707.3308.
- [39] C. S. Huang and X. H. Wu, hep-ph/0707.1268.
- [40] N. V. Krasnikov, hep-ph/0707.1419.
- [41] A. Lenz, hep-ph/0707.1535.
- [42] D. Choudhury and D. K. Ghosh, hep-ph/0707.2074.
- [43] H. Zhang, C. S. Li and Z. Li, hep-ph/0707.2132.
- [44] Y. Nakayama, hep-ph/0707.2451.
- [45] N. G. Deshpande, X. G. He and J. Jiang, hep-ph/0707.2959.
- [46] N. G. Deshpande, S. D. H. Hsu and J. Jiang, hep-ph/0708.2735.
- [47] A. Delgado, J. R. Espinosa and M. Quiros, hep-ph/0707.4309.
- [48] M. Neubert, hep-ph/0708.0036.
- [49] S. Hannestad, G. Raffelt and Y. Y. Y. Wong, hep-ph/0708.1404.
- [50] P. K. Das, hep-ph/0708.2812.
- [51] S. Das, S. Mohanty and K. Rao, hep-ph/0709.2583.
- [52] G. Bhattacharyya, D. Choudhury and D. K. Ghosh, hep-ph/0708.2835.
- [53] D. Majumdar, hep-ph/0708.3485.

- [54] A. T. Alan and N. K. Pak, hep-ph/0708.3802.
- [55] A. Freitas and D. Wyler, hep-ph/0708.4339.
- [56] I. Gogoladze, N. Okada and Q. Shafi, hep-ph/0708.4405.
- [57] T. i. Hur, P. Ko and X. H. Wu, hep-ph/0709.0629.
- [58] L. Anchordoqui and H. Goldberg, hep-ph/0709.0678.
- [59] S. Majhi, hep-ph/0709.1960.
- [60] J. McDonald, hep-ph/0709.2350.
- [61] M. C. Kumar, P. Mathews, V. Ravindran and A. Tripathi, hep-ph/0709.2478.
- [62] K. M. Cheung, W. Y. Keung, T. C. Yuan, hep-ph/0710.2230
- [63] A. Kobakhidze, hep-ph/0709.3782
- [64] G. J. Ding and M. L. Yan, hep-ph/0709.3435.
- [65] A.B. Balantekin , K.O. Ozansoy, hep-ph/0710.0028.
- [66] E. D. Commins et.al. *Phys. Rev. A* **50**, 2960 (1994).
- [67] J. Bailey et al, *Journ. Phys.* **G4**, 345 (1978);
- [68] Particle Data Group, D. E. Groom et.al. *European Phys. Journ.* **C15**, 1 (2000).
- [69] I. B. Khriplovich, *Yad. Fiz.* **44**, 1019 (1986) ; (*Sov. J. Nucl. Phys.* **44**, 659 (1986)).
- [70] I. B. Khriplovich, *Phys. Lett.* **B173**, 193 (1986).
- [71] A. Czarnecki and B. Krause, *Acta Phys. Polon.* **B28**, 829 (1997).
- [72] A. Czarnecki and B. Krause, *Phys. Rev. Lett* **78**, 4339 (1997).
- [73] B. Dutta, R. N. Mohapatra, *Phys. Rev.* **D68**, 113008 (2003).
- [74] E. Iltan, *Phys. Rev.* **D64**, 013013 (2001).
- [75] E. Iltan, *JHEP* **065**, 0305 (2003).
- [76] E. Iltan, *JHEP* **0404**, 018 (2004).

[77] E. Iltan, *Eur. Phys. J.* **C44**, 411 (2005).

[78] E. Iltan, *Eur. Phys. J.* **C51**, 689 (2007).

[79] E. Iltan, hep-ph/0708.3765.




Figure 1: One loop diagram contributes to the EDM of charged leptons due to scalar unparticle. Wavy (solid) line represents the electromagnetic field (lepton field) and double dashed line the scalar unparticle field.




Figure 2: d_e with respect to the scale parameter d_u for the energy scale $\Lambda_u = 10 TeV$, the coupling $\lambda_{ee} = 0.01$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for total (diagonal, off-diagonal) part.




Figure 3: d_e with respect to the energy scale Λ_u , for the coupling $\lambda_{ee} = 0.01$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.4$, $d_u = 1.6$, $d_u = 1.8$.




Figure 4: d_e with respect to the coupling λ for the energy scale $\Lambda_u = 10 TeV$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.4$, $d_u = 1.6$, $d_u = 1.8$.




Figure 5: The same as Fig. 2 but for d_{μ} and the coupling $\lambda_{\mu\mu} = 0.1$




Figure 6: d_{μ} with respect to the energy scale Λ_u , for the coupling $\lambda_{\mu\mu} = 0.1$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.1$, $d_u = 1.4$, $d_u = 1.6$.




Figure 7: d_μ with respect to the coupling λ for the energy scale $\Lambda_u = 10 TeV$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.1$, $d_u = 1.4$, $d_u = 1.6$.




Figure 8: The same as Fig. 2 but for d_τ and the coupling $\lambda_{\tau\tau} = 1.0$.




Figure 9: d_τ with respect to the energy scale Λ_u for the coupling $\lambda_{\tau\tau} = 1.0$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.1$, $d_u = 1.4$, $d_u = 1.6$.




Figure 10: d_τ with respect to the coupling λ for the energy scale $\Lambda_u = 10 \text{ TeV}$ and the intermediate value of the CP parameter $\sin\theta = 0.5$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.1$, $d_u = 1.4$, $d_u = 1.6$.




Figure 11: d_e with respect to the CP parameter $\sin\theta$, for the energy scale $\Lambda_u = 10 \text{ TeV}$ and the coupling $\lambda = 0.01$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.4$, $d_u = 1.6$, $d_u = 1.8$.




Figure 12: d_μ with respect to the CP parameter $\sin\theta$, for the energy scale $\Lambda_u = 10 \text{ TeV}$ and the coupling $\lambda = 0.1$. Here the solid (dashed, small dashed) line represents the EDM for $d_u = 1.1$, $d_u = 1.4$, $d_u = 1.6$.




Figure 13: The same as the Fig.12 but for d_τ and the coupling $\lambda = 1$.