

General analysis of the rare $B_c \rightarrow D_s^* \ell^+ \ell^-$ decay beyond the standard model

U. O. Yilmaz ¹ and G. Turan ^{2 *}

¹ Physics Department, Mersin University 33343 Ciftlikkoy Mersin, Turkey

² Physics Department, Middle East Technical University 06531 Ankara, Turkey

Abstract

The general analysis of the rare $B_c \rightarrow D_s^* \ell^+ \ell^-$ decay is presented by using the most general, model independent effective Hamiltonian. The dependencies of the branching ratios, longitudinal, normal and transversal polarization asymmetries for ℓ^- and the combined asymmetries for ℓ^- and ℓ^+ on the new Wilson coefficients are investigated. Our analysis shows that the lepton polarization asymmetries are very sensitive to the scalar and tensor type interactions, which will be very useful in looking for new physics beyond the standard model.

*e-mail: gsevgur@metu.edu.tr

1 Introduction

Rare B meson decays, induced by flavor-changing neutral current (FCNC) $b \rightarrow s, d$ transitions, are excellent places to search for new physics because they appear at the same order as the standard model (SM). In rare B meson decays, effects of the new physics may appear in two different manners, either through the new contributions to the Wilson coefficients existing in the SM or through the new structures in the effective Hamiltonian which are absent in the SM.

There have been many investigations of the new physics through the study of rare radiative, leptonic and semileptonic decays of $B_{u,d,s}$ mesons induced by FCNC transitions of $b \rightarrow s, d$ [1] since the CLEO observation of $b \rightarrow s \gamma$ [2]. The studies will be even more complete if similar decays for B_c are also included.

The study of the B_c meson is by itself quite interesting too, due to its outstanding features [3]-[5]. It is the lowest bound state of two heavy quarks (b and c) with explicit flavor that can be compared with the charmonium ($c\bar{c}$ - bound state) and bottomium ($b\bar{b}$ - bound state) which have implicit flavor. The implicit-flavor states decay strongly and electromagnetically whereas the B_c meson decays weakly. The major difference between the weak decay properties of B_c and $B_{u,d,s}$ is that those of the latter ones are described very well in the framework of the heavy quark limit, which gives some relations between the form factors of the physical process. In case of B_c meson, the heavy flavor and spin symmetries must be reconsidered because both b and c are heavy.

From the experimental side, the running B factories in KEK and SLAC continue to collect data samples and encourage the study of rare B meson decays. It is believed that in future experiments at hadronic colliders, such as the BTeV and LHC-B most of rare B_c decays should be accessible.

One of the efficient ways in establishing new physics beyond the SM is the measurement of the lepton polarization [7]-[16]. In this work we present a study of the branching ratio and lepton polarizations in the exclusive $B_c \rightarrow D_s^* \ell^+ \ell^-$ ($\ell = \mu, \tau$) decay for a general form of the effective Hamiltonian including all possible form of interactions in a model independent way without forcing concrete values for the Wilson coefficients corresponding to any specific model.

It is well known that the theoretical study of the inclusive decays is rather easy but their experimental investigation is difficult. However for the exclusive decays the situation is contrary to the inclusive case, i.e., their experimental detection is very easy but theoretical investigation has its own drawbacks. This is due to the fact that for description of the exclusive decay form factors, i.e., the matrix elements of the effective Hamiltonian between initial and final meson states, are needed. This problem is related to the nonperturbative sector of the QCD and it can only be solved in framework of the nonperturbative approaches.

These matrix elements have been studied in framework of different approaches, such as light front, constituent quark models [5], and a relativistic quark model proposed in ref. [6]. In this work we will use the weak decay form factors calculated in ref. [6].

The paper is organized as follows. In section 2, we first give the effective Hamiltonian for the quark level process $b \rightarrow s \ell^+ \ell^-$ and the definitions of the form factors, and then introduce the corresponding matrix element. In section 3, we present the model independent

expressions for the longitudinal, transversal and normal polarizations of leptons. We also give the lepton-antilepton combined asymmetries. Section 4 is devoted to the numerical analysis and discussion of our results.

2 Effective Hamiltonian

In the standard effective Hamiltonian approach, the $B_c \rightarrow D_s^* \ell^+ \ell^-$ decay is described at the quark level by the $b \rightarrow s \ell^+ \ell^-$ process, which can be written in terms of twelve model independent four-Fermi interactions as follows [11],

$$\begin{aligned} \mathcal{H}_{eff} = & \frac{G\alpha}{\sqrt{2}\pi} V_{ts} V_{tb}^* \left\{ C_{SL} \bar{s} i \sigma_{\mu\nu} \frac{q^\nu}{q^2} L b \bar{\ell} \gamma^\mu \ell + C_{BR} \bar{s} i \sigma_{\mu\nu} \frac{q^\nu}{q^2} R b \bar{\ell} \gamma^\mu \ell \right. \\ & + C_{LL}^{tot} \bar{s}_L \gamma_\mu b_L \bar{\ell}_L \gamma^\mu \ell_L + C_{LR}^{tot} \bar{s}_L \gamma_\mu b_L \bar{\ell}_R \gamma^\mu \ell_R + C_{RL} \bar{s}_R \gamma_\mu b_R \bar{\ell}_L \gamma^\mu \ell_L \\ & + C_{RR} \bar{s}_R \gamma_\mu b_R \bar{\ell}_R \gamma^\mu \ell_R + C_{LRLR} \bar{s}_L b_R \bar{\ell}_L \ell_R + C_{RLLR} \bar{s}_R b_L \bar{\ell}_L \ell_R \\ & + C_{LRRL} \bar{s}_L b_R \bar{\ell}_R \ell_L + C_{RLRL} \bar{s}_R b_L \bar{\ell}_R \ell_L + C_T \bar{s} \sigma_{\mu\nu} b \bar{\ell} \sigma^{\mu\nu} \ell \\ & \left. + i C_{TE} \epsilon^{\mu\nu\alpha\beta} \bar{s} \sigma_{\mu\nu} b \bar{\ell} \sigma_{\alpha\beta} \ell \right\}, \end{aligned} \quad (1)$$

where the chiral projection operators L and R in (1) are defined as

$$L = \frac{1 - \gamma_5}{2}, \quad R = \frac{1 + \gamma_5}{2},$$

and C_X are the coefficients of the four-Fermi interactions. The coefficients, C_{SL} and C_{BR} , are the nonlocal Fermi interactions which correspond to $-2m_s C_7^{eff}$ and $-2m_b C_7^{eff}$ in the SM, respectively. The following four terms in Eq. (1) are the vector type interactions with coefficients C_{LL} , C_{LR} , C_{RL} and C_{RR} . Two of these vector interactions containing C_{LL}^{tot} and C_{LR}^{tot} already exist in the SM in combinations of the form $(C_9^{eff} - C_{10})$ and $(C_9^{eff} + C_{10})$. Therefore we write

$$\begin{aligned} C_{LL}^{tot} &= C_9^{eff} - C_{10} + C_{LL}, \\ C_{LR}^{tot} &= C_9^{eff} + C_{10} + C_{LR}, \end{aligned}$$

so that C_{LL}^{tot} and C_{LR}^{tot} describe the sum of the contributions from SM and the new physics. The terms with coefficients C_{LRLR} , C_{RLLR} , C_{LRRL} and C_{RLRL} describe the scalar type interactions. The remaining two terms with the coefficients C_T and C_{TE} describe the tensor type interactions.

After giving the general form of four-Fermi interaction for the $b \rightarrow s \ell^+ \ell^-$ transition, we now need to estimate the matrix element for the $B_c \rightarrow D_s^* \ell^+ \ell^-$ decay. These can be expressed in term of invariant form factors as follows:

$$\begin{aligned} \langle D_s^*(p_{D^*}, \varepsilon) | \bar{s} \gamma_\mu (1 \pm \gamma_5) b | B_c(p_{B_c}) \rangle = \\ -\epsilon_{\mu\nu\lambda\sigma} \varepsilon^{*\nu} p_{D^*}^\lambda q^\sigma \frac{2V(q^2)}{m_{B_c} + m_{D^*}} \pm i \varepsilon_\mu^* (m_{B_c} - m_{D^*}) A_0(q^2) \\ \mp i (p_{B_c} + p_{D^*})_\mu (\varepsilon^* q) \frac{A_+(q^2)}{m_{B_c} + m_{D^*}} \mp i q_\mu (\varepsilon^* q) \frac{A_-(q^2)}{m_{B_c} + m_{D^*}}, \end{aligned} \quad (2)$$

$$\begin{aligned}
\langle D_s^*(p_{D^*}, \varepsilon) | \bar{s} i \sigma_{\mu\nu} q^\nu (1 \pm \gamma_5) b | B(p_{B_c}) \rangle = \\
2\epsilon_{\mu\nu\lambda\sigma} \varepsilon^{*\nu} p_{D^*}^\lambda q^\sigma g(q^2) \pm i \left[\varepsilon_\mu^* (m_{B_c}^2 - m_{D^*}^2) - (p_{B_c} + p_{D^*})_\mu (\varepsilon^* q) \right] a_0(q^2) \\
\pm i (\varepsilon^* q) \left[q_\mu - (p_{B_c} + p_{D^*})_\mu \frac{q^2}{m_{B_c}^2 - m_{D^*}^2} \right] \frac{(m_{B_c}^2 - m_{D^*}^2)}{q^2} (a_+(q^2) - a_0(q^2)) ,
\end{aligned} \tag{3}$$

$$\begin{aligned}
\langle D_s^*(p_{D^*}, \varepsilon) | \bar{s} \sigma_{\mu\nu} b | B(p_{B_c}) \rangle = \\
i\epsilon_{\mu\nu\lambda\sigma} \left[-g(q^2) \varepsilon^{*\lambda} (p_{B_c} + p_{D^*})^\sigma + \frac{1}{q^2} (m_{B_c}^2 - m_{D^*}^2) \varepsilon^{*\lambda} q^\sigma (g(q^2) - a_0(q^2)) \right. \\
\left. - \frac{2}{q^2} \left(g(q^2) - a_+(q^2) \right) (\varepsilon^* q) p_{D^*}^\lambda q^\sigma \right] .
\end{aligned} \tag{4}$$

$$\begin{aligned}
\langle D_s^*(p_{D^*}, \varepsilon) | \bar{s} (1 \pm \gamma_5) b | B(p_{B_c}) \rangle = \frac{1}{m_b} \Big[\mp (\varepsilon^* q) (m_{B_c} - m_{D_s^*}) \\
+ A_0(q^2) - A_+(q^2) - \frac{q^2}{m_{B_c}^2 - m_{D^*}^2} A_-(q^2) \Big] .
\end{aligned} \tag{5}$$

where $q = p_{B_c} - p_{D_s^*}$ is the momentum transfer and ε is the polarization vector of D_s^* meson. The matrix element $\langle D_s^* | \bar{s} (1 \pm \gamma_5) b | B \rangle$ is calculated by contracting both sides of Eq. (2) with q^μ and using equation of motion.

By using Eqs. (1)–(5), we can now write the matrix element of the $B_c \rightarrow D_s^* \ell^+ \ell^-$ decay as

$$\begin{aligned}
\mathcal{M}(B_c \rightarrow D_s^* \ell^+ \ell^-) = \frac{G\alpha}{4\sqrt{2}\pi} V_{tb} V_{ts}^* \\
\times \left\{ \bar{\ell} \gamma^\mu (1 - \gamma_5) \ell \left[-2A_1 \epsilon_{\mu\nu\lambda\sigma} \varepsilon^{*\nu} p_{D_s^*}^\lambda q^\sigma - iB_1 \varepsilon_\mu^* + iB_2 (\varepsilon^* q) (p_{B_c} + p_{D_s^*})_\mu + iB_3 (\varepsilon^* q) q_\mu \right] \right. \\
+ \bar{\ell} \gamma^\mu (1 + \gamma_5) \ell \left[-2C_1 \epsilon_{\mu\nu\lambda\sigma} \varepsilon^{*\nu} p_{D_s^*}^\lambda q^\sigma - iD_1 \varepsilon_\mu^* + iD_2 (\varepsilon^* q) (p_{B_c} + p_{D_s^*})_\mu + iD_3 (\varepsilon^* q) q_\mu \right] \\
+ \bar{\ell} (1 - \gamma_5) \ell \left[iB_4 (\varepsilon^* q) \right] + \bar{\ell} (1 + \gamma_5) \ell \left[iB_5 (\varepsilon^* q) \right] \\
+ 4\bar{\ell} \sigma^{\mu\nu} \ell \left(iC_T \epsilon_{\mu\nu\lambda\sigma} \right) \left[-2g \varepsilon^{*\lambda} (p_{B_c} + p_{D_s^*})^\sigma + B_6 \varepsilon^{*\lambda} q^\sigma - B_7 (\varepsilon^* q) p_{D_s^*}^\lambda q^\sigma \right] \\
\left. + 16C_{TE} \bar{\ell} \sigma_{\mu\nu} \ell \left[-2g \varepsilon^{*\mu} (p_{B_c} + p_{D_s^*})^\nu + B_6 \varepsilon^{*\mu} q^\nu - B_7 (\varepsilon^* q) p_{D_s^*}^\mu q^\nu \right] \right\} ,
\end{aligned} \tag{6}$$

where

$$\begin{aligned}
A_1 &= (C_{LL}^{tot} + C_{RL}) \frac{V}{m_{B_c} + m_{D_s^*}} - (C_{BR} + C_{SL}) \frac{g}{q^2} , \\
B_1 &= (C_{LL}^{tot} - C_{RL}) (m_{B_c} - m_{D_s^*}) A_0 - (C_{BR} - C_{SL}) (m_{B_c}^2 - m_{D_s^*}^2) \frac{a_0}{q^2} , \\
B_2 &= \frac{C_{LL}^{tot} - C_{RL}}{m_{B_c} + m_{D_s^*}} A_+ - (C_{BR} - C_{SL}) \frac{a_+}{q^2} , \\
B_3 &= (C_{LL}^{tot} - C_{RL}) \frac{A_-}{m_{B_c} + m_{D_s^*}} + 2(C_{BR} - C_{SL}) (a_+ - a_0) \frac{m_{B_c}^2 - m_{D_s^*}^2}{q^4} ,
\end{aligned}$$

$$\begin{aligned}
B_4 &= -(C_{LRRL} - C_{RLRL}) \frac{m_{B_c} - m_{D_s^*}}{m_b} \left(A_0 - A_- - \frac{q^2}{m_{B_c}^2 - m_{D_s^*}^2} \right), \\
B_5 &= -(C_{LRLR} - C_{RLLR}) \frac{m_{B_c} - m_{D_s^*}}{m_b} \left(A_0 - A_- - \frac{q^2}{m_{B_c}^2 - m_{D_s^*}^2} \right), \\
B_6 &= (m_{B_c}^2 - m_{D_s^*}^2) \frac{g - a_0}{q^2}, \\
B_7 &= \frac{2}{q^2} (g - a_0 - a_+) , \\
C_1 &= A_1 (C_{LL}^{tot} \rightarrow C_{LR}^{tot}, C_{RL} \rightarrow C_{RR}), \\
D_1 &= B_1 (C_{LL}^{tot} \rightarrow C_{LR}^{tot}, C_{RL} \rightarrow C_{RR}), \\
D_2 &= B_2 (C_{LL}^{tot} \rightarrow C_{LR}^{tot}, C_{RL} \rightarrow C_{RR}), \\
D_3 &= B_3 (C_{LL}^{tot} \rightarrow C_{LR}^{tot}, C_{RL} \rightarrow C_{RR}). \tag{7}
\end{aligned}$$

3 Lepton polarizations

We now like to calculate the final lepton polarizations for the $B_c \rightarrow D_s^* \ell^+ \ell^-$ decay. For this we will use the convention followed by the earlier works, such as [10],[11] and define the following orthogonal unit vectors, $S_i^{-\mu}$ in the rest frame of ℓ^- and $S_i^{+\mu}$ in the rest frame of ℓ^+ , for the polarization of the leptons along the longitudinal ($i = L$), transverse ($i = T$) and normal ($i = N$) directions:

$$\begin{aligned}
S_L^{-\mu} &\equiv (0, \vec{e}_L^-) = \left(0, \frac{\vec{p}_-}{|\vec{p}_-|} \right), \\
S_N^{-\mu} &\equiv (0, \vec{e}_N^-) = \left(0, \frac{\vec{p} \times \vec{p}_-}{|\vec{p} \times \vec{p}_-|} \right), \\
S_T^{-\mu} &\equiv (0, \vec{e}_T^-) = \left(0, \vec{e}_N^- \times \vec{e}_L^- \right), \\
S_L^{+\mu} &\equiv (0, \vec{e}_L^+) = \left(0, \frac{\vec{p}_+}{|\vec{p}_+|} \right), \\
S_N^{+\mu} &\equiv (0, \vec{e}_N^+) = \left(0, \frac{\vec{p} \times \vec{p}_+}{|\vec{p} \times \vec{p}_+|} \right), \\
S_T^{+\mu} &\equiv (0, \vec{e}_T^+) = \left(0, \vec{e}_N^+ \times \vec{e}_L^+ \right), \tag{8}
\end{aligned}$$

where \vec{p}_\pm and \vec{p} are the three momenta of ℓ^\pm and D_s^* meson in the center of mass (CM) frame of the $\ell^+ \ell^-$ system, respectively. The longitudinal unit vectors S_L^- and S_L^+ are boosted to CM frame of $\ell^+ \ell^-$ by Lorentz transformation,

$$\begin{aligned}
S_{L,CM}^{-\mu} &= \left(\frac{|\vec{p}_-|}{m_\ell}, \frac{E_\ell \vec{p}_-}{m_\ell |\vec{p}_-|} \right), \\
S_{L,CM}^{+\mu} &= \left(\frac{|\vec{p}_-|}{m_\ell}, -\frac{E_\ell \vec{p}_-}{m_\ell |\vec{p}_-|} \right), \tag{9}
\end{aligned}$$

while vectors of perpendicular directions are not changed by boost.

The differential decay rate of the $B_c \rightarrow D_s^* \ell^+ \ell^-$ decay for any spin direction \vec{n}^\pm of the ℓ^\pm can be written in the following form

$$\frac{d\Gamma(\vec{n}^\pm)}{ds} = \frac{1}{2} \left(\frac{d\Gamma}{ds} \right)_0 \left[1 + \left(P_L^\pm \vec{e}_L^\pm + P_N^\pm \vec{e}_N^\pm + P_T^\pm \vec{e}_T^\pm \right) \cdot \vec{n}^\pm \right], \quad (10)$$

where \vec{n}^\pm is the unit vector in the ℓ^\pm rest frame and $s = q^2/m_{B_c}^2$. Here, the superscripts $+$ and $-$ correspond to ℓ^+ and ℓ^- cases, the subscript 0 corresponds to the unpolarized decay rate, whose explicit form is given by

$$\begin{aligned} \left(\frac{d\Gamma}{ds} \right)_0 &= \frac{G^2 \alpha^2 m_{B_c}}{2^{14} \pi^5} |V_{tb} V_{ts}^*|^2 \sqrt{\lambda} v \\ &\times \left\{ \frac{32}{3} m_{B_c}^4 \lambda \left[(m_{B_c}^2 s - m_\ell^2) (|A_1|^2 + |C_1|^2) + 6 m_\ell^2 \text{Re}(A_1 C_1^*) \right] \right. \\ &+ 96 m_\ell^2 \text{Re}(B_1 D_1^*) - \frac{4}{r} m_{B_c}^2 m_\ell \lambda \text{Re}[(B_1 - D_1)(B_4^* - B_5^*)] \\ &+ \frac{8}{r} m_{B_c}^2 m_\ell^2 \lambda \left(\text{Re}[B_1(-B_3^* + D_2^* + D_3^*)] + \text{Re}[D_1(B_2^* + B_3^* - D_3^*)] - \text{Re}(B_4 B_5^*) \right) \\ &+ \frac{4}{r} m_{B_c}^4 m_\ell (1-r) \lambda \left(\text{Re}[(B_2 - D_2)(B_4^* - B_5^*)] + 2 m_\ell \text{Re}[(B_2 - D_2)(B_3^* - D_3^*)] \right) \\ &- \frac{8}{r} m_{B_c}^4 m_\ell^2 \lambda (2 + 2r - s) \text{Re}(B_2 D_2^*) + \frac{4}{r} m_{B_c}^4 m_\ell s \lambda \text{Re}[(B_3 - D_3)(B_4^* - B_5^*)] \\ &+ \frac{4}{r} m_{B_c}^4 m_\ell^2 s \lambda |B_3 - D_3|^2 + \frac{2}{r} m_{B_c}^2 (m_{B_c}^2 s - 2 m_\ell^2) \lambda (|B_4|^2 + |B_5|^2) \\ &- \frac{8}{3rs} m_{B_c}^2 \lambda \left[m_\ell^2 (2 - 2r + s) + m_{B_c}^2 s (1 - r - s) \right] \left[\text{Re}(B_1 B_2^*) + \text{Re}(D_1 D_2^*) \right] \\ &+ \frac{4}{3rs} \left[2 m_\ell^2 (\lambda - 6rs) + m_{B_c}^2 s (\lambda + 12rs) \right] (|B_1|^2 + |D_1|^2) \\ &+ \frac{4}{3rs} m_{B_c}^4 \lambda \left(m_{B_c}^2 s \lambda + m_\ell^2 [2\lambda + 3s(2 + 2r - s)] \right) (|B_2|^2 + |D_2|^2) \\ &+ \frac{32}{r} m_{B_c}^6 m_\ell \lambda^2 \text{Re}[(B_2 + D_2)(B_7 C_{TE})^*] \\ &- \frac{32}{r} m_{B_c}^4 m_\ell \lambda (1 - r - s) \left(\text{Re}[(B_1 + D_1)(B_7 C_{TE})^*] + 2 \text{Re}[(B_2 + D_2)(B_6 C_{TE})^*] \right) \\ &+ \frac{64}{r} (\lambda + 12rs) m_{B_c}^2 m_\ell \text{Re}[(B_1 + D_1)(B_6 C_{TE})^*] \\ &+ \frac{256}{3rs} |g|^2 |C_T|^2 m_{B_c}^2 \left(4 m_\ell^2 [\lambda(8r - s) - 12rs(2 + 2r - s)] \right. \\ &+ m_{B_c}^2 s [\lambda(16r + s) + 12rs(2 + 2r - s)] \Big) \\ &+ \frac{1024}{3rs} |g|^2 |C_{TE}|^2 m_{B_c}^2 \left(8 m_\ell^2 [\lambda(4r + s) + 12rs(2 + 2r - s)] \right. \\ &+ m_{B_c}^2 s [\lambda(16r + s) + 12rs(2 + 2r - s)] \Big) \\ &- \frac{128}{r} m_{B_c}^2 m_\ell [\lambda + 12r(1 - r)] \text{Re}[(B_1 + D_1)(g C_{TE})^*] \\ &+ \frac{128}{r} m_{B_c}^4 m_\ell \lambda (1 + 3r - s) \text{Re}[(B_2 + D_2)(g C_{TE})^*] + 512 m_{B_c}^4 m_\ell \lambda \text{Re}[(A_1 + C_1)(g C_T)^*] \end{aligned} \quad (11)$$

$$\begin{aligned}
& + \frac{16}{3r} m_{B_c}^2 \left(4(m_{B_c}^2 s + 8m_\ell^2) |C_{TE}|^2 + m_{B_c}^2 s v^2 |C_T|^2 \right) \times \left(4(\lambda + 12rs) |B_6|^2 \right. \\
& + m_{B_c}^4 \lambda^2 |B_7|^2 - 4m_{B_c}^2 (1 - r - s) \lambda \operatorname{Re}(B_6 B_7^*) - 16 [\lambda + 12r(1 - r)] \operatorname{Re}(g B_6^*) \\
& \left. + 8m_{B_c}^2 (1 + 3r - s) \lambda \operatorname{Re}(g B_7^*) \right) \Big\} ,
\end{aligned}$$

where $\lambda = 1 + r^2 + s^2 - 2r - 2s - 2rs$, $r = m_{D_s^*}^2 / m_{B_c}^2$ and $v = \sqrt{1 - \frac{4m_\ell^2}{sm_{B_c}^2}}$ is the lepton velocity.

The polarizations P_L^\pm , P_T^\pm and P_N^\pm in Eq. (10) are defined by the equation

$$P_i^\pm(q^2) = \frac{\frac{d\Gamma}{dq^2}(\vec{n}^\pm = \vec{e}_i^\pm) - \frac{d\Gamma}{dq^2}(\vec{n}^\pm = -\vec{e}_i^\pm)}{\frac{d\Gamma}{dq^2}(\vec{n}^\pm = \vec{e}_i^\pm) + \frac{d\Gamma}{dq^2}(\vec{n}^\pm = -\vec{e}_i^\pm)} ,$$

for $i = L, N, T$, i.e., P_L^\pm and P_T^\pm represents the charged lepton ℓ^\pm longitudinal and transversal asymmetries in the decay plane, respectively, and P_N^\pm is the normal component to both of them. After some lengthy algebra, we get for the longitudinal polarization of the ℓ^\pm

$$\begin{aligned}
P_L^\pm &= \frac{4}{\Delta} m_{B_c}^2 v \left\{ \mp \frac{1}{3r} \lambda^2 m_{B_c}^4 [|B_2|^2 - |D_2|^2] + \frac{1}{r} \lambda m_\ell \operatorname{Re}[(B_1 - D_1)(B_4^* + B_5^*)] \right. \\
&- \frac{1}{r} \lambda m_{B_c}^2 m_\ell (1 - r) \operatorname{Re}[(B_2 - D_2)(B_4^* + B_5^*)] \mp \frac{8}{3} \lambda m_{B_c}^4 s [|A_1|^2 - |C_1|^2] \\
&- \frac{1}{2r} \lambda m_{B_c}^2 s [|B_4|^2 - |B_5|^2] - \frac{1}{r} \lambda m_{B_c}^2 m_\ell s \operatorname{Re}[(B_3 - D_3)(B_4^* + B_5^*)] \\
&\pm \frac{2}{3r} \lambda m_{B_c}^2 (1 - r - s) [\operatorname{Re}(B_1 B_2^*) - \operatorname{Re}(D_1 D_2^*)] \mp \frac{1}{3r} (\lambda + 12rs) [|B_1|^2 - |D_1|^2] \\
&\mp \frac{256}{3} \lambda m_{B_c}^2 m_\ell \left(\operatorname{Re}[A_1^*(C_T \mp C_{TE})g] - \operatorname{Re}[C_1^*(C_T \pm C_{TE})g] \right) \\
&+ \frac{4}{3r} \lambda^2 m_{B_c}^4 m_\ell \left(\operatorname{Re}[B_2^*(C_T \mp 4C_{TE})B_7] + \operatorname{Re}[D_2^*(C_T \pm 4C_{TE})B_7] \right) \\
&- \frac{8}{3r} \lambda m_{B_c}^2 m_\ell (1 - r - s) \left(\operatorname{Re}[B_2^*(C_T \mp 4C_{TE})B_6] + \operatorname{Re}[D_2^*(C_T \pm 4C_{TE})B_6] \right) \\
&- \frac{4}{3r} \lambda m_{B_c}^2 m_\ell (1 - r - s) \left(\operatorname{Re}[B_1^*(C_T \mp 4C_{TE})B_7] + \operatorname{Re}[D_1^*(C_T \pm 4C_{TE})B_7] \right) \\
&+ \frac{8}{3r} (\lambda + 12rs) m_\ell \left(\operatorname{Re}[B_1^*(C_T \mp 4C_{TE})B_6] + \operatorname{Re}[D_1^*(C_T \pm 4C_{TE})B_6] \right) \\
&- \frac{16}{3r} m_\ell [\lambda + 12r(1 - r)] \left(\operatorname{Re}[B_1^*(C_T \mp 4C_{TE})g] + \operatorname{Re}[D_1^*(C_T \pm 4C_{TE})g] \right) \quad (12) \\
&+ \frac{16}{3r} \lambda m_{B_c}^2 m_\ell (1 + 3r - s) \left(\operatorname{Re}[B_2^*(C_T \mp 4C_{TE})g] + \operatorname{Re}[D_2^*(C_T \pm 4C_{TE})g] \right) \\
&+ \frac{16}{3r} \lambda^2 m_{B_c}^6 s |B_7|^2 \operatorname{Re}(C_T C_{TE}^*) \\
&+ \frac{64}{3r} (\lambda + 12rs) m_{B_c}^2 s |B_6|^2 \operatorname{Re}(C_T C_{TE}^*)
\end{aligned}$$

$$\begin{aligned}
& - \frac{64}{3r} \lambda m_{B_c}^4 s (1 - r - s) \operatorname{Re}(B_6 B_7^*) \operatorname{Re}(C_T C_{TE}^*) \\
& + \frac{128}{3r} \lambda m_{B_c}^4 s (1 + 3r - s) \operatorname{Re}(B_7 g^*) \operatorname{Re}(C_T C_{TE}^*) \\
& - \frac{256}{3r} m_{B_c}^2 s [\lambda + 12r(1 - r)] \operatorname{Re}(B_6 g^*) \operatorname{Re}(C_T C_{TE}^*) \\
& + \frac{256}{3r} m_{B_c}^2 [\lambda(4r + s) + 12r(1 - r)^2] |g|^2 \operatorname{Re}(C_T C_{TE}^*) \Big\} ,
\end{aligned}$$

where Δ is the term inside curly brackets of Eq. (11).

Similarly, we find for the transverse polarization P_T^\pm

$$\begin{aligned}
P_T^- &= \frac{\pi}{\Delta} m_{B_c} \sqrt{s\lambda} \Big\{ - 8m_{B_c}^2 m_\ell \operatorname{Re}[(A_1 + C_1)(B_1^* + D_1^*)] \\
& + \frac{1}{r} m_{B_c}^2 m_\ell (1 + 3r - s) [\operatorname{Re}(B_1 D_2^*) - \operatorname{Re}(B_2 D_1^*)] \\
& + \frac{1}{rs} m_\ell (1 - r - s) [|B_1|^2 - |D_1|^2] \\
& + \frac{2}{rs} m_\ell^2 (1 - r - s) [\operatorname{Re}(B_1 B_5^*) - \operatorname{Re}(D_1 B_4^*)] \\
& - \frac{1}{r} m_{B_c}^2 m_\ell (1 - r - s) \operatorname{Re}[(B_1 + D_1)(B_3^* - D_3^*)] \\
& - \frac{2}{rs} m_{B_c}^2 m_\ell^2 \lambda [\operatorname{Re}(B_2 B_5^*) - \operatorname{Re}(D_2 B_4^*)] \\
& + \frac{1}{rs} m_{B_c}^4 m_\ell (1 - r) \lambda [|B_2|^2 - |D_2|^2] + \frac{1}{r} m_{B_c}^4 m_\ell \lambda \operatorname{Re}[(B_2 + D_2)(B_3^* - D_3^*)] \\
& - \frac{1}{rs} m_{B_c}^2 m_\ell [\lambda + (1 - r - s)(1 - r)] [\operatorname{Re}(B_1 B_2^*) - \operatorname{Re}(D_1 D_2^*)] \\
& + \frac{1}{rs} (1 - r - s) (2m_\ell^2 - m_{B_c}^2 s) [\operatorname{Re}(B_1 B_4^*) - \operatorname{Re}(D_1 B_5^*)] \\
& + \frac{1}{rs} m_{B_c}^2 \lambda (2m_\ell^2 - m_{B_c}^2 s) [\operatorname{Re}(D_2 B_5^*) - \operatorname{Re}(B_2 B_4^*)] \\
& - \frac{16}{rs} \lambda m_{B_c}^2 m_\ell^2 \operatorname{Re}[(B_1 - D_1)(B_7 C_{TE})^*] \\
& + \frac{16}{rs} \lambda m_{B_c}^4 m_\ell^2 (1 - r) \operatorname{Re}[(B_2 - D_2)(B_7 C_{TE})^*] \\
& + \frac{8}{r} \lambda m_{B_c}^4 m_\ell \operatorname{Re}[(B_4 - B_5)(B_7 C_{TE})^*] \\
& + \frac{16}{r} \lambda m_{B_c}^4 m_\ell^2 \operatorname{Re}[(B_3 - D_3)(B_7 C_{TE})^*] \\
& + \frac{32}{rs} m_\ell^2 (1 - r - s) \operatorname{Re}[(B_1 - D_1)(B_6 C_{TE})^*] \\
& - \frac{32}{rs} m_{B_c}^2 m_\ell^2 (1 - r) (1 - r - s) \operatorname{Re}[(B_2 - D_2)(B_6 C_{TE})^*] \\
& - \frac{16}{r} m_{B_c}^2 m_\ell (1 - r - s) \operatorname{Re}[(B_4 - B_5)(B_6 C_{TE})^*] \\
& - \frac{32}{r} m_{B_c}^2 m_\ell^2 (1 - r - s) \operatorname{Re}[(B_3 - D_3)(B_6 C_{TE})^*] \Big\} \tag{13}
\end{aligned}$$

$$\begin{aligned}
& - 16m_{B_c}^2 \left(4m_\ell^2 \operatorname{Re}[A_1^*(C_T + 2C_{TE})B_6] - m_{B_c}^2 s \operatorname{Re}[A_1^*(C_T - 2C_{TE})B_6] \right) \\
& + 16m_{B_c}^2 \left(4m_\ell^2 \operatorname{Re}[C_1^*(C_T - 2C_{TE})B_6] - m_{B_c}^2 s \operatorname{Re}[C_1^*(C_T + 2C_{TE})B_6] \right) \\
& + \frac{32}{s} m_{B_c}^2 (1-r) \left(4m_\ell^2 \operatorname{Re}[A_1^*(C_T + 2C_{TE})g] - m_{B_c}^2 s \operatorname{Re}[A_1^*(C_T - 2C_{TE})g] \right) \\
& - \frac{32}{s} m_{B_c}^2 (1-r) \left(4m_\ell^2 \operatorname{Re}[C_1^*(C_T - 2C_{TE})g] - m_{B_c}^2 s \operatorname{Re}[C_1^*(C_T + 2C_{TE})g] \right) \\
& + \frac{64}{rs} m_{B_c}^2 m_\ell^2 (1-r)(1+3r-s) \operatorname{Re}[(B_2 - D_2)(gC_{TE})^*] \\
& + \frac{64}{r} m_{B_c}^2 m_\ell^2 (1+3r-s) \operatorname{Re}[(B_3 - D_3)(gC_{TE})^*] \\
& + \frac{32}{r} m_{B_c}^2 m_\ell (1+3r-s) \operatorname{Re}[(B_4 - B_5)(gC_{TE})^*] \\
& + \frac{64}{rs} [m_{B_c}^2 rs - m_\ell^2 (1+7r-s)] \operatorname{Re}[(B_1 - D_1)(gC_{TE})^*] \\
& - \frac{32}{s} (4m_\ell^2 + m_{B_c}^2 s) \operatorname{Re}[(B_1 + D_1)(gC_T)^*] \\
& - 2048m_{B_c}^2 m_\ell \operatorname{Re}[(C_T g)(B_6 C_{TE})^*] \\
& + \frac{4096}{s} m_{B_c}^2 m_\ell (1-r) |g|^2 \operatorname{Re}(C_T C_{TE}^*) \Big\} ,
\end{aligned}$$

and

$$\begin{aligned}
P_T^+ &= \frac{\pi}{\Delta} m_{B_c} \sqrt{s\lambda} \Big\{ - 8m_{B_c}^2 m_\ell \operatorname{Re}[(A_1 + C_1)(B_1^* + D_1^*)] \\
& - \frac{1}{r} m_{B_c}^2 m_\ell (1+3r-s) [\operatorname{Re}(B_1 D_2^*) - \operatorname{Re}(B_2 D_1^*)] \\
& - \frac{1}{rs} m_\ell (1-r-s) [|B_1|^2 - |D_1|^2] \\
& + \frac{1}{rs} (2m_\ell^2 - m_{B_c}^2 s) (1-r-s) [\operatorname{Re}(B_1 B_5^*) - \operatorname{Re}(D_1 B_4^*)] \\
& + \frac{1}{r} m_{B_c}^2 m_\ell (1-r-s) \operatorname{Re}[(B_1 + D_1)(B_3^* - D_3^*)] \\
& - \frac{1}{rs} m_{B_c}^2 \lambda (2m_\ell^2 - m_{B_c}^2 s) [\operatorname{Re}(B_2 B_5^*) - \operatorname{Re}(D_2 B_4^*)] \\
& - \frac{1}{rs} m_{B_c}^4 m_\ell (1-r) \lambda [|B_2|^2 - |D_2|^2] - \frac{1}{r} m_{B_c}^4 m_\ell \lambda \operatorname{Re}[(B_2 + D_2)(B_3^* - D_3^*)] \\
& + \frac{1}{rs} m_{B_c}^2 m_\ell [\lambda + (1-r-s)(1-r)] [\operatorname{Re}(B_1 B_2^*) - \operatorname{Re}(D_1 D_2^*)] \\
& + \frac{2}{rs} m_\ell^2 (1-r-s) [\operatorname{Re}(B_1 B_4^*) - \operatorname{Re}(D_1 B_5^*)] \\
& + \frac{2}{rs} m_{B_c}^2 m_\ell^2 \lambda [\operatorname{Re}(D_2 B_5^*) - \operatorname{Re}(B_2 B_4^*)] \\
& + \frac{16}{rs} \lambda m_{B_c}^2 m_\ell^2 \operatorname{Re}[(B_1 - D_1)(B_7 C_{TE})^*] \\
& - \frac{16}{rs} \lambda m_{B_c}^4 m_\ell^2 (1-r) \operatorname{Re}[(B_2 - D_2)(B_7 C_{TE})^*] \Big\}
\end{aligned}$$

$$\begin{aligned}
& - \frac{8}{r} \lambda m_{B_c}^4 m_\ell \operatorname{Re}[(B_4 - B_5)(B_7 C_{TE})^*] \\
& - \frac{16}{r} \lambda m_{B_c}^4 m_\ell^2 \operatorname{Re}[(B_3 - D_3)(B_7 C_{TE})^*] \\
& - \frac{32}{rs} m_\ell^2 (1 - r - s) \operatorname{Re}[(B_1 - D_1)(B_6 C_{TE})^*] \\
& + \frac{32}{rs} m_{B_c}^2 m_\ell^2 (1 - r)(1 - r - s) \operatorname{Re}[(B_2 - D_2)(B_6 C_{TE})^*] \\
& + \frac{16}{r} m_{B_c}^2 m_\ell (1 - r - s) \operatorname{Re}[(B_4 - B_5)(B_6 C_{TE})^*] \\
& + \frac{32}{r} m_{B_c}^2 m_\ell^2 (1 - r - s) \operatorname{Re}[(B_3 - D_3)(B_6 C_{TE})^*] \\
& + 16 m_{B_c}^2 \left(4 m_\ell^2 \operatorname{Re}[A_1^*(C_T - 2C_{TE})B_6] - m_{B_c}^2 s \operatorname{Re}[A_1^*(C_T + 2C_{TE})B_6] \right) \\
& - 16 m_{B_c}^2 \left(4 m_\ell^2 \operatorname{Re}[C_1^*(C_T + 2C_{TE})B_6] - m_{B_c}^2 s \operatorname{Re}[C_1^*(C_T - 2C_{TE})B_6] \right) \\
& - \frac{32}{s} m_{B_c}^2 (1 - r) \left(4 m_\ell^2 \operatorname{Re}[A_1^*(C_T - 2C_{TE})g] - m_{B_c}^2 s \operatorname{Re}[A_1^*(C_T + 2C_{TE})g] \right) \\
& + \frac{32}{s} m_{B_c}^2 (1 - r) \left(4 m_\ell^2 \operatorname{Re}[C_1^*(C_T + 2C_{TE})g] - m_{B_c}^2 s \operatorname{Re}[C_1^*(C_T - 2C_{TE})g] \right) \\
& - \frac{64}{rs} m_{B_c}^2 m_\ell^2 (1 - r)(1 + 3r - s) \operatorname{Re}[(B_2 - D_2)(g C_{TE})^*] \\
& - \frac{64}{r} m_{B_c}^2 m_\ell^2 (1 + 3r - s) \operatorname{Re}[(B_3 - D_3)(g C_{TE})^*] \\
& - \frac{32}{r} m_{B_c}^2 m_\ell (1 + 3r - s) \operatorname{Re}[(B_4 - B_5)(g C_{TE})^*] \\
& - \frac{64}{rs} [m_{B_c}^2 rs - m_\ell^2 (1 + 7r - s)] \operatorname{Re}[(B_1 - D_1)(g C_{TE})^*] \\
& - \frac{32}{s} (4 m_\ell^2 + m_{B_c}^2 s) \operatorname{Re}[(B_1 + D_1)(g C_T)^*] \\
& - 2048 m_{B_c}^2 m_\ell \operatorname{Re}[(C_T g)(B_6 C_{TE})^*] \\
& + \frac{4096}{s} m_{B_c}^2 m_\ell (1 - r) |g|^2 \operatorname{Re}(C_T C_{TE}^*) \Big\} .
\end{aligned} \tag{14}$$

Finally for normal asymmetries we get

$$\begin{aligned}
P_N^- &= \frac{1}{\Delta} \pi v m_{B_c}^3 \sqrt{s\lambda} \Big\{ 8 m_\ell \operatorname{Im}[(B_1^* C_1) + (A_1^* D_1)] \\
& - \frac{1}{r} m_{B_c}^2 \lambda \operatorname{Im}[(B_2^* B_4) + (D_2^* B_5)] \\
& + \frac{1}{r} m_{B_c}^2 m_\ell \lambda \operatorname{Im}[(B_2 - D_2)(B_3^* - D_3^*)] \\
& - \frac{1}{r} m_\ell (1 + 3r - s) \operatorname{Im}[(B_1 - D_1)(B_2^* - D_2^*)] \\
& + \frac{1}{r} (1 - r - s) \operatorname{Im}[(B_1^* B_4) + (D_1^* B_5)] \Big\}
\end{aligned}$$

$$\begin{aligned}
& - \frac{1}{r} m_\ell (1 - r - s) \text{Im}[(B_1 - D_1)(B_3^* - D_3^*)] \\
& - \frac{8}{r} m_{B_c}^2 m_\ell \lambda \text{Im}[(B_4 + B_5)(B_7 C_{TE})^*] \\
& + \frac{16}{r} m_\ell (1 - r - s) \text{Im}[(B_4 + B_5)(B_6 C_{TE})^*] \\
& - \frac{32}{r} m_\ell (1 + 3r - s) \text{Im}[(B_4 + B_5)(g C_{TE})^*] \\
& - 16 m_{B_c}^2 s \left(\text{Im}[A_1^*(C_T - 2C_{TE})B_6] + \text{Im}[C_1^*(C_T + 2C_{TE})B_6] \right) \\
& + 32 m_{B_c}^2 (1 - r) \left(\text{Im}[A_1^*(C_T - 2C_{TE})g] + \text{Im}[C_1^*(C_T + 2C_{TE})g] \right) \\
& + 32 \left(\text{Im}[B_1^*(C_T - 2C_{TE})g] - \text{Im}[D_1^*(C_T + 2C_{TE})g] \right) \\
& + 512 m_\ell \left(|C_T|^2 - 4 |C_{TE}|^2 \right) \text{Im}(B_6^* g) \Big\} ,
\end{aligned} \tag{15}$$

and

$$\begin{aligned}
P_N^+ &= \frac{1}{\Delta} \pi v m_{B_c}^3 \sqrt{s \lambda} \Big\{ - 8 m_\ell \text{Im}[(B_1^* C_1) + (A_1^* D_1)] \\
& + \frac{1}{r} m_{B_c}^2 \lambda \text{Im}[(B_2^* B_5) + (D_2^* B_4)] \\
& + \frac{1}{r} m_{B_c}^2 m_\ell \lambda \text{Im}[(B_2 - D_2)(B_3^* - D_3^*)] \\
& - \frac{1}{r} m_\ell (1 + 3r - s) \text{Im}[(B_1 - D_1)(B_2^* - D_2^*)] \\
& - \frac{1}{r} (1 - r - s) \text{Im}[(B_1^* B_5) + (D_1^* B_4)] \\
& - \frac{1}{r} m_\ell (1 - r - s) \text{Im}[(B_1 - D_1)(B_3^* - D_3^*)] \\
& + \frac{8}{r} m_{B_c}^2 m_\ell \lambda \text{Im}[(B_4 + B_5)(B_7 C_{TE})^*] \\
& - \frac{16}{r} m_\ell (1 - r - s) \text{Im}[(B_4 + B_5)(B_6 C_{TE})^*] \\
& + \frac{32}{r} m_\ell (1 + 3r - s) \text{Im}[(B_4 + B_5)(g C_{TE})^*] \\
& - 16 m_{B_c}^2 s \left(\text{Im}[A_1^*(C_T + 2C_{TE})B_6] + \text{Im}[C_1^*(C_T - 2C_{TE})B_6] \right) \\
& + 32 m_{B_c}^2 (1 - r) \left(\text{Im}[A_1^*(C_T + 2C_{TE})g] + \text{Im}[C_1^*(C_T - 2C_{TE})g] \right) \\
& - 32 \left(\text{Im}[B_1^*(C_T + 2C_{TE})g] - \text{Im}[D_1^*(C_T - 2C_{TE})g] \right) \\
& + 512 m_\ell \left(|C_T|^2 - 4 |C_{TE}|^2 \right) \text{Im}(B_6^* g) \Big\} .
\end{aligned} \tag{16}$$

From Eqs. (12)-(16), we observe that for longitudinal and normal polarizations, the difference between ℓ^+ and ℓ^- lepton asymmetries results from the scalar and tensor type interactions. Similar situation takes place for transverse polarization asymmetries in the

$m_\ell \rightarrow 0$ limit. From this, we can conclude that their experimental study may provide essential information about new physics.

Another source of useful information about new physics can be a combined analysis of the lepton and antilepton polarizations, since in the SM $P_L^- + P_L^+ = 0$, $P_N^- + P_N^+ = 0$ and $P_T^- - P_T^+ \approx 0$ [11]. Using Eqs. (12)-(16) we get

$$\begin{aligned}
P_L^- + P_L^+ &= \frac{4}{\Delta} m_{B_c}^2 v \left\{ \frac{2}{r} m_\ell \lambda \operatorname{Re}[(B_1 - D_1)(B_4^* + B_5^*)] \right. \\
&- \frac{2}{r} m_{B_c}^2 m_\ell \lambda (1 - r) \operatorname{Re}[(B_2 - D_2)(B_4^* + B_5^*)] \\
&- \frac{1}{r} m_{B_c}^2 s \lambda (|B_4|^2 - |B_5|^2) - \frac{2}{r} m_{B_c}^2 m_\ell s \lambda \operatorname{Re}[(B_3 - D_3)(B_4^* + B_5^*)] \\
&+ \frac{8}{3r} m_{B_c}^4 m_\ell \lambda^2 \operatorname{Re}[(B_2 + D_2)(B_7 C_T)^*] \\
&+ \frac{32}{3r} m_{B_c}^6 s \lambda^2 |B_7|^2 \operatorname{Re}(C_T C_{TE}^*) \\
&- \frac{8}{3r} m_{B_c}^2 m_\ell \lambda (1 - r - s) \operatorname{Re}[(B_1 + D_1)(B_7 C_T)^*] \\
&- \frac{16}{3r} m_{B_c}^2 m_\ell \lambda (1 - r - s) \operatorname{Re}[(B_2 + D_2)(B_6 C_T)^*] \\
&- \frac{128}{3r} m_{B_c}^4 s \lambda (1 - r - s) \operatorname{Re}(B_6 B_7^*) \operatorname{Re}(C_T C_{TE}^*) \\
&+ \frac{16}{3r} m_\ell (\lambda + 12rs) \operatorname{Re}[(B_1 + D_1)(B_6 C_T)^*] \\
&+ \frac{128}{3r} m_{B_c}^2 s (\lambda + 12rs) |B_6|^2 \operatorname{Re}(C_T C_{TE}^*) \\
&+ \frac{512}{3r} m_{B_c}^2 [\lambda(4r + s) + 12r(1 - r)^2] |g|^2 \operatorname{Re}(C_T C_{TE}^*) \\
&- \frac{512}{3r} m_{B_c}^2 s [\lambda + 12r(1 - r)] \operatorname{Re}(g B_6^*) \operatorname{Re}(C_T C_{TE}^*) \\
&+ \frac{256}{3r} m_{B_c}^4 s \lambda (1 + 3r - s) \operatorname{Re}(g B_7^*) \operatorname{Re}(C_T C_{TE}^*) \\
&+ \frac{512}{3} m_{B_c}^2 m_\ell \lambda \operatorname{Re}[(A_1 + C_1)(g C_{TE})^*] \\
&- \frac{32}{3r} m_\ell [\lambda + 12r(1 - r)] \operatorname{Re}[(B_1 + D_1)(g C_T)^*] \\
&+ \left. \frac{32}{3r} m_{B_c}^2 m_\ell \lambda (1 + 3r - s) \operatorname{Re}[(B_2 + D_2)(g C_T)^*] \right\}. \tag{17}
\end{aligned}$$

For the case of transverse polarization, it is the difference of the lepton and antilepton polarizations that is relevant and it can be calculated from Eqs. (13) and (14)

$$\begin{aligned}
P_T^- - P_T^+ &= \frac{\pi}{\Delta} m_{B_c} \sqrt{s \lambda} \left\{ \frac{2}{rs} m_{B_c}^4 m_\ell (1 - r) \lambda [|B_2|^2 - |D_2|^2] \right. \\
&+ \left. \frac{1}{r} m_{B_c}^4 \lambda \operatorname{Re}[(B_2 + D_2)(B_4^* - B_5^*)] \right\}
\end{aligned}$$

$$\begin{aligned}
& + \frac{2}{r} m_{B_c}^4 m_\ell \lambda \operatorname{Re}[(B_2 + D_2)(B_3^* - D_3^*)] \\
& + \frac{2}{r} m_{B_c}^2 m_\ell (1 + 3r - s) [\operatorname{Re}(B_1 D_2^*) - \operatorname{Re}(B_2 D_1^*)] \\
& + \frac{2}{rs} m_\ell (1 - r - s) [|B_1|^2 - |D_1|^2] \\
& - \frac{1}{r} m_{B_c}^2 (1 - r - s) \operatorname{Re}[(B_1 + D_1)(B_4^* - B_5^*)] \\
& - \frac{2}{r} m_{B_c}^2 m_\ell (1 - r - s) \operatorname{Re}[(B_1 + D_1)(B_3^* - D_3^*)] \\
& - \frac{2}{rs} m_{B_c}^2 m_\ell [\lambda + (1 - r)(1 - r - s)] [\operatorname{Re}(B_1 B_2^*) - \operatorname{Re}(D_1 D_2^*)] \\
& - \frac{32}{rs} m_{B_c}^2 m_\ell^2 \lambda \operatorname{Re}[(B_1 - D_1)(B_7 C_{TE})^*] \\
& + \frac{32}{rs} m_{B_c}^4 m_\ell^2 \lambda (1 - r) \operatorname{Re}[(B_2 - D_2)(B_7 C_{TE})^*] \\
& + \frac{16}{r} m_{B_c}^4 m_\ell \lambda \operatorname{Re}[(B_4 - B_5)(B_7 C_{TE})^*] \\
& + \frac{32}{r} m_{B_c}^4 m_\ell^2 \lambda \operatorname{Re}[(B_3 - D_3)(B_7 C_{TE})^*] \\
& + \frac{64}{rs} m_\ell^2 (1 - r - s) \operatorname{Re}[(B_1 - D_1)(B_6 C_{TE})^*] \\
& - \frac{64}{rs} m_{B_c}^2 m_\ell^2 (1 - r)(1 - r - s) \operatorname{Re}[(B_2 - D_2)(B_6 C_{TE})^*] \\
& - \frac{32}{r} m_{B_c}^2 m_\ell (1 - r - s) \operatorname{Re}[(B_4 - B_5)(B_6 C_{TE})^*] \\
& - \frac{64}{r} m_{B_c}^2 m_\ell^2 (1 - r - s) \operatorname{Re}[(B_3 - D_3)(B_6 C_{TE})^*] \\
& + 32 m_{B_c}^4 s v^2 \operatorname{Re}[(A_1 - C_1)(B_6 C_T)^*] \\
& + \frac{64}{r} m_{B_c}^2 m_\ell (1 + 3r - s) \operatorname{Re}[(B_4 - B_5)(g C_{TE})^*] \\
& - 64 m_{B_c}^4 (1 - r) v^2 \operatorname{Re}[(A_1 - C_1)(g C_T)^*] \\
& + \frac{128}{rs} [m_{B_c}^2 r s - m_\ell^2 (1 + 7r - s)] \operatorname{Re}[(B_1 - D_1)(g C_{TE})^*] \\
& + \frac{128}{rs} m_{B_c}^2 m_\ell^2 (1 - r)(1 + 3r - s) \operatorname{Re}[(B_2 - D_2)(g C_{TE})^*] \\
& + \frac{128}{r} m_{B_c}^2 m_\ell^2 (1 + 3r - s) \operatorname{Re}[(B_3 - D_3)(g C_{TE})^*] \Big\} .
\end{aligned} \tag{18}$$

In the same manner it follows from Eqs. (15) and (16)

$$\begin{aligned}
P_N^- + P_N^+ &= \frac{1}{\Delta} \pi v m_{B_c}^3 \sqrt{s \lambda} \Big\{ - \frac{2}{r} m_\ell (1 + 3r - s) \operatorname{Im}[(B_1 - D_1)(B_2^* - D_2^*)] \\
& - \frac{2}{r} m_\ell (1 - r - s) \operatorname{Im}[(B_1 - D_1)(B_3^* - D_3^*)] \\
& - \frac{1}{r} (1 - r - s) \operatorname{Im}[(B_1 - D_1)(B_4^* - B_5^*)]
\end{aligned}$$

$$\begin{aligned}
& + \frac{2}{r} m_{B_c}^2 m_\ell \lambda \text{Im}[(B_2 - D_2)(B_3^* - D_3^*)] \\
& + \frac{1}{r} m_{B_c}^2 \lambda \text{Im}[(B_2 - D_2)(B_4^* - B_5^*)] \\
& + 32 m_{B_c}^2 s \text{Im}[(A_1 + C_1)(B_6 C_T)^*] \\
& + 1024 m_\ell (|C_T|^2 - |4C_{TE}|^2) \text{Im}(B_6^* g) \\
& - 64 m_{B_c}^2 (1 - r) \text{Im}[(A_1 + C_1)(g C_T)^*] \\
& + 128 \text{Im}[(B_1 + D_1)(g C_{TE})^*] \Big\} .
\end{aligned} \tag{19}$$

It can be seen from Eq. (17) that in $P_L^- + P_L^+$ the terms containing the SM contribution, i.e., terms containing C_{BR} , C_{SL} , C_{LL}^{tot} and C_{LR}^{tot} completely cancels. For this reason, a measurement of the nonzero value of $P_L^- + P_L^+$ in future experiments, may be an indication of the discovery of new physics beyond the SM.

Before going into the details of our numerical analysis we like to note a final point about the numerical calculations of the polarization asymmetries. As seen from the Eqs.(12)-(19), all the expressions of the lepton polarizations depend on both $s = q^2/m_{B_c}^2$ and the new Wilson coefficients. However, it may be experimentally easier to study the dependence of the polarizations of each lepton on the new Wilson coefficients only. For this reason we eliminate s dependence by considering their averaged forms over the allowed kinematical region. The averaged lepton polarizations are defined as

$$\langle P_i \rangle = \frac{\int_{(2m_\ell/m_{B_c})^2}^{(1-m_{D_s^*}/m_{B_c})^2} P_i \frac{d\mathcal{B}}{ds} ds}{\int_{(2m_\ell/m_{B_c})^2}^{(1-m_{D_s^*}/m_{B_c})^2} \frac{d\mathcal{B}}{ds} ds} . \tag{20}$$

4 Numerical analysis and discussion

We here present our numerical analysis about the branching ratios and averaged polarization asymmetries $\langle P_L^- \rangle$, $\langle P_T^- \rangle$ and $\langle P_N^- \rangle$ of ℓ^- for the $B_c \rightarrow D^* \ell^+ \ell^-$ decays with $\ell = \mu, \tau$, as well as the lepton-antilepton combined asymmetries $\langle P_L^- + P_L^+ \rangle$, $\langle P_T^- - P_T^+ \rangle$ and $\langle P_N^- + P_N^+ \rangle$. We first give the input parameters used in our numerical analysis :

$$\begin{aligned}
& m_{B_c} = 6.50 \text{ GeV} , \ m_{D_s^*} = 2.112 \text{ GeV} \ m_b = 4.8 \text{ GeV} , \ m_\mu = 0.105 \text{ GeV} , \ m_\tau = 1.77 \text{ GeV} , \\
& |V_{tb} V_{ts}^*| = 0.0385 , \ \alpha^{-1} = 129 , \ G_F = 1.17 \times 10^{-5} \text{ GeV}^{-2} \\
& \tau_{B_c} = 0.46 \times 10^{-12} \text{ s} .
\end{aligned} \tag{21}$$

The values of the individual Wilson coefficients that appear in the SM are listed in Table (1). The values for the mass and the lifetime of the B_c meson given above in Eq.(21) were reported by CDF Collaboration [18]. Recently, CDF quoted a new value $m_{B_c} = 6.2857 \pm 0.0053 \pm 0.0012 \text{ GeV}$ [19]. Also, D0 has observed B_c and reported the preliminary results $m_{B_c} = 5.95_{-0.13}^{+0.14} \pm 0.34 \text{ GeV}$ and $\tau_{B_c} = 0.45_{-0.10}^{+0.12} \pm 0.12$ [20]. However, we observed that our numerical results are not sensitive to the numerical values of m_{B_c} more than 3–5%.

| C_1 | C_2 | C_3 | C_4 | C_5 | C_6 | C_7^{eff} | C_9 | C_{10} |
|--------|--------|--------|--------|--------|--------|--------------------|--------|----------|
| -0.248 | +1.107 | +0.011 | -0.026 | +0.007 | -0.031 | -0.313 | +4.344 | -4.624 |

Table 1: Values of the SM Wilson coefficients at $\mu \sim m_b$ scale.

We note that the value of the Wilson coefficient C_9^{eff} in Table (1) corresponds only to the short-distance contributions. C_9^{eff} also receives long-distance contributions due to conversion of the real $\bar{c}c$ into lepton pair $\ell^+\ell^-$ and they are usually absorbed into a redefinition of the short-distance Wilson coefficients:

$$C_9^{\text{eff}}(\mu) = C_9(\mu) + Y(\mu) , \quad (22)$$

where

$$\begin{aligned} Y(\mu) &= Y_{\text{reson}} + h(y, s)[3C_1(\mu) + C_2(\mu) + 3C_3(\mu) + C_4(\mu) + 3C_5(\mu) + C_6(\mu)] \\ &- \frac{1}{2}h(1, s)(4C_3(\mu) + 4C_4(\mu) + 3C_5(\mu) + C_6(\mu)) \\ &- \frac{1}{2}h(0, s)[C_3(\mu) + 3C_4(\mu)] \\ &+ \frac{2}{9}(3C_3(\mu) + C_4(\mu) + 3C_5(\mu) + C_6(\mu)) , \end{aligned} \quad (23)$$

with $y = m_c/m_b$, and the functions $h(y, s)$ arises from the one loop contributions of the four quark operators O_1, \dots, O_6 and their explicit forms can be found in [21]. It is possible to parametrize the resonance $\bar{c}c$ contribution $Y_{\text{reson}}(s)$ in Eq.(23) using a Breit-Wigner shape with normalizations fixed by data which is given by [22]

$$\begin{aligned} Y_{\text{reson}}(s) &= -\frac{3}{\alpha_{em}^2}\kappa \sum_{V_i=\psi_i} \frac{\pi\Gamma(V_i \rightarrow \ell^+\ell^-)m_{V_i}}{sm_{B_c}^2 - m_{V_i} + im_{V_i}\Gamma_{V_i}} \\ &\times [(3C_1(\mu) + C_2(\mu) + 3C_3(\mu) + C_4(\mu) + 3C_5(\mu) + C_6(\mu))] , \end{aligned} \quad (24)$$

where the phenomenological parameter κ is usually taken as ~ 2.3 .

As for the values of the new Wilson coefficients, they are the free parameters in this work, but it is possible to establish ranges out of experimentally measured branching ratios of the semileptonic and also purely leptonic rare B-meson decays

$$\begin{aligned} BR(B \rightarrow K \ell^+\ell^-) &= (0.75_{-0.21}^{+0.25} \pm 0.09) \times 10^{-6} , \\ BR(B \rightarrow K^* \mu^+\mu^-) &= (0.9_{-0.9}^{+1.3} \pm 0.1) \times 10^{-6} , \end{aligned}$$

reported by Belle and Babar collaborations [23]. It is now also available an upper bound of pure leptonic rare B-decays in the $B^0 \rightarrow \mu^+\mu^-$ mode [24]:

$$BR(B^0 \rightarrow \mu^+\mu^-) \leq 2.0 \times 10^{-7} .$$

Being in accordance with this upper limit and also the above mentioned measurements of the branching ratios for the semileptonic rare B-decays, we take in this work all new Wilson coefficients as real and varying in the region $-4 \leq C_X \leq 4$.

Among the new Wilson coefficients that appear in Eq.(1), those related to the helicity-flipped counter-parts of the SM operators, namely, C_{RL} and C_{RR} , vanish in all models with minimal flavor violation in the limit $m_s \rightarrow 0$. However, there are some MSSM scenarios in which there are finite contributions from these vector operators even for a vanishing s-quark mass. In addition, scalar type interactions can also contribute through the neutral Higgs diagrams in e.g. multi-Higgs doublet models and MSSM for some regions of the parameter spaces of the related models. In literature there exists studies to establish ranges out of constraints under various precision measurements for these coefficients (see e.g. [25]) and our choice for the range of the new Wilson coefficients are in agreement with these calculations.

To make some numerical predictions, we also need the explicit forms of the form factors $A_0, A_+, A_-, V, a_0, a_+$ and g . In our work we have used the results of [6], in which q^2 dependencies of the form factors are given as

$$F(q^2) = \frac{F(0)}{(1 - as + bs^2)^2},$$

where the values of parameters $F(0)$, a and b for the $B_c \rightarrow D_s^*$ decay are listed in Table 2.

| | $F(0)$ | a | b |
|-------|--------|------|-------|
| A_0 | 0.279 | 1.30 | 0.149 |
| A_+ | 0.156 | 2.16 | 1.15 |
| A_- | -0.321 | 2.41 | 1.51 |
| V | 0.290 | 2.40 | 1.49 |
| a_0 | 0.178 | 1.21 | 0.125 |
| a_+ | 0.178 | 2.14 | 1.14 |
| g | 0.179 | 2.51 | 1.67 |

Table 2: B_c meson decay form factors in a relativistic constituent quark model.

We present the results of our analysis in a series of figures. Before the discussion of these figures, we give our SM predictions for the longitudinal, transverse and the normal components of the lepton polarizations for $B_c \rightarrow D_s^* \ell^+ \ell^-$ decay for μ (τ) channel for reference:

$$\begin{aligned} \langle P_L^- \rangle &= 0.6211 (0.6321), \\ \langle P_T^- \rangle &= 0.0017 (0.0468), \\ \langle P_N^- \rangle &= -0.0837 (-0.17). \end{aligned}$$

Figs. (1) and (2) give dependence of the integrated branching ratio (BR) on the new Wilson coefficients for the $B_c \rightarrow D_s^* \mu^+ \mu^-$ and $B_c \rightarrow D_s^* \tau^+ \tau^-$ decays, respectively. From

these figures we see that BR depends strongly on the tensor interactions and weakly on the vector interactions, while it is completely insensitive to the scalar type of interactions. It is also clear from these figures that dependence of the BR on the new Wilson coefficients is symmetric with respect to the zero point for the muon final state, but such a symmetry is not observed for the tau final state for the tensor interactions.

In Figs. (3) and (4), we present the dependence of averaged longitudinal polarization $\langle P_L^- \rangle$ of ℓ^- and the combined averaged $\langle P_L^- + P_L^+ \rangle$ for $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay on the new Wilson coefficients. We observe that $\langle P_L^- \rangle$ is more sensitive to the existence of the tensor type interactions while the combined average $\langle P_L^- + P_L^+ \rangle$ is to that of scalar type interactions only. The fact that $\langle P_L^- + P_L^+ \rangle$ does not exhibit any dependence on the vector type of interactions are already an expected result since vector type interactions are cancelled when the longitudinal polarization asymmetry of the lepton and antilepton is considered together. We also note that the values of $\langle P_L^- \rangle$ becomes substantially different from the SM value (at $C_X = 0$) as C_X becomes different from zero, which indicates that measurement of the longitudinal lepton polarization in $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay can be very useful to investigate new physics beyond the SM. From Fig. (3), we see that, the contributions coming from all types of interactions to $\langle P_L^- \rangle$ are positive and it is an increasing (decreasing) functions of both C_T and C_{TE} for their negative (positive) values. We observe from Fig. (4) that $\langle P_L^- + P_L^+ \rangle$ becomes zero at $C_X = 0$, which conforms the SM results, and its dependence on C_X is symmetric with respect to this zero value. It is also interesting to note that $\langle P_L^- + P_L^+ \rangle$ is positive for all values of C_{LRLR} and C_{RLLR} , while it is negative for remaining scalar type interactions.

Figs. (5) and (6) are the same as Figs. (3) and (4), but for $B_c \rightarrow D_s^* \tau^+ \tau^-$. Similar to the muon case, $\langle P_L^- \rangle$ is more sensitive to the tensor interactions than others. Contributions to $\langle P_L^- \rangle$ from all type of interactions are positive for all values of C_X except for C_{TE} : in region $0.25 \lesssim C_{TE} < 4$, $\langle P_L^- \rangle$ changes the sign and becomes negative. As for the main interesting point in Fig. (6), although $\langle P_L^- + P_L^+ \rangle$ for $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay depends only on scalar interactions, for $B_c \rightarrow D_s^* \tau^+ \tau^-$ decay it is also and very sensitively dependent on tensor type of interactions. It is also interesting to note that $\langle P_L^- + P_L^+ \rangle$ changes sign: it takes positive (negative) values for the negative (positive) values of C_T and C_{TE} . Thus, one can provide valuable information about the new physics by determining the sign and the magnitude of $\langle P_L^- + P_L^+ \rangle$. We finally note that as in case of muon final state, in tau final state too, $\langle P_L^- + P_L^+ \rangle$ becomes zero at $C_X = 0$ and confirms the SM result.

In Figs. (7) and (8), we present the dependence of averaged transverse polarization $\langle P_T^- \rangle$ of ℓ^- and the combined averaged $\langle P_T^- - P_T^+ \rangle$ for $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay on the new Wilson coefficients. From these figures, it is seen that for the $\langle P_T^- \rangle$, there appears strong dependence on tensor and scalar interactions and also a weak dependence on vector interactions. On the other hand, vector contributions to the $\langle P_T^- - P_T^+ \rangle$ is negligible and main contribution comes from the tensor interactions and C_{LRRL} and C_{RLRL} components of the scalar interactions. As seen from Figs. (7) and (8), both $\langle P_T^- \rangle$ and $\langle P_T^- - P_T^+ \rangle$ are positive (negative) for the negative (positive) values of C_T and C_{RLRL} , except in a region about the zero values of the coefficients, $-1 \lesssim C_X \lesssim 1$, while their behavior with respect to C_{TE} and C_{LRRL} are opposite. Therefore, determination of the sign and magnitude of these observables can also give useful information about existence of new physics.

Figs. (9) and (10) are the same as Figs. (7) and (8), but for $B_c \rightarrow D_s^* \tau^+ \tau^-$. We see

from Fig. (9) that the $\langle P_T^- \rangle$ is quite sensitive to all types of interactions and behavior of scalar interaction is identical for coefficients C_{LRRL}, C_{LRLR} and C_{RLLR}, C_{RLRL} in pairs. It can be seen from Fig. (10) that although tensor and scalar interactions are dominant for $\langle P_T^- - P_T^+ \rangle$, the dependence of vector interactions are also more sizable as compared with the case of muon final state. In addition, change in sign of $\langle P_T^- \rangle$ and $\langle P_T^- - P_T^+ \rangle$ are observed depending on the change in the tensor and scalar interaction coefficients, whose measure may provide useful tools for new physics.

In Figs.(11) and (12), we present the dependence of averaged normal polarization $\langle P_N^- \rangle$ of ℓ^- and the combined averaged $\langle P_N^- + P_N^+ \rangle$ for $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay on the new Wilson coefficients. We see from Fig. (11) that $\langle P_N^- \rangle$ strongly depends on the tensor interactions. Its dependence on the scalar type of interactions is moderate and identical for the coefficients C_{LRRL}, C_{LRLR} and C_{RLLR}, C_{RLRL} in pairs. As seen from Fig. (12), the behavior of $\langle P_N^- + P_N^+ \rangle$ is determined by the tensor interactions only. We also observe that $\langle P_N^- + P_N^+ \rangle$ is positive (negative) when $C_T < 0$ ($C_T > 0$) while its behavior with respect to C_{TE} is opposite. Further, $\langle P_N^- + P_N^+ \rangle$ becomes zero at $C_X = 0$ as expected in the SM.

Figs. (13) and (14) are the same as Figs. (11) and (12), but for $B_c \rightarrow D_s^* \tau^+ \tau^-$. We first note that as being opposite to the muon final state case, here $\langle P_N^- \rangle$ depends on all types of interactions although dependence on tensor interaction is stronger. We also observe that $\langle P_N^- \rangle$ always takes the positive values except when $C_{TE} \lesssim -0.25$ and $C_T \gtrsim 2$. As seen from Fig. (14), $\langle P_N^- + P_N^+ \rangle$ depends only on the tensor interactions and its behavior is the same as that of the muon final state case.

In conclusion, we present the most general analysis of the lepton polarization asymmetries in the rare $B_c \rightarrow D_s^* \ell^+ \ell^-$ decay using the general, model independent form of the effective Hamiltonian. The dependence of the longitudinal, transversal and normal polarization asymmetries of ℓ^- and their combined asymmetries on the new Wilson coefficients are studied. It is found that the lepton polarization asymmetries are very sensitive to the existence of the tensor and scalar type interactions. Moreover, $\langle P_T \rangle$ and $\langle P_N \rangle$ change their signs as the new Wilson coefficients vary in the region of interest. This conclusion is valid also for the combined polarization effects $\langle P_L^- + P_L^+ \rangle$, $\langle P_T^- - P_T^+ \rangle$ and $\langle P_N^- + P_N^+ \rangle$ for the same decay channel. It is well known that in the SM, $\langle P_L^- + P_L^+ \rangle = \langle P_T^- - P_T^+ \rangle = \langle P_N^- + P_N^+ \rangle \simeq 0$ in the limit $m_\ell \rightarrow 0$. Therefore any deviation from this relation and determination of the sign of polarization is decisive and effective tool in looking for new physics beyond the SM.

Acknowledgments

This work was partially supported by Mersin University under Grant No: BAP-FEF-FB (UOY) 2006-3.

References

- [1] See, for a recent review, A. Ali, *Int. J. Mod. Phys.* **A20** (2005) 5080.
- [2] CLEO Collaboration, M. S. Alam, *et. al.*, *Phys. Rev. Lett.* **74** (1995) 2885.
- [3] P. Colangelo and F. De Fazio, *Phys. Rev.* **D61** (2000) 034012.
- [4] M. A. Ivanov, J. G. Körner, and P. Santorelli, *Phys. Rev.* **D63** (2001) 074010.
- [5] C. Q. Geng, C. W. Hwang, and C. C. Liu, *Phys. Rev.* **D65** (2002) 094037.
- [6] A. Faessler, Th. Gutsche, M. A. Ivanov, J. G. Körner, and V. E. Lyubovitskij, *Eur.Phys.J.direct* **C4** (2002) 18.
- [7] T. M. Aliev, M. Savcı, *Phys. Lett.* **B452** (1999) 318;
T. M. Aliev, A. Özpıneci, H. Koru and M. Savcı, *Phys. Lett.* **B410** (1997) 216.
- [8] Y. G. Kim, P. Ko and J. S. Lee, *Nucl. Phys.* **B544** (1999) 64.
- [9] J. L. Hewett, *Phys. Rev.* **D53** (1996) 4964.
- [10] F. Krüger and L. M. Sehgal, *Phys. Lett.* **B380** (1996) 199.
- [11] S. Fukae, C. S. Kim and T. Yoshikawa, *Phys. Rev.* **D61** (2000) 074015.
- [12] T. M. Aliev, M. Savcı, *Phys. Lett.* **B481** (2000) 275.
- [13] T. M. Aliev, D. A. Demir, M. Savcı, *Phys. Rev.* **D62** (2000) 074016.
- [14] F. Krüger and E. Lunghi, *Phys. Rev.* **D63** (2001) 014013.
- [15] Qi-Shu Yan, Chao-Shang Huang, Liao Wei, Shou-Hua Zhu, *Phys. Rev.* **D62** (2000) 094023.
- [16] D. Guetta and E. Nardi, *Phys. Rev.* **D58** (1998) 012001.
- [17] P. Colangelo, F. De Fazio, P. Santorelli and E. Scrimieri, *Phys. Rev.* **D53** (1996) 3672;
Erratum, *ibid.* **D57** (1998) 3186.
- [18] F. Abe *et al.*, CDF Collaboration, *Phys. Rev.* **D58**, (1998) 112004; *Phys. Rev. Lett.* **81**, (1998) 2432.
- [19] A. Abulencia *et al.*, CDF Collaboration, *Phys. Rev. Lett.* **96**, (2006) 082002.
- [20] M. D. Corcoran, CDF Collaboration, hep-ex/0506061.
- [21] A. J. Buras and M. Münz, *Phys. Rev. D* **52** (1995) 186; M. Misiak, *Nucl. Phys.*, **B 393**(1993) 23; erratum *ibid.* **B 439** (1995) 461.
- [22] A. Ali, T. Mannel and T. Morozumi, *Phys. Lett.* **B 273** (1991) 505;

- [23] K. Abe, *et al.*, BELLE Collaboration, *Phys. Rev. Lett.* **88** (2002) 021801; B. Aubert, *et al.*, BaBar Collaboration, *Phys. Rev. Lett.* **88** (2002) 241801.
- [24] V. Halyo, *hep-ex/0207010*.
- [25] Chao-Shang Huang and Xiao-Hong Wu, *Nucl. Phys.* **B 665** (2003) 304.




Figure 1: The dependence of the integrated branching ratio for the $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay on the new Wilson coefficients.




Figure 2: The dependence of the integrated branching ratio for the $B_c \rightarrow D_s^* \tau^+ \tau^-$ decay on the new Wilson coefficients.




Figure 3: The dependence of the averaged longitudinal polarization $\langle P_L^- \rangle$ of ℓ^- for the $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay on the new Wilson coefficients.




Figure 4: The dependence of the combined averaged longitudinal lepton polarization $\langle P_L^- + P_L^+ \rangle$ for the $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay on the new Wilson coefficients.




Figure 5: The same as Fig. (3), but for the $B_c \rightarrow D_s^* \tau^+ \tau^-$ decay.




Figure 6: The same as Fig. (4), but for the $B_c \rightarrow D_s^* \tau^+ \tau^-$ decay.




Figure 7: The dependence of the averaged transverse polarization $\langle P_T^- \rangle$ of ℓ^- for the $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay on the new Wilson coefficients.




Figure 8: The dependence of the combined averaged transverse lepton polarization $\langle P_T^- - P_T^+ \rangle$ for the $B_c \rightarrow D_s^* \gamma \mu^+ \mu^-$ decay on the new Wilson coefficients.




Figure 9: The same as Fig. (7), but for the $B_c \rightarrow D_s^* \tau^+ \tau^-$ decay.




Figure 10: The same as Fig. (8), but for the $B_c \rightarrow D_s^* \tau^+ \tau^-$ decay.




Figure 11: The dependence of the averaged normal polarization $\langle P_N^- \rangle$ of ℓ^- for the $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay on the new Wilson coefficients.




Figure 12: The dependence of the combined averaged normal lepton polarization $\langle P_N^- + P_N^+ \rangle$ for the $B_c \rightarrow D_s^* \mu^+ \mu^-$ decay on the new Wilson coefficients.




Figure 13: The same as Fig.(10), but for the $B_c \rightarrow D_s^* \tau^+ \tau^-$ decay.




Figure 14: The same as Fig.(11), but for the $B_c \rightarrow D_s^* \tau^+ \tau^-$ decay.