

The complex relation between production and scattering amplitudes

Eef van Beveren

*Centro de Física Teórica, Departamento de Física, Universidade de Coimbra
P-3004-516 Coimbra, Portugal*

<http://cft.fis.uc.pt/eef>

George Rupp

*Centro de Física das Interações Fundamentais, Instituto Superior Técnico
Universidade Técnica de Lisboa, Edifício Ciência, P-1049-001 Lisboa, Portugal*

george@ist.utl.pt

PACS number(s): 11.80.Gw, 11.55.Ds, 13.75.Lb, 12.39.Pn

February 9, 2022

Abstract

The unitarity relation $\Im m(A) = T^* A$ is derived for a three-body production amplitude A that consists of a complex linear combination of elements of the two-body scattering amplitude T . We conclude that the unitarity relation does not impose a realness condition on the coefficients in the expansion of A in terms of T .

Under the spectator assumption, we deduced in Ref. [1] that the three-particle production amplitude A consists of a complex linear combination of elastic and inelastic matrix elements of the two-body scattering amplitude T . Furthermore, in Ref. [2] we showed that such a two-particle production amplitude can reasonably describe experiment in an energy region where no additional resonances from possible rescattering with the spectator particle exist. Moreover, no need to treat a sizable fraction of the experimental signal as background was noticed.

The result of Ref. [1] agrees to some extent with the expression proposed in Refs. [3, 4]. Like in our Ref. [1], the authors of Ref. [4] based their ansatz on the OZI rule [5] and the spectator picture, finding that the production amplitude can be written as a linear combination of the elastic and inelastic two-body scattering amplitudes, with coefficients that do not carry any singularities, but are rather supposed to depend smoothly on the total CM energy of the system.

However, Ref. [4] concluded from the unitarity relation

$$\Im m(A) = T^* A \quad (1)$$

that the production amplitude must be given by a *real* linear combination of the elements of the transition matrix. A similar conclusion, based on a K -matrix parametrisation, can be found in

Ref. [6]. In contrast, we arrive at a different conclusion, namely that the coefficients must be *complex*, in agreement with experiment [7–9] as well as with the work of the Ishidas [10, 11].

Relation (1), which can also be found in Ref. [12], basically stems from the operator relations $AV = (1 + TG)V = V + TGV = T$, the symmetry of T , the realness of V , and the unitarity of $1 + 2iT$, which gives $\Im m(A)V = \Im m(AV) = \Im m(T) = T^*T = T^*AV$. This leads, for non-singular potentials V , to relation (1).

Now we shall show that A and T satisfy the unitarity relation (1) *despite* the complexness of the coefficients in the expansion of A in terms of T . Thereto, we are going to strip the expressions of Ref. [1] of all details which might obscure the simplicity of our arguments. Hence, let $Z_k(E)$ ($k = 1, 2, \dots, n$) represent a vector of complex non-singular expressions¹, being smooth functions of the energy E , where n represents the number of coupled scattering channels under consideration, and let the relation between A and T be given by²

$$A_k = \Re e(Z_k) + i \sum_{\ell} Z_{\ell} T_{k\ell} . \quad (2)$$

We then find for the imaginary part of the production amplitude

$$\Im m(A_k) = \frac{1}{2i} (A_k - A_k^*) = \sum_{\ell} \{ \Re e(Z_{\ell}) \Re e(T_{k\ell}) - \Im m(Z_{\ell}) \Im m(T_{k\ell}) \} . \quad (3)$$

Next, we substitute on the right-hand side of Eq. (3) the identity $\Re e(T_{k\ell}) = T_{k\ell}^* + i\Im m(T_{k\ell})$, and furthermore insert the unitarity condition for T , *i.e.*, $\Im m(T_{k\ell}) = \sum_{\ell'} T_{\ell'\ell} T_{k\ell'}^*$, so as to obtain

$$\begin{aligned} \Im m(A_k) &= \sum_{\ell} \{ \Re e(Z_{\ell}) (T_{k\ell}^* + i\Im m(T_{k\ell})) - \Im m(Z_{\ell}) \Im m(T_{k\ell}) \} \\ &= \sum_{\ell} \left\{ \Re e(Z_{\ell}) T_{k\ell}^* + i Z_{\ell} \sum_{\ell'} T_{\ell'\ell} T_{k\ell'}^* \right\} . \end{aligned} \quad (4)$$

Finally, we interchange ℓ and ℓ' in the second term on the right-hand side of Eq. (4), leaving us, also using Eq. (2), with

$$\Im m(A_k) = \sum_{\ell} T_{k\ell}^* \left\{ \Re e(Z_{\ell}) + i \sum_{\ell'} Z_{\ell'} T_{\ell\ell'} \right\} = \sum_{\ell} T_{k\ell}^* A_{\ell} . \quad (5)$$

This completes the proof that A , as defined in Eq. (2), satisfies the unitarity condition (1). Consequently, relation (1) does not impose a realness condition on the coefficients in Eq. (2).

The first term on the right-hand side of relation (2) was not considered in Refs. [3, 4, 6]. However, in the works of Graves-Morris [13] and Aitchison & collaborators [14–16], the possible existence of an additional real contribution was anticipated. In Refs. [1, 2], this follows straightforwardly from the reasonable assumption that a produced meson pair originates from an initial $q\bar{q}$ pair. As a consequence, the observed phenomenological necessity [9] to employ complex coefficients in experimental analyses of production processes does *not* allow by itself to draw conclusions on the inevitability of including rescattering diagrams with the spectator particle in theoretical approaches.

¹ In Appendix A we give the precise relation between the expressions used in Ref. [1] and Z_k .

²Note that for $\Re e(Z_k) = 0$ one obtains an expansion with real coefficients, as in Refs. [3, 4].

Acknowledgements

We wish to thank I. J. R. Aitchison, D. V. Bugg, C. Hanhart and M. R. Pennington for useful discussions. This work was supported in part by the *Fundação para a Ciência e a Tecnologia* of the *Ministério da Ciência, Tecnologia e Ensino Superior* of Portugal, under contract PDCT/FP/63907/2005.

A Precise definition of $Z_k(E)$

In Ref. [1] we discussed the partial-wave expansion of the amplitudes for two-meson production — together with a spectator particle — and scattering, assuming $q\bar{q}$ pair creation. Hence, the coefficients bear reference to the partial wave ℓ and the flavor content α of the quark pair. We obtained [1] the following relation between production and scattering partial-wave amplitudes:

$$A_{\alpha i}^{(\ell)} = g_{\alpha i} j_{\ell}(p_i r_0) \sqrt{\mu_i p_i} + i \sum_{\nu} g_{\alpha \nu} \sqrt{\mu_{\nu} p_{\nu}} h_{\ell}^{(1)}(p_{\nu} r_0) T_{i\nu}^{(\ell)}. \quad (6)$$

Accordingly, we must define

$$Z_{\alpha k}^{(\ell)}(E) = g_{\alpha k} h_{\ell}^{(1)}(p_k r_0) \sqrt{\mu_k p_k}. \quad (7)$$

In the latter equations, j_{ℓ} and $h_{\ell}^{(1)}$ stand for the spherical Bessel function and Hankel function of the first kind, respectively. These are smooth functions of the total CM energy, just like μ_k and p_k , which are the reduced mass and relative linear momentum of the two-meson system in the k -th channel, respectively. The constants $g_{\alpha k}$ stand for the intensities of the $q\bar{q} \rightarrow MM$ couplings. A distance scale ~ 0.6 fm (for light quarks) is represented by r_0 . In the text we have stripped Z of a reference to ℓ and α .

Note, moreover, as can be easily seen from expressions (2) and (6), that the singularity structures of the production and scattering amplitudes are identical, since $\Re(Z_k)$, which is proportional to the spherical Bessel function in Eq. (6), is a smooth function of the total invariant mass.

References

- [1] E. van Beveren and G. Rupp, *Relating multichannel scattering and production amplitudes in a microscopic OZI-based model*, arXiv:0706.4119.
- [2] E. van Beveren and G. Rupp, *S-wave and P-wave pion-pion and kaon-pion contributions to three-body decay processes in the Resonance-Spectrum Expansion*, J. Phys. G **34**, 1789 (2007) [arXiv:hep-ph/0703286].
- [3] S. U. Chung, J. Brose, R. Hackmann, E. Klempt, S. Spanier and C. Strassburger, *Partial wave analysis in K matrix formalism*, Annalen Phys. **507**, 404 (1995).
- [4] K. L. Au, D. Morgan and M. R. Pennington, *Meson dynamics beyond the quark model: A study of final-state interactions*, Phys. Rev. D **35**, 1633 (1987).

- [5] S. Okubo, *Φ meson and unitary symmetry model*, Phys. Lett. **5**, 165 (1963);
 G. Zweig, *An SU_3 model for strong interaction symmetry and its breaking*, CERN Reports TH-401 and TH-412 (1963);
 see also *Developments in the Quark Theory of Hadrons*, Vol. 1, pp. 22–101 (1981), edited by
 D. B. Lichtenberg and S. P. Rosen;
 J. Iizuka, K. Okada and O. Shito, *Systematics and phenomenology of boson mass levels (3)*,
 Prog. Theor. Phys. **35**, 1061 (1966).
- [6] M. Boglione and M. R. Pennington, *Towards a model independent determination of the $\phi \rightarrow f_0\gamma$ coupling*, Eur. Phys. J. C **30**, 503 (2003) [arXiv:hep-ph/0303200].
- [7] J. M. Link *et al.* [FOCUS Collaboration], *Dalitz plot analysis of D_s^+ and D^+ decay to $\pi^+\pi^-\pi^+$ using the K -matrix formalism*, Phys. Lett. B **585**, 200 (2004) [arXiv:hep-ex/0312040].
- [8] P. Dini [FOCUS Collaboration], *Dalitz plot analyses from FOCUS*, Int. J. Mod. Phys. A **20**, 482 (2005).
- [9] J. M. Link *et al.* [FOCUS Collaboration] and M. R. Pennington, *Dalitz plot analysis of the $D^+ \rightarrow K^-\pi^+\pi^+$ decay in the FOCUS experiment*, Phys. Lett. B **653**, 1 (2007) [arXiv:0705.2248 [hep-ex]].
- [10] M. Ishida, S. Ishida and T. Ishida, *Relation between scattering and production amplitudes: Concerning σ particle in $\pi\pi$ system*, Prog. Theor. Phys. **99**, 1031 (1998) [arXiv:hep-ph/9805319].
- [11] M. Ishida, *The present status on sigma and kappa meson properties: Relation between scattering and production amplitudes*, AIP Conf. Proc. **688**, 18 (2004).
- [12] U. G. Meißner and J. A. Oller, *$J/\psi \rightarrow \phi\pi\pi$ (K anti- K) decays, chiral dynamics and OZI violation*, Nucl. Phys. A **679**, 671 (2001) [arXiv:hep-ph/0005253].
- [13] P. R. Graves-Morris, *S-matrix theory of final-state interactions*, Nuovo Cim. A **50**, 681 (1967).
- [14] M. G. Bowler, M. A. V. Game, I. J. R. Aitchison and J. B. Dainton, *Diffraction dissociation, the Deck mechanism and diffractive resonance production*, Nucl. Phys. B **97**, 227 (1975).
- [15] I. J. R. Aitchison and M. G. Bowler, *Rescattering effects in the Deck model*, J. Phys. G **3**, 1503 (1977).
- [16] I. J. R. Aitchison, *Are there important unitarity corrections to the isobar model?*, Phys. Lett. B **84**, 349 (1979).