

Associated production of the scalars and new gauge bosons from a little Higgs model at the LHC

Chong-Xing Yue, Nan Zhang, Li Ding, and Shi-Hai Zhu

Department of Physics, Liaoning Normal University, Dalian 116029, P. R. China *

February 9, 2022

Abstract

The littlest Higgs model with T-parity (LHT model) predicts the existence of the T-odd scalars (ϕ^\pm , ϕ^0 , and ϕ^p). We consider production of these new particles associated with T-odd gauge bosons at the LHC . We find that the partonic process $q\bar{q}' \rightarrow \phi^+ B_H$ can generate a number of the characteristic signal events with a charged lepton and large missing energy at the LHC .

PACS number:12.60.Cn, 14.80.Cp, 14.70.Pw

*E-mail: cxyue@lnnu.edu.cn

1. Introduction

The mechanism of electroweak symmetry breaking ($EWSB$) and origin of the fermion mass remain unknown in the current particle physics despite the success of the standard model (SM) tested by high energy experimental data. In next year, the *CERN* Large Hadron Collider (LHC) will begin operation, which has an increase of a factor of seven in energy and a factor of 100 in luminosity over the Fermilab Tevatron. The LHC can be seen as the machine extremely well suit to the study of these problems and is expected to directly probe possible new physics beyond the SM up to a few TeV, which might provide some striking evidence of new physics.

Little Higgs theory [1] is proposed as an interesting solution to the so called hierarchy problem of the SM and can be regarded as one of the important candidates for new physics beyond the SM . Among these models, the littlest Higgs (LH) model [2] is one of the simplest and phenomenological viable models, which has all essential features. Most phenomenological analysis about the original little Higgs models are given in the context of the LH model [3, 4]. They have shown that some of new particles predicted by the LH model can generate characteristic signatures at present and in future collider experiments.

It is well known that the LH model suffers from severe constraints from precision electroweak measurement, which could only be satisfied by fine-tuning the model parameters [5]. To avoid this serious problem, a new discrete symmetry called T-parity has been introduced. The LH model with T-parity (LHT model) [6] is one of the attractive little Higgs models, which satisfies the electroweak precision data in most of the parameter space. In the LHT model, all of the SM particles are assigned with an T-even parity, while all of the new particles are assigned with an T-odd parity, except for the little Higgs partner of the top quark. As a consequence, all of the T-odd particles can only be generated in pairs. The electroweak precision data allow for a relatively low value of new particle mass scale $f \sim 500 GeV$ [7, 8]. Thus, the LHT model can produce rich phenomenology at present and in future high energy experiments [9, 10, 11].

Most of the previous works on studying the phenomenology of the LHT model focus their attention on the new fermions (T-odd fermions or T-even heavy top quark T^+) and

new gauge bosons (W_H^\pm, B_H, Z_H). In this letter, we will discuss single production of the scalars (ϕ^0, ϕ^p , and ϕ^\pm) predicted by the *LHT* model via possible processes at the *LHC* with the center-of-mass (c. m.) energy $\sqrt{S} = 14TeV$. In section 2, we briefly summarize the mass and coupling formula related to our calculation. Single production of the new scalars in association with new gauge bosons at the *LHC* are discussed in section 3. The relevant phenomenological analysis are also given in this section. In section 4, we give our conclusions.

2. The relevant mass and coupling formula

Similar to the *LH* model, the *LHT* model [6] is based on an $SU(5)/SO(5)$ global symmetry breaking pattern. A subgroup $[SU(2)_1 \times U(1)_1] \times [SU(2)_2 \times U(1)_2]$ of the $SU(5)$ global symmetry is gauged, and at the scale f it is broken into the *SM* electroweak symmetry $SU(2)_L \times U(1)_Y$. T-parity is an automorphism that exchanges the $[SU(2)_1 \times U(1)_1]$ and $[SU(2)_2 \times U(1)_2]$ gauge symmetries. Consequently, four of the 14 Nambu-Goldstone(*NG*) bosons are eaten by the T-odd heavy gauge bosons (W_H^\pm, Z_H, B_H) associated with the broken symmetry. The remaining *NG* bosons decompose into a T-even doublet H , considered to be the *SM* Higgs doublet, and a complex T-odd $SU(2)$ triplet Φ . There is a relation between the triplet and doublet Higgs boson masses, which is approximately expressed as:

$$M_\Phi = \frac{\sqrt{2}m_H}{\nu}f, \quad (1)$$

where $\nu=246\text{GeV}$ is the electroweak scale and f is the scale of the gauge symmetry breaking of the *LHT* model.

After *EWSB*, at the order of ν^2/f^2 , the mass expressions of the T-odd heavy gauge bosons B_H, Z_H and W_H are given by

$$M_{B_H} = \frac{g'f}{\sqrt{5}}[1 - \frac{5\nu^2}{8f^2}], \quad M_{Z_H} \approx M_{W_H} = gf[1 - \frac{\nu^2}{8f^2}]. \quad (2)$$

Here g' and g are the *SM* $U(1)_Y$ and $SU(2)_L$ gauge coupling constants, respectively. Because of the smallness of g' , the T-odd gauge boson B_H is the lightest T-odd particle, which can be seen as an attractive dark matter candidate [9].

To implement T-parity in the fermion sector, one introduces two $SU(2)$ fermion doublets for each SM fermion doublet [6, 7, 8]. The T-even linear combination is associated with the SM $SU(2)$ doublet, while the T-odd combination is associated with its T-parity partner. Its mass is at order of the symmetry breaking scale f . Assuming universal and flavor diagonal Yukawa coupling constant k , the masses of the T-odd fermions u_- and d_- can be approximately written as:

$$M_{u_-} \approx \sqrt{2}kf(1 - \frac{\nu^2}{8f^2}), \quad M_{d_-} \approx \sqrt{2}kf. \quad (3)$$

The couplings of these new particles to the SM particles (T-even), which are related our calculation, can be approximately written as [8, 11]:

$$\phi^0 Z^\mu B_H^\nu : \frac{ig'^2}{2\sqrt{2}S_W} \frac{\nu^2}{f} g^{\mu\nu}, \quad \phi^0 Z^\mu Z_H^\nu : -\frac{ig^2}{2\sqrt{2}C_W} \frac{\nu^2}{f} g^{\mu\nu}; \quad (4)$$

$$\phi^p W_H^{-\mu} W^{+\nu} : \frac{g^2}{3\sqrt{2}} \frac{\nu^2}{f} g^{\mu\nu}, \quad \phi^p u^i \bar{u}_-^j : \frac{k}{12} \frac{\nu^2}{f^2} (V_{Hu})_{ij} P_L; \quad (5)$$

$$\phi^+ B_H^\mu W^{-\nu} : \frac{igg'}{4} \frac{\nu^2}{f} g^{\mu\nu}, \quad \phi^+ W_H^{-\mu} Z^\nu : -\frac{ig^2}{12C_W} (1 + 2S_W^2) \frac{\nu^2}{f} g^{\mu\nu}; \quad (6)$$

$$\phi^+ W_H^{-\mu} \gamma^\nu : \frac{ieg}{6} \frac{\nu^2}{f} g^{\mu\nu}, \quad \phi^+ Z_H^\mu W^{-\nu} : \frac{ig^2}{12} \frac{\nu^2}{f} g^{\mu\nu}; \quad (7)$$

$$Z_H^\mu d_-^i \bar{d}^i : \frac{ig}{2} (V_{Hd})_{ij} \gamma^\mu P_L, \quad B_H^\mu d_-^i \bar{d}^i : \frac{ig'}{10} (V_{Hd})_{ij} \gamma^\mu P_L; \quad (8)$$

$$W_H^{+\mu} \bar{u}_-^i d^j : \frac{ig}{\sqrt{2}} (V_{Hd})_{ij} \gamma^\mu P_L, \quad \Phi^+ d_-^i \bar{u}^j : \frac{k}{12} \frac{\nu^2}{f^2} (V_{Hu})_{ij} P_L. \quad (9)$$

Here ϕ^+ , ϕ^- , ϕ^0 , and ϕ^p are the components of the triplet scalar Φ , which have the same mass at the leading order. k is the Yukawa-type coupling constant for the mirror fermions. $S_W = \sin \theta_W$, θ_W is the Weinberg angle. $(V_{Hu})_{ij}$ and $(V_{Hd})_{ij}$ are the matrix elements of the CKM-like unitary mixing matrices V_{Hu} and V_{Hd} , respectively, which satisfy $V_{Hu}^\dagger V_{Hd} = V_{CKM}$.

In the following section, we will use above Feynman rules to calculate production of the T-odd scalars (ϕ^p , ϕ^0 , ϕ^+ and ϕ^-) in association with new gauge bosons at the LHC .

3. Associated production of the T-odd scalars and new gauge bosons at the LHC

In the LHT model [6, 8], T-parity explicitly forbids the tree level contributions from the new gauge bosons to the observables involving only the SM particles and forbids

the interactions that induce triplet vacuum expectation value (VEV). Corrections to electroweak observables are only generated at one loop level or beyond, which are naturally small. As a result, the electroweak precision data allow for a relatively low value of the symmetry breaking scale $f \sim 500 GeV$ [7]. Thus, the new particle mass might be relatively low, which induces that the T-partners of the SM particles can be copiously produced at the LHC as well as future e^+e^- linear collider (ILC) [8, 10, 11]. In this section, we will consider production of the T-odd scalars in association with the T-odd new gauge bosons at the LHC .




Figure 1: Feynman diagrams for the production of the new scalars at the LHC .

From Feynman rules given in section 2, we can see that the T-odd scalars can be produced at the LHC via the following partonic processes:

$$q\bar{q} \rightarrow \phi^0 Z_H, \quad q\bar{q} \rightarrow \phi^0 B_H, \quad q\bar{q} \rightarrow \phi^- W_H^+; \quad (10)$$

$$q\bar{q}' \rightarrow \phi^p W_H^+, \quad q\bar{q}' \rightarrow \phi^+ Z_H, \quad qq' \rightarrow \phi^+ B_H. \quad (11)$$

The relevant Feynman diagrams are shown in Fig.1. One can see from Fig.1 that the partonic processes $q\bar{q} \rightarrow \phi V_H$ ($V_H = Z_H, B_H$, or W_H) can only be induced via the s-channel exchange of the SM gauge boson Z or γ , while the partonic processes $q\bar{q}' \rightarrow \phi V_H$ can proceed by the s-channel exchange of the SM gauge bosons W^\pm and the t-channel exchange of the T-odd mirror fermion q_- .

Considering that the coupling ϕqq_- is proportion to the factor ν^2/f^2 , the t-channel contributions to the process $q\bar{q}' \rightarrow \phi V_H$ should be suppressed by the factor ν^4/f^4 . However, to completely consider production cross section of the T-odd scalars at the LHC , we will include both the s-channel and t-channel contributions in our numerical calculation.

At the LHC , production of the Higgs boson in association with W or Z boson, $pp \rightarrow WH/ZH + X$, is one of the interesting processes for detecting the SM Higgs boson H with a mass below about $135 GeV$, where H decays into $b\bar{b}$ final states are dominate [12]. At leading order, production of a Higgs boson in association with W or Z boson, $pp \rightarrow VH + X$ ($V = W, Z$), proceeds through $q\bar{q}'$ annihilation $q\bar{q}' \rightarrow V^* \rightarrow VH$ [13]. Production of the T-odd scalar in association with a new gauge boson at the LHC is similar with WZ/ZH production. Thus, we can easily give the production cross section $\sigma(\phi V_H)$ for the process $pp \rightarrow q\bar{q}' + X \rightarrow \phi V_H + X$ ($V_H = W_H, Z_H, B_H$) at the LHC . In our numerical calculation, we will use $CTEQ6L$ parton distribution functions ($PDF's$) [14] for the quark distribution functions.

To obtain numerical results, we need to specify the relevant SM input parameters, which are $\alpha_e = 1/128.8$, $S_W^2 = 0.2315$, $M_W = 80.45 GeV$, and $M_Z = 91.187 GeV$ [15]. Considering the experimental constraints on the SM Higgs boson mass m_H [16], we will take $m_H = 120 GeV$. Except these SM input parameters, the cross section $\sigma(\phi V_H)$ for associated production of the new scalar and the new gauge boson depends on the free parameters k , $(V_{Hu})_{ij}$, $(V_{Hd})_{ij}$, and f . The matrix element $(V_{Hd})_{ij}$ can be determined in flavor violating processes and the matrix element $(V_{Hu})_{ij}$ is then determined through $V_{Hu} = V_{Hd} V_{CKM}^+$. Following Ref.[11], we will take $k = 1$, $(V_{Hd})_{ij}(i = j) = \frac{1}{\sqrt{2}}$, and assume

the scale parameter f as a free parameter in our numerical estimation.

Our numerical results are shown in Fig.2 and Fig.3. From these figures, one can see that the production cross section quickly decreases as f increases. This is because the masses of the T-odd scalars and gauge bosons increase as f increases. Since the new gauge boson B_H is lighter than the new gauge boson Z_H or W_H^\pm , the production cross sections of the new scalars in association with B_H are generally larger than those for the new scalars in association with gauge boson Z_H or W_H^\pm . However, in most of the parameter space of the *LHT* model, all of the production cross sections are small. Even for the process $q\bar{q}' \rightarrow \phi^+ B_H$, its production cross section is only in the range of $1.014fb \sim 0.024fb$ for $500GeV \leq f \leq 800GeV$. With the high luminosity option of the *LHC*, around $300fb^{-1}$, there will be several and up to hundreds $\phi^+ B_H$ events.




Figure 2: Production cross sections for the partonic processes $q\bar{q}' \rightarrow \phi^+ B_H$ and $q\bar{q} \rightarrow \phi^0 B_H$ as functions of the scale parameter f .

From the discussion given in Sec.2, we can see that the charged scalar ϕ^+ mainly decays to $W^+ B_H$ and there will be $Br(\phi^+ \rightarrow W^+ B_H) \approx 1$ for $m_H = 120GeV$ and $k = 1$ [11].

In the case of W^+ decaying leptonically, the partonic process $q\bar{q}' \rightarrow \phi^+ B_H$ could give rise to the distinct signal events with one charged lepton and large missing energy, i.e. $q\bar{q}' \rightarrow \phi^+ B_H \rightarrow W^+ B_H B_H \rightarrow l^+ \nu B_H B_H$. Thus, as long as $f \leq 800 \text{ GeV}$, there will be several tens of observed $l^+ \nu B_H B_H$ events to be generated at the *LHC* with $\sqrt{S} = 14 \text{ TeV}$ and a yearly integrated luminosity of $\mathcal{L} = 300 \text{ fb}^{-1}$.




Figure 3: Same as Fig.2 but for the partonic processes $q\bar{q} \rightarrow \phi^0 Z_H$, $q\bar{q} \rightarrow \phi^- W_H^+$, $q\bar{q}' \rightarrow \phi^p W_H^+$, and $q\bar{q}' \rightarrow \phi^+ Z_H$.

4. Conclusions

The *LHT* model is one of the attractive little Higgs models, which provides a possible dark matter candidate. In this model, all dangerous tree level contributions to electroweak observables are forbidden by T-parity and hence the corrections to electroweak observables are very small. So the low scale parameter f ($f \geq 500 \text{ GeV}$) is allowed by the electroweak precision data and this model might produce rich phenomenology at present and in future high energy experiments.

In this letter, we consider production of the T-odd scalars (ϕ^\pm , ϕ^0 , and ϕ^p) in associ-

ation with T-odd gauge bosons at the *LHC*. Since all production processes are proceed mainly via s-channel processes with highly virtual gauge boson propagators, the production cross sections are small in most of the parameter space of the *LHT* model. However, the process $q\bar{q}' \rightarrow \phi^+ B_H$ can generate the characteristic signal events with a charged lepton and large missing energy. There will be several tens of observed events to be generated at the *LHC*. Therefore, detecting the signature of ϕ^+ is challenging for the *LHC*, depending on its luminosity.

Acknowledgments

This work was supported in part by the Program for New Century Excellent Talents in University(NCET-04-0290), and the National Natural Science Foundation of China under the Grants No.10675057.

References

- [1] M. Schmaltz and D. Tucker-Smith, *Ann. Rev. Nucl. Part. Sci.* **55**, 229(2005).
- [2] N. Arkani-Hamed, A. G. Cohen, E. Katz, A. E. Nelson, *JHEP* **0207**, 034(2002).
- [3] T. Han, H. E. Logan, and L. T. Wang, *JHEP* **0601**, 099(2006).
- [4] M. Perelstein, *Prog. Part. Nucl. Phys.* **58**, 247(2007).
- [5] J. L. Hewett, F. J. Petriello, and T. G. Rizzo, *JHEP* **0310**, 062(2003); C. Csaki, J. Hubisz, G. D. Kribs, P. Meade, and J. Terning, *Phys. Rev.* **D67**, 115002(2003); C. Csaki et al., *Phys. Rev.* **D68**, 035009(2003); R. Casalbuoni, A. Deandrea, M. Oertel, *JHEP* **0402**, 032(2004); Mu-Chun Chen and S. Dawson, *Phys. Rev.* **D70**, 015003(2004); Chong-Xing Yue and Wei Wang, *Nucl. Phys. B* **683**, 48(2004); W. Kilian and J. Reuter, *Phys. Rev.* **D70**, 015004(2004); T. Gregoire, D. R. Smith, and

- J. G. Wacker, *Phys. Rev. D***69**, 115008(2004); Mu-Chun Chen, *Mod. Phys. Lett. A***21**, 621(2006).
- [6] H. C. Cheng and I. Low, *JHEP* **0309**, 051(2003); *JHEP* **0408**, 061(2004); I. Low, *JHEP* **0410**, 067(2004).
- [7] J. Hubisz, P. Meade, A. Noble, and M. Perelstein, *JHEP* **0601**, 135(2006).
- [8] J. Hubisz, P. Meade, *Phys. Rev. D***71**, 035016(2005).
- [9] A. Birkedal, A. Noble, M. Perelstein, A. Spary, *Phys. Rev. D***74**, 035002(2006); M. Asano et al., *Phys. Rev. D***75**, 063506(2007); C. S. Chen, Kingman Cheung, and T. C. Yuan, *Phys. Lett. B***644**, 158(2007); M. Perelstein A. Spray, *Phys. Rev. D***75**, 083519(2007).
- [10] C. S. Chen, Kingman Cheung, and T. -C. Yuan, *Phys. Lett. B***644**, 158(2007); J. Hubisz, S. J. Lee, and G. Paz, *JHEP* **0606**, 041(2006); M. Blanke et al., *JHEP* **0612**, 003(2006); A. J. Buras, A. Poschenrieder, S. Uhlig, and W. A. Bardeen, *JHEP* **0611**, 062(2006); A. Freitas and D. Wyler, *JHEP* **0611**, 061(2006); C. O. Dib, R. Rosenfeld, and A. Zerwekh, *JHEP* **0605**, 074(2006); H. C. Cheng, I. Low, and L. T. Wang, *Phys. Rev. D***74**, 055001(2006); C. R. Chen, K. Tobe, and C.-P. Yuan, *Phys. Lett. B***640**, 263(2006); D. Choudhury, D. Kumar Ghosh, *JHEP* **0708**, 084(2007); V. Barger, Wai-Yee Keung, Y. Gao, *arXiv:0707.3648*.
- [11] A. Belyaev, Chuan-Ren Chen, K. Tobe, and C.-P. Yuan, *Phys. Rev. D***74**, 115020(2006); M. Blake et al., *JHEP* **0701**, 066(2007).
- [12] For example, see: D. Rainwater, *hep-ph/0702124*.
- [13] S. L. Glashow, D. V. Nanoponlos, and A. Yildiz, *Phys. Rev. D***18**, 1724(1978); Z. Kunszt, Z. Trócsány, and W. J. Stjrling, *Phys. Lett. B***271**, 247(1991).
- [14] J. Pumplin et al., *JHEP* **0207**, 012(2002); D. Stump et al., *JHEP* **0310**, 046(2003).
- [15] W. M. Yao et al.[Particle Data Group], *J. Phys. G***33**, 1(2006).

- [16] The LEP Collaborations, the LEP Electroweak Working Group, and the SLD Heavy Flavour Group, *A Combination of Preliminary Electroweak Measurements and Constraints on the Standard Model*, *hep-ex/0412015*.