# Predictive pion-quark BCS relation and Thornber-Feynman high-T<sub>c</sub> gap

### M.D. Scadron<sup>a</sup> B.A. Green<sup>b</sup> <sup>a</sup>Physics Department, University of Arizona, Tucson, AZ 85721 <sup>b</sup>11912 N. Centaurus PI., Tucson AZ 85737

A pion-quark pairing temperature is defined by a BCS-like relation identified from a quarklevel Goldberger-Treiman relation with a Nambu scalar mass "gap parameter" taken in the lowmass limit. This intuitive relation predicts the associated "experimental" lattice-gauge pairing temperature. The opposite high-mass limit predicts the sigma mass, and notably has a predictive analog in high-T<sub>c</sub> superconductivity in the stable nondispersive energy gap as defined by Thornber-Feynman polaron dynamics. A predictive analog pion-quark BCS relation with a Nambu-mass gap parameter has a further parallel in the nondispersive high-T<sub>c</sub> superconductivity gap as defined from Thornber-Feynman (TF) polaron dynamics [1]. This BCSlike relation for the pion  $\bar{q}q$  binding energy and pair-breaking "temperature", as defined from a quark-level Goldberger-Treiman relation (GTR) in the low-mass limit [1], predicts this latest calculated temperature as follows. The GTR for the pion-nucleon coupling constant  $g_{\pi NN}$  is  $g_{\pi NN}=m_Ng_A/f_{\pi}$  in terms of the nucleon mass  $m_N$  (939 MeV), pion decay constant in the chiral pairing limit  $f_{\pi}$ -90 MeV, and axial current form factor  $g_A$ =1 for structureless quarks in the constituent quark model [2]. For the constituent quark this relation becomes  $g_{\pi qq}=m_q/f_{\pi}$  where  $m_q\approx$ 939 MeV/3≈313 MeV and  $g_A$  is unity for structureless quarks, so an "experimental" two-flavor quark-level GTR is

$$g_{\pi qq} = m_q / f_{\pi} \sim 313 \text{ MeV}/90 \text{ MeV} \approx 3.5.$$
 (1)

Noting that the righthand side of Eq. 1 numerically approximates the BCS ratio, we cast it into "BCS form" by multiplying the numerator and denominator by two, i.e.,

$$g_{\pi qq} \approx 2m_q/2f_{\pi} \sim 3.5.$$
 (2)

The numerator  $2m_q$  is recognized as the scalar meson mass identified from the Nambu scalar mass (Eq. A-3) in the Appendix as an analog superconducting gap  $2\Delta$ . From this approximate BCS ratio  $2\Delta/k_BT_c\sim3.5$  in Eq. 2 for  $T_c\sim3-4$  K we identify the numerator ( $2m_q$ ) with a pair binding energy  $2\Delta$ , where the constituent quark masses are largely taken up in the binding as with the pion [3]. This further suggests the BCS denominator  $2f_\pi$  is an intuitive  $\overline{q}q$  pairing or chiral

restoration "temperature"  $T_c$ , i.e.,  $T_c=2f_{\pi}\sim 180$  MeV [4], which in fact is compatible with the recent "experimental" value of 173±8 MeV obtained from a computer "lattice" calculation for two quark flavors [5]. Notably this intuitive  $T_c$  determination from the quark-level GTR complements the earlier ones in Ref. 4 in the framework of chiral symmetry breaking.

This Nambu scalar mass-pion BCS connection in the low-mass limit  $(2m_q \rightarrow 2\Delta, 2f_\pi \rightarrow T_c)$  is further supported by independent calculations of the BCS ratio and  $g_{\pi qq}$  as analytic expressions that are similar in both magnitude and form. Namely for acoustical phonons the gap ratio becomes [1]

$$2\Delta/k_{\rm B}T_{\rm c} = 2\pi e^{-\gamma} \approx 3.528, \tag{3}$$

where  $\gamma$  is the Euler constant (0.5772), and for the pion-quark coupling constant with color number N<sub>c</sub>=3 we find [6,4]

$$g_{\pi qq} \approx 2\pi / N_c^{1/2} = 2\pi / 3^{1/2} \approx 3.628.$$
 (4)

Note this numerical nearness of Eqs. 3 and 4 to the BCS ratio in Eqs. 1,2.

The above identification of the double quark mass  $2m_q$  in Eq. 2 with the pion  $\overline{q}q$  pairing energy  $2\Delta$  is reinforced by theoretical and experimental determinations of the pion charge radius  $r_{\pi}$  from a "fused" quark pair that accords with the tight-binding massless Nambu-Goldstone pion implicit in the GTR, Eq. 1. Hence  $r_{\pi}$  is specified by a *single* quark mass suggestive of the fused pair as  $r_{\pi}=\hbar c/m_q=197.3$  MeV-fm/313 MeV $\approx$ 0.63 fm ( $\hbar c=197.3$  MeV-fm) [7] from the "common sense" quark mass 313 MeV=m<sub>N</sub>/3 as expected with three constituent non-strange quarks in a nucleus. Similarly, vector meson dominance (VMD) [8] with rho-meson mass 770 MeV specifies a pion charge radius  $6^{1/2}$ (197.3 MeV-fm)/770 MeV $\approx$ 0.63 fm.

Not surprisingly, the numerical closeness of 3.528 and 3.628 in Eqs. 3 and 4 is not accidental. In both cases the energy scales with the momentum as E=p (in units c=1), but for different reasons. For the acoustic-phonon BCS coupling in Eq. 3 the linear scaling is in the low-energy limit due to the large Fermi surface (FS), which follows from the relation  $\Delta E=(p/m)\Delta p$  with constant of proportionality p/m defined from energy-momentum changes  $\Delta E$  and  $\Delta p$ . Hence p/m is approximately constant on the FS because the changes  $\Delta p/p$  are small. Whereas in Eq. 4 the linear scaling is due to tight binding, where the fused pion mass vanishes in the relativistic energy-mass relation  $E^2=p^2+m^2$  as the mass converts to binding energy, so E is approximately p in this (tight binding) limit.

The Nambu scalar mass gap parameter Eq. A-3 in the opposite high-mass limit predicts the sigma mass as shown in the Appendix. The further analog in the flatband high-T<sub>c</sub> superconductivity gap  $2\Delta_{HTC}$  in this limit, paralleling the long-range low-T<sub>c</sub> gap parameter Eq. A-2 [1], is *experimentally* specified by the non-dispersive (massive) energy-constrained TF quasiparticle shift E<sub>LO</sub> (the longitu-dinal-optical (LO) phonon mode energy) [9]. This mobile, stable, gapped quasiparticle in paralleling the high-T<sub>c</sub> particle arises from nonadiabatic (conserved) internal polaron dynamics in such low-symmetry Frohlich media [10] as defined by a "universal" TF mobility [1] (e.g., in photoconducting alkali halides and related transition-metal oxides [12]). Accordingly the optimum, constrained high-T<sub>c</sub> gap analogous to Eq. A-2 is empirically

$$2\Delta_{\rm HTC}=E_{\rm LO},$$
 (5a)

where in the cuprates this gap as defined by angle-resolved photoemission spectroscopy (ARPES) data in fact equals the resonance-softened in-plane Cu-O mode energy  $E_{LO}^{R}$  in satisfying Eq. 5a directly or from combined cuprate data [1], so the above TF gap relation becomes

$$2\Delta_{\rm HTC} = E_{\rm LO}^{\rm R}$$
. (5b)

In particular for optimum-doped Bi<sub>2</sub>Sr<sub>2</sub>CaCu<sub>3</sub>0<sub>8</sub> (Bi-2212) this ARPES energy shift  $2\Delta_{SC}$  in Fig. 1(a) is 74 meV (1 meV=  $10^{-3}$  eV), where combining with resonancesoftened La<sub>2-x</sub>Sr<sub>x</sub>CuO<sub>4</sub> (La-214) Cu-O mode data  $E_{LO}^R \approx 70$  meV from Table I. Whereas for YBa<sub>2</sub>Cu<sub>3</sub>O<sub>7</sub> (Y-123) the ARPES gap Eq. 5b is predicted by this Y-123 data alone, with the shift  $2\Delta_{HTC}$  and  $E_{LO}^R$  from Table I and Fig. 1(b) respectively 58 meV and 56 meV [20]. Notably Eq. 5b is further given by a directional resonance coupling of these TF dynamics to the ARPES distribution [1].

Moreover Eq. 5 in addition to a Nambu-sigma mass analog constitutes an empirical "slowing" relation, where the simple energy and coupling independence reflects nondispersive, nonlinear inelastic-Frohlich slowing, and the conservation or irreversibility derives from the system asymmetry [1]. Such energetic "bunching" from these nonlocal dynamics, in setting the (TF) polaron apart from that due to Feynman [10], is consistent with the short-range stable normal-state pairing. Similarly, such nonlinear extension of this Frohlich "action" [11b] facilitates its energetic sharing in the many-particle state.

Significantly an extra-high- $T_c$  multi-layer cubic BCS cuprate possibly observed as a minority phase is defined from this empirical TF Nambu-mass

analog gap relation Eq. 5b for the non-cubic cuprates, and in particular for the BCS-like/cubic alkali-fullerene  $Rb_2CsCo_{60}$  [21]. Hence in the latter where  $T_c$ =40 K the TF shift of 13 meV from Table I yields the expected near-BCS ratio,

$$2\Delta_{\text{HTC}}/T_{\text{c}}k_{\text{B}}$$
=(13 meV/40 K)(11.6 K/meV) $\approx$ 3.8, (6a)

paralleling a low-T<sub>c</sub> Nambu-analog gap mode in NbSe<sub>2</sub> with gap ratio 3.4 in Eq. A-1. Whereas for the (non-cubic) cuprates non-BCS gap ratios are specified, viz., for Bi-2212 with T<sub>c</sub>=95 K and  $2\Delta_{HTC}$ =74 meV experimentally from the TF-ARPES relation Eq. 5b, the ratio is

$$2\Delta_{\rm HTC}/T_{\rm c}k_{\rm B}$$
=74 meV/95 K)(11.6 K/meV)≈9.0 (6b)

(k<sub>B</sub>=1 meV/11.6 K). Similarly for Y-123 where T<sub>c</sub>=93 K and  $2\Delta_{HTC}$ =58 meV again given by Eq. 5b a decreased ratio is specified as

$$2\Delta_{\text{HTC}}/T_{c}k_{\text{B}}=58 \text{ meV}/93 \text{ K})(11.6 \text{ K/meV})\approx7.2,$$
 (6c)

where this decrease would reflect the relative uniformity of the Y-123 structure. Notably for the purposes of our projected ideal BCS cuprate, the diversity in the optimum high-T<sub>c</sub> gap of these structures of 13 meV, 74 meV and 58 meV is *fundamentally* specified by the TF shift Eq. 5. (Albeit the accompanying T<sub>c</sub> increases in the cuprates are offset by the large ratios reflecting non-BCS localization associated with the low lattice symmetry).

Added basis for this ideal BCS cuprate comes from such high-T<sub>c</sub> superconductivity associated with cubic symmetry *in general* as in optimum Rb<sub>2</sub>CsC<sub>60</sub> above. Hence in Ba<sub>0.6</sub>K<sub>0.4</sub>BiO<sub>3</sub> with  $2\Delta_{HTC}$ =70 cm<sup>-1</sup> and T<sub>c</sub>=30 K [22] (recall  $\hbar$  c=197.3 MeV-fm=0.01973 meV-cm) the gap ratio is

 $2\Delta_{\text{HTC}}/T_{c}k_{\text{B}}=(2\pi)(70 \text{ cm}^{-1}/30 \text{ K})(11.6 \text{ K/meV})(0.01973 \text{ meV-cm})\approx 3.4,$  (7a)

and in Rb<sub>3</sub>C<sub>60</sub> with  $\Delta$ <sub>SC</sub>/k<sub>B</sub>=53 K, or  $\Delta$ =4.6 meV and T<sub>c</sub>=29.4 K [23] the ratio is

$$2\Delta_{\text{HTC}}/T_{c}k_{\text{B}}=2(4.6 \text{ meV}/29.4 \text{ K})(11.6 \text{ K/meV})\approx 3.6.$$
 (7b)

These Nambu-analog TF gap and BCS ratio specifications from Eqs. 5b, 6 and 7 provide an empirical basis for defining the ideal BCS cuprate, e.g., the gap specifications imply the TF shift is *fundamental* and therefore also holds for the record-high-T<sub>c</sub> 164 K multi-layer structure HgBa<sub>2</sub>Ca<sub>2</sub>Cu<sub>3</sub>O<sub>8</sub> under applied hydrostatic pressure [24]. Hence taking the resonance LO-mode energy  $E_{LO}^{R}$  in Eq. 5b in this case as ~65 meV, as estimated from an average of the Y-123 and La-214 data, the gap ratio is logically reduced from 7-9 in Bi-2212 and Y-123 to

$$2\Delta_{HTC}/T_ck_B \sim (65 \text{ meV}/164 \text{ K})(11.6 \text{ K/meV})=4.6,$$
 (8)  
where this reduction is consistent with symmetry increases toward a cubic BCS  
structure from the added planes and hydrostatic pressure.

In Ref. 1 a ~250 K infinite-layer cubic cuprate superconductor was projected from this predictive TF gap Eq. 5b plus BCS condition generally observed in cubic high-T<sub>c</sub> structures (cf. Eqs. 6a and 7). For such a Bi-2212 base structure this T<sub>c</sub> limit from Eq. 6b as a result of added planes and applied pressure is thus

$$T_c = 74 \text{ meV}(11.6 \text{ K/meV})/3.5=245 \text{ K}.$$
 (9a)

Stated in reverse, the BCS ratio limit is reached for the ideal 245 K  $T_c$  structure as

$$2\Delta_{\text{HTC}}/T_{c}k_{\text{B}} = (74 \text{ meV}/245 \text{ K})(11.6 \text{ K/meV})=3.5.$$
 (9b)

Notably this  $T_c$  is consistent with numerous observations of possible minorityphase superconductivity near this temperature. Summarizing this Nambu-analog gap ratio progression toward the extrahigh-T<sub>c</sub> structure, starting with the low-T<sub>c</sub> analog NbSe<sub>2</sub> with a ratio 3.4 in Eq. A-1, the sequence continues in the optimum high-T<sub>c</sub> structures with ratios 3.8, 9.0 and 7.2, all given by the TF gap Eq. 5. Continuing to higher T<sub>c</sub> a TF-lattice symmetry gap ratio pattern peaking at 9.0 emerges with increasing T<sub>c</sub> as 3.8, 9.0, 7.2, 4.6(?)  $\rightarrow$  3.5(?), with (?) denoting the TF-BCS-like projections.

In conclusion, the latest calculated pion-quark pairing temperature is predicted by an intuitive BCS relation identified from a quark-level Goldberger-Treiman relation with a Nambu scalar-mass gap parameter taken in the lowmass limit. Whereas in the high-mass limit this parameter predicts the sigma mass, and moreover has an analog in high-T<sub>c</sub> superconductivity in the stable localized energy gap as specified by the Thornber-Feynman (TF) polaron with a strongly-constrained shift from conserved dynamics paralleling the high-T<sub>c</sub> case. Notably an extra-high-T<sub>c</sub> BCS cuprate possibly observed as a minority phase is defined by a high-T<sub>c</sub> gap ratio sequence based on these empirical Nambu-analog TF gaps.

#### Appendix: Particle physics analog of a low-T<sub>c</sub> gap mode

We outline here how a massive "gap mode" appearing at  $T_c$  near  $2\Delta$  in the BCS superconductor NbSe<sub>2</sub> constitutes a particle physics analog leading in the low-mass limit to the pion pairing BCS relation Eq. 2. A near-BCS gap ratio is defined in NbSe<sub>2</sub> from  $2\Delta$ =17 cm<sup>-1</sup> (with  $\omega$ = $2\pi v$ ) and  $T_c$ =7.2 K [25],

 $2\Delta/T_ck_B = [(2\pi)17 \text{ cm}^{-1}/7.2 \text{ K}][0.01973 \text{ meV-cm}] [11.6 \text{ K/meV}] \approx 3.4$ , (A-1)

suggesting a nominal global pairing over the Fermi surface. The particle physics connection of this gap mode follows from its identification by Littlewood and Varma with the Nambu scalar mass [26]. This mode arises from a coupling of the superconductivity/gap to an amplitude mode of the charge-density-wave (CDW) (created from optical-phonon induced oscillation of the CDW gap at a local zone face). Hence as a local paired bound state near 2∆ with mass M, or

it has an analog in the Nambu scalar mass [27] identified in Eq. 2, i.e.,

$$2m_{q} = m_{s}.$$
 (A-3)

This "gap parameter" Eq. A-3 in the high-mass limit predicts the sigma meson mass  $m_{\sigma}$ , as also concluded in Ref. 28 from a chiral-breaking analog mode of Nambu and Jona-Lasinio [27], where  $m_q$  is the constituent quark mass as verified experimentally by the scalar sigma with mass  $m_{\sigma}\approx600$  MeV [29]. Independent determination of the quark mass  $m_q$  of 325 MeV in the chiral limit from the linear  $\sigma$  model [6] (or simply  $m_N/3\approx313$  MeV from the 3 quarks in the nucleus) yields 626-650 MeV for  $m_{\sigma}$ . Equation A-3 expresses the dispersion-theoretic limit in the dispersion range with squared four-momentum  $q^2=4m^2$  to 0, where the quark-anti-quark pair can be considered as "touching" with  $q^2=(2m_q)^2$ .

A BCS-particle physics analog of Eq. A-2 in the opposite zero-mass limit in the text gives the predictive BCS-like relation for the pion with the  $\bar{q}q$  pair considered as "fused", i.e., a tightly-bound pair at  $q^2=m_{\pi}^2=0$ , or at the VMD value  $r_{\pi}\approx 0.63$  fm.

#### References

1. B. A. Green and M. D. Scadron, Physica B **305** 175 (2001).

2. Quoting from S. Weinberg in Phys. Rev. Lett. **65** 1181 (1990), "An explanation is offered why quarks in the constituent quark model should be particles with axial coupling  $g_A=1$  and no anomalous magnetic moment".

Y. Nambu, Phys. Rev. Lett. **4** 380 (1960); J. Goldstone, Nuovo Cim. **19** 154 (1961).

4. D. Bailin, J. Cleymans and M.D. Scadron, Phys. Rev. D 31 164 (1985); J.

Cleymans, A. Kocic and M.D. Scadron, Phys. Rev. D 39 323 (1989); N. Bilic, J.

Claymans and M.D. Scadron, Int. J. Mod. Physics A 10 1169 (1995); M. D.

Scadron and P. Zenczykowski, Hadronic Jour. 26 503 (2003).

5. F. Karsch, Quark Matter Conf. 2001, Stonybrook, Jan. 2001, hep-ph/0103314;F. Karsch et al., Nucl. Phys. B 605 579 (2001).

6. R. Delbourgo and M. D. Scadron, Mod. Phys. Lett. A 10 251 (1995); hepph/9910242; see also V. Elias and M. D. Scadron, Phys. Rev. Lett. **53** 1129 (1984).

This is the chiral limit, whereas the experimentally measured radius is
0.672±0.008 fm; see. e.g., Particle Data Group, Phys. Lett. B 592 499 (2004).

 J. J. Sakurai, Ann. Phys. **11** 1 (1960); ibid., *Advanced Quantum Mechanics*, Addison-Wesley (1967).

9. K.K. Thornber and R.P. Feynman, Phys. Rev. B 1 4099 (1970).

10. This physics due the system asymmetry sets the TF quasiparticle apart from the commonly used dispersive negative-energy Feynman polaron [11].

- 11. R.P. Feynman, Phys. Rev. 97 660 (1955); also R.P. Feynman and A. R.
- Hibbs, Quantum Mechanics and Path Integrals, McGraw-Hill, N.Y. (1965), Chap.
- 12. Ref. 1 and references to polar mobility data therein.
- 13. M. R. Norman, et al., Phys. Rev. Lett. **79** 3506 (1997).
- 14. D. H. Lu, et al., Phys. Rev. Lett. 86 4370 (2001).
- 15. L. Pintschovius et al., Physica C 185-189 156 (1991); W. Reichardt et al., op.
- cit. **162-164** 464 (1989).
- 16. R. J. McQueeney et al., Phys. Rev. Lett. 82 628 (1999).
- 17. J. –H. Chung et al., Phys. Rev. B 67 014517 (2003).
- 18. O. Gunnarson, Rev. Mod. Phys. 69 575 (1997), Sect. III.B.
- 19. Op. cit., Sect. II.A.
- 20. See Ref. 1, Note Added in the Proof.
- 21. B. A. Green, Phys. Lett. A 227 372 (1997).
- 22. C. -K. Loong et al., Phys. Rev. Lett. 62 2628 (1989).
- 23. R. F. Kiefl et al., Phys. Rev. Lett. 70 3987 (1993).
- 24. L. Gao et al., Phys. Rev. B 50 4260 (1994).
- 25. R. Sooryakumar and M.V. Klein, Phys. Rev. Lett. 45 660 )(1980); Phys. Rev.
- B **23** 3213 (1981); **23** 3223 (1981).

26. P.B. Littlewood and C.M. Varma, Phys. Rev. B 26 4883 (1982); ibid., Phys.

Rev. Lett. 47 811 (1981).

27. Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** 345 (1961).

28. R. Delbourgo and M.D. Scadron, Phys. Rev. Lett. 48 379 (1982); M.D.

Scadron, Ann. Phys. 148 257 (1983).

29. Particle Data Group, S. Eidelman et al., Phys. Lett. B 592 1 (2004).

	Optimum gap, 2∆ (meV)	In-plane LO-phonon energy (meV) E <sub>LO</sub> E <sub>LO</sub> R		Critical temperature, T <sub>c</sub> (K)
Bi <sub>2</sub> Sr <sub>2</sub> CaCu <sub>2</sub> O <sub>8</sub> (Bi-2212)	74 [12]	80 [15]	<u> </u>	95
La <sub>1.85</sub> Sr <sub>0.15</sub> CuO <sub>4</sub> (La-214)	15	83 [16]	70 [16]	40
YBa <sub>2</sub> Cu <sub>3</sub> O <sub>7</sub> (Y-123)	58 <sup>a</sup> [14]	68 <sup>a</sup> [17]	56 <sup>a</sup> [17]	93
Rb <sub>2</sub> CsC <sub>60</sub>	12 <sup>b</sup>	13 [18]		40 [19]

<sup>a</sup>a-axis data in direction of reduced screening across charge stripes. <sup>b</sup>This gap with the observed  $T_c$  of 40 K corresponds to the BCS gap ratio generally observed in these structures.

**Table I**. Optimum gap  $2\Delta$  and non-resonance and resonance in-plane LO-

phonon energies  $E_{LO}$  and  $E_{LO}^{R}$  in the (1,0,0) direction in Bi-2212, La-214 and Y-

123 from ARPES and inelastic neutron scattering data, respectively. Note the

 $2\Delta$ -E<sub>LO</sub><sup>R</sup> resonance match according to the TF polar coupling relation Eq. 5b from

such resonance "softening" of the mid-zone Cu-O mode, and similarly in

Rb<sub>2</sub>CsC<sub>60</sub> according to Eq. 5a. The small gap in La-214 reflects the short-range

order.

## Figure 1

(a) ARPES distribution of Bi-2212 at ( $\pi$ ,0) momentum location on the Fermi surface that defines the optimum energy gap  $\Delta$  for the normal and superconducting states (from Ref. 13). (b) High-density a-axis superconducting peak defining the gap of Y-123 at ( $\pi$ ,0), with the corresponding Bi-2212 peak (from Ref. 14).

