

# Moments of the heavy-quark parton distribution function from QCD sum rules

A.G.Oganesian

Institute of Theoretical and Experimental Physics  
B.Chermushkinskaya 25, 117218, Moscow, Russia

## Abstract

The moments of the heavy quark-parton distribution functions in a heavy pseudoscalar meson are calculated from QCD sum rules. Expanding these sum rules in the inverse heavy quark mass we obtain the heavy-mass limits of the moments. Comparison with the finite mass results reveals that while the heavy mass expansion works reasonably well for the  $b$  quark, one has to take into account terms of higher than  $(1/m_c)^2$  order for the  $c$  quark. This result can provide a quantitative assessment of  $c$  and  $b$  quark fragmentation models based on the heavy-quark mass limit.

## 1

The knowledge of the heavy quark fragmentation functions is very important for description of various processes of heavy hadron production. There exist many different theoretical models and approaches, in which these functions are parameterized and their perturbative part calculated in QCD (see, for example [1]-[4]). Some of these models (e.g. [4]) use expansions, based on the heavy-quark mass limit or on HQET. That is why it seems interesting to reconsider the moments of the heavy-quark fragmentation functions, obtained long ago from QCD sum rules in [5]. In this approach the moments of the heavy-quark fragmentation functions into heavy-light meson (e.g., the  $c$ -quark fragmentation into a  $D$  meson)

$$M_n^D = \int_0^1 dz z^{n-1} D^c(z, Q^2) \quad (1)$$

and the moments of heavy-quark parton distribution functions in the same meson

$$M_n^c = \int_0^1 dx x^{n-1} c(x, Q^2) \quad (2)$$

were equated (in the scaling limit) following the familiar relation [7] ( see also [6] and [5] for details). The moments  $M_n^c$  were directly calculated from QCD sum rules. The advantage

of this method is the possibility to estimate the heavy parton distribution functions (hence, also fragmentation functions) in full QCD, in terms of universal parameters, such as quark masses and condensate densities. One can use this result to fix the parameters of different models, as was discussed in [5].

In addition, as will be shown here it is possible to check whether the expansion in the inverse heavy-quark mass is reliable for the description of the fragmentation functions, especially in the case of the  $c$ -quark. In what follows I address this question considering the fragmentation of heavy quarks to pseudoscalar heavy-light meson. A more complete study including mesons with different quantum numbers and assessing the heavy-quark spin symmetry violation in the fragmentation functions will be presented in the paper being in preparation.

Let me briefly remind the method and results of the paper [5]. For definiteness the  $c$ -quark fragmentation to  $D$  meson is considered, hence one needs the QCD sum rule for  $c$  quark-parton distribution in  $D$ -meson. One starts from the four-point correlator:

$$\Pi_{\mu\nu}(p_1, p_2; q_1, q_2) = i \int e^{ip_1x+iq_1y-ip_2z} d^4x d^4y d^4z \langle 0 | T \{ j_5(x) j_\mu^{em}(y) j_\nu^{em}(0) j_5^\dagger(z) \} | 0 \rangle \quad (3)$$

where the two  $c$ -quark electromagnetic currents,  $j_\mu^{em} = \bar{c}\Gamma_\mu c$ , are correlated with the two heavy-light pseudoscalar currents  $j_5 = \bar{c}\Gamma_5 u$  that interpolate  $D$  meson:

The above correlator is considered in the deep spacelike region at  $t = (p_1 - p_2)^2 = 0$ ,  $p_i^2, q_i^2 < 0$ , where all internal quarks are highly virtual and the operator-product expansion (OPE) is applicable. We take into account the contributions of the unit operator (bare loop), quark condensate  $\langle 0 | \bar{\psi}\psi | 0 \rangle$  and quark-gluon condensate  $\langle 0 | \bar{\psi}G_{\mu\nu}\sigma^{\mu\nu}\psi | 0 \rangle$ . The two latter contributions are important, their coefficients being proportional to the heavy quark mass. The gluon- and four-quark-condensate contributions are neglected, because the estimates show, that they are very small.

Importantly, the correlator is calculated as a function of two independent external momenta squared  $p_1^2, p_2^2$ . In this way one can reliably extract the ground state  $D$ -meson contribution both in the initial and final states, making the resulting sum rules more accurate. Hereafter we will consider the invariant amplitude multiplying the kinematical structure  $g_{\mu\nu}$ :  $\Pi_{\mu\nu} = g_{\mu\nu}\Pi + \dots$ . Let us express this amplitude in terms of double dispersion relation over  $p_1^2$  and  $p_2^2$ , i.e:

$$\Pi = \frac{1}{\pi^2} \int \frac{du_1}{u_1 - p_1^2} \int \frac{du_2}{u_2 - p_2^2} \rho(u_1, u_2, s) \quad (4)$$

We will consider the derivatives of the amplitude  $\Pi$  over  $s = (p_1 + q_1)^2$ . Furthermore, using the dispersion representation over  $s$ , one can write the spectral density  $\rho$  in the form

$$\frac{1}{n!} \frac{d^n}{ds^n} \rho(u_1, u_2, s) = \int \frac{ds_1}{(s_1 - s)^{n+1}} f^0(u_1, u_2, s_1). \quad (5)$$

As usual in the derivation of a QCD sum rule, one should saturate the amplitude (5) by hadronic states and equate it to the result of OPE (the bare loop and operators of dimensions 3 and 5, as mentioned above). One easily finds the contribution of the ground ( $D$ -meson) states in the initial and final hadron channels to the equation (5):

$$\rho = Im_{u_1} Im_{u_2} \Pi = -\pi^2 g_D^2 m_D^4 \delta(u_1 - m_D^2) \delta(u_2 - m_D^2) T_1 \quad (6)$$

where  $g_D m_D^2 = i\langle 0 | \bar{c} \gamma_5 c | c \rangle$  and  $T_1$  is defined by following relation

$$\int e^{iq_1 x} d^4 x \langle D | T \{ j_\mu^{em}(x) j_\nu^{em}(0) \} | D \rangle = g_{\mu\nu} T_1 + \dots \quad (7)$$

The above amplitude corresponds to the amplitude of the virtual photon scattering on  $D$  meson and is therefore obviously related with the  $c$ -quark parton distribution (structure function of  $D$  meson) by well known relation

$$Im_s T_1 = -\pi c(x). \quad (8)$$

Now one should substitute eq. (6-8) into eq. 5, then perform two independent Borel transformation on  $p_1^2$  and  $p_2^2$ , with the Borel masses  $M_{1Bor}^2$  and  $M_{2Bor}^2$ , respectively. We will equate the Borel masses  $M_{1Bor}^2 = M_{2Bor}^2 = M^2$  only in the final sum rule. Finally, the hadronic part of the sum rule can now be written as

$$\begin{aligned} \frac{1}{n!} \frac{d^n}{ds^n} \Pi &= -\pi^2 g_D^2 m_D^4 \exp[-2m_D^2/M_{Bor}^2] \frac{1}{\pi} \int_{m_c^2} \frac{ds_1}{(s_1 - s)^{n+1}} Im_s T_1(Q^2, s_1) \\ &= -\pi^2 g_D^2 m_D^4 \exp[-m_D^2/M_{Bor}^2] \frac{1}{\pi} \int_0^1 \frac{dx}{x^2} C(x, Q^2) (m_D^2 - Q^2 + Q^2/x - s)^{-n-1} = \end{aligned} \quad (9)$$

Here we adopt quark-hadron duality approximation for the hadron continuum contribution, so that it cancels with the appropriate part of the bare loop contribution. Following [5], we choose the value of the variable  $s$  in the unphysical region as  $s = m_D^2 - Q^2$ , and transform the integrals in the first line of eq.(9) into the moments of the parton distribution. of the  $c$  quark in  $D$  meson. Finally, equating the hadronic representation (7) to the OPE result, one can write down the following sum rules for these moments:

$$\begin{aligned} M_n &= g_D^{-2} m_D^{-4} \left( \frac{1}{\pi^2} \int_{m^2}^{s_0} du_1 \exp \left[ \frac{-2(u_1 - m_D^2)}{M^2} \right] \int_{z_-}^{z_+} dz f_0 * (1 + z - m_D^2/Q^2)^{-n-1} \right. \\ &\quad \left. + \frac{ma}{4\pi^2} L \eta^{-n-1} \exp \left[ \frac{-2(m^2 - m_D^2)}{M^2} \right] + R_6 \right) \end{aligned} \quad (10)$$

where

$$f_0 = \frac{3}{8} (s_1 - m^2) \left[ \left( 1 + z + \frac{s_1 + 2m^2}{Q^2} \right) \left( (1 + z - s_1/Q^2)^2 + \frac{4s_1}{Q^2} \right)^{-1/2} - \frac{4m^2}{Q^2} \left( 1 + \frac{4m^2}{Q^2} \right)^{-1/2} \right] \quad (11)$$

and

$$\begin{aligned} R_6 &= \frac{mm_0^2 a}{4\pi^2} \frac{1}{2M^2} \left[ 4/M^2 - 2m^2/M^4 + 1/Q^2 + \frac{(n+1)}{\eta} \left( \frac{2}{3Q^2} - 1/M^2 - \frac{2m^2}{M^2 Q^2} \right) \right. \\ &\quad \left. - \frac{(n+1)(n+2)m^2}{Q^4 \eta^2} \right] \eta^{-n-1} \exp \left[ \frac{-2(m^2 - m_D^2)}{M^2} \right] \end{aligned} \quad (12)$$

where

$$z_+ = u_1(1/Q^2 + 1/(2m^2)) - 1/2 + (1 + 4m^2/Q^2)^{1/2}(u_1/m^2 - 1)/2$$

,

$$z_- = u_1(1/Q^2 + 1/(2m^2)) - 1/2 - (1 + 4m^2/Q^2)^{1/2}(u_1/m^2 - 1)/2$$

.

Here  $a = -(2\pi)^4 \langle 0 | \bar{\psi}\psi | 0 \rangle$ ,  $\eta = (1 - (m^2 - m_D^2)/Q^2)$ . The factor  $L = \ln(m/\lambda)/\ln(\mu\lambda)^{4/9}$  accounts for the anomalous dimension of the quark condensate, normalized at point  $\mu$ . We adopt the values of condensates [8], [9], normalizing them at  $\mu = 1\text{GeV}$ . In this case the factor  $L$  is close to the unity.

In [5], it was shown that the first few moments of the sum rules are well behaved and the following numerical results were obtained:

$$M_2 = 0.85, M_3 = 0.75, M_4 = 0.67$$

at  $Q_0^2 = 20\text{GeV}^2$

$$M_2 = 0.9, M_3 = 0.83, M_4 = 0.78$$

at  $Q_0^2 = 10\text{GeV}^2$ .

These moments can be use as an input in the evolution equations, at some initial point  $Q_0^2 = (3 - 5)m_D^2$ , hence it is possible to predict the structure functions also at large  $Q^2$ .

In what follows we will use the sum rules on order to address only one question, if it is possible to use the heavy quark limit for the heavy quark parton distribution (and, correspondingly, fragmentation function). For that, we expand the sum rules for the moments in the inverse mass of the heavy quark.

Substituting in eq(9) instead of the  $g_D^2 m_D^4$  the sum rule obtained from the two-point correlator [10]

$$g_D^2 m_D^4 = \exp(m_D^2/M_B^2) \left( \frac{3}{8\pi^2} \int_{m^2} du (u - m^2) \exp(-u/M_B^2)/s \right. \\ \left. + \frac{ma}{4\pi^2} \left[ L + \frac{m_0^2}{2M_B^2} (2 - m^2/M^2) \exp(-m^2/M_B^2) \right] \right) \quad (13)$$

One can then rewrite the sum rules for  $M_n$  as

$$M_n = \frac{R_0(n) + R_4(n) + R_6(n)}{K_0 + K_4 + K_6} \quad (14)$$

where the terms  $R_d(n)$  in the numerator and  $K_d$  in the denominator originate from the dimension  $d = 0, 4, 6$  contributions to the OPE of the four-point and two-point correlator, respectively. Following [5] we adopt the relation  $M_B^2 = M^2/2$  between the Borel masses in the four-point and two-point sum rules. Note that the exponential factor containing  $m_D^2$  cancels in the ratio of the two sum rules.

To investigate the heavy-quark mass limit  $m \rightarrow \infty$  limit of (14) we employ the standard scaling relations for the heavy hadron mass, continuum threshold and Borel parameter:

$$m_D = m + \bar{\Lambda}, \quad s_0 = m^2 + 2m\omega, \quad M_B^2 = 2\tau m \quad (15)$$

and in addition assuming the scaling  $Q^2 = \gamma m_D^2$ , where in the heavy quark limit the parameters  $\bar{\Lambda}, \omega, \tau$  and  $\gamma$  do not depend of the heavy quark mass.

The terms entering the numerator of eq.(15) have the following expressions:

$$R_0(n) = \int_0^{z_m} \frac{z dz}{2\tau} e^{-z} I_0(n) \quad (16)$$

where

$$I_0(n) = \int_{t_-}^{t_+} \left[ \frac{1 + 4(\rho/\gamma) + t/m + 4(\varphi/m)(\rho/\gamma)}{\sqrt{(1 + t/m)^2 + 4(\rho/\gamma) + 8(\varphi/m)(\rho/\gamma)}} - \frac{4(\rho/\gamma)}{v} \right] \frac{dt}{\lambda^{n+1}} \quad (17)$$

,  $\varphi = z\tau$ ,  $\delta = \bar{\Lambda}/m$ ,  $z_m = \omega/\tau$ , and

$$\rho = m^2/m_D^2 = 1/(1 + 2\delta + \delta^2)$$

$$\lambda = 1 + t/m + (\rho - 1)/\gamma + 2(\varphi\rho)/m$$

$$v = \sqrt{1 + 4\rho/\gamma}$$

$$t_+ = z\tau(1 + v)$$

$$t_- = z\tau(1 - v)$$

$$y = \bar{\Lambda}/\tau$$

Expanding in the powers of  $1/m$ , we obtain the first three terms for  $R_0$ :

$$R_0(n) \simeq D_2 + (n+1)\delta \frac{2D_2 - (2+\gamma)D_3}{\gamma} + \delta^2(n+1)(n+2) \left[ \frac{2(n+2) + 3\gamma}{\gamma^2(n+2)} D_2 - \frac{2}{\gamma} \left( 1 + \frac{2(n+1)}{(n+2)\gamma} \right) D_3 + \frac{2D_4}{3\gamma^2} (\gamma^2 + 4\gamma + 3 + \frac{2\gamma}{(n+1)(n+2)}) \right] \quad (18)$$

Here  $D_n$  is defined as

$$D_n = n! E_n(\omega/\tau)/y^{n-2}$$

and

$$E_n(z) = 1 - e^{-z} - ze^{-z} - z^2 e^{-z}/(2!) - \dots - z^n e^{-z}/(n!)$$

For the contribution of  $d = 4$  operators we obtain:

$$R_4(n) = \frac{a}{12\tau^3} L \frac{1}{\kappa^{n+1}}, \quad (19)$$

where  $\kappa = 1 + (\rho - 1)/\gamma$ . After expansion the first three terms are:

$$R_4(n) \simeq \frac{a}{12\tau^3} L \left( 1 + 2\delta(n+1)/\gamma + \delta^2 \frac{(n+1)(n+2)}{\gamma} [2/\gamma - 3/(n+2)] \right) \quad (20)$$

Finally, the contribution of  $d = 6$  operators transforms into

$$R_6(n) = \frac{a}{12\tau^3} \frac{m_0^2}{16\tau^2} \left[ -1 + 8\epsilon - 2\epsilon \frac{(n+1)}{\kappa} (1 + 2\rho/\gamma) + 8\epsilon^2 \left( \frac{\rho}{\gamma} + \frac{2(n+1)}{3\kappa} \frac{\rho}{\gamma} - \frac{(n+1)(n+2)}{2\kappa^2} \frac{\rho^2}{\gamma^2} \right) \right] \frac{1}{\kappa^{n+1}}, \quad (21)$$

where

$$\epsilon = \tau/m = \delta/y$$

. After expansion the above expression retaining again the first three terms yields:

$$R_6(n) \simeq \frac{a}{12\tau^3} \frac{m_0^2}{16\tau^2} \left[ -1 + \frac{2\delta}{y} \left( 4 - (n+1)[1 + (y+2)/\gamma] \right) - \frac{2\delta^2(n+1)}{y\gamma} \left( (n+2)[2 + (y+4)/\gamma + 2/(y\gamma)] - 3y - 24 - (8/3)(2n+5)/y \right) \right] \quad (22)$$

And, finally, for the denominator in (14) we obtain

$$K = \int_0^{z_m} \frac{z^2 dz}{1 + 2\varphi/m} e^{-z} + \frac{a}{12\tau^3} L + \frac{a}{12\tau^3} \frac{m_0^2}{16\tau^2} (-1 + 4\delta/y) \quad (23)$$

$$K^* = \int_0^{z_m} \frac{z^2 dz}{(1 + 2\varphi/m)^2} \left( 1 + \frac{4\varphi}{3m} \right) e^{-z} + \frac{a}{12\tau^3} L + \frac{a}{12\tau^3} \frac{m_0^2}{16\tau^2} (-1) \quad (24)$$

resulting in the heavy quark limit in

$$K \simeq D_2 - 2D_3\delta - 4D_4\delta^2 + \frac{a}{12\tau^3} \left( L - \frac{m_0^2}{16\tau^2} \right) + \frac{a}{12\tau^3} \frac{m_0^2}{16\tau^2} \frac{4\delta}{y} \quad (25)$$

$$K^* \simeq D_2 - \frac{8}{3}D_3\delta - \frac{20}{3}D_4\delta^2 + \frac{a}{12\tau^3} \left( L - \frac{m_0^2}{16\tau^2} \right) \quad (26)$$

One can easily see, that in the  $m \rightarrow \infty$  limit  $M_n = 1$  for all  $n$ . This limiting case corresponds to  $c(x) = \delta(1 - x)$ , and is accordance with a simple physical picture of the infinitely heavy quark inside heavy hadron carrying the whole momentum. Interestingly, the sum rule (12) predict the deviation from unity,  $\Delta = 1 - M_n$  related with the finiteness of the heavy quark mass for both parton distributions and fragmentation function. From

the equations presented before it is easy and instructive to calculate the actual value of  $\Delta$  in the first and second orders of expansion and compare it with the exact value of  $\Delta$  to examine the accuracy of the expansion in the inverse heavy quark mass. For numerical analysis we choose  $\bar{\Lambda} = 0.6$  GeV,  $\omega = 1.4$  GeV and  $\tau = 0.6$  GeV. One can see, that these values reproduce the  $D$  meson mass (assuming  $m_c \simeq 1.3$  GeV), as well as the continuum threshold  $s_0 = 6\text{GeV}^2$ , and Borel mass  $M_B^2 = 1 - 2\text{GeV}^2$  in the ballpark of input parameters used in the sum rules in  $D$  channel (see e.g., [10]). Before we discuss the numerical results, it should be noted, that the results for moments, obtained in [5], which we have cited before, was obtained in slightly another way of numerical analysis:- the ratio of the neighboring moments was considered. As was discuss in [5], this two methods give very close results (the difference less than 5-10%, which is within the accuracy of our sum rules approach).

The numerical results for heavy-quark mass expansion of the first and second moments are shown in Fig. 1 and Fig. 2, respectively, where they are plotted as a function of the dimensionless ratio  $k = m/m_c$ , for two different values of the ratio  $\gamma = Q^2/M_D^2 = 2, 4$ . One can easily see, that even at relatively large masses of order of the  $b$ -quark mass ( $k \sim 3$ ) the heavy mass limit and even the first-order expansion in inverse mass are not reliable. Including the second  $O(1/m^2)$  term one improves the situation for the  $b$  quark, but the approximate result for the moments still remains very far from the exact answer in the case of  $c$ -quark. We conclude that the heavy quark limit is not a reliable approximation for the parton distributions and fragmentation functions of  $c$ -quark.

The author thanks B.L. Ioffe and A. Khodjamirian for useful discussions. This work was supported in part by the Russian Foundation of Basic Research, project no. 06-02-16905a and the funds from EC to the project "Study of the Strong Interacting Matter" under contract N0. R113-CT-2004-506078.

## References

- [1] C. Peterson, D. Schlatter, I. Schmitt and P. M. Zerwas, Phys. Rev. D **27** (1983) 105.
- [2] M. Suzuki, Phys. Rev. D **33**, 676 (1986).
- [3] P. D. B. Collins and T. P. Spiller, J. Phys. G **11**, 1289 (1985).
- [4] E. Braaten, K. m. Cheung, S. Fleming and T. C. Yuan, Phys. Rev. D **51**, 4819 (1995).
- [5] G. L. Balaian, A. Y. Khodjamirian and A. G. Oganesian, Sov. J. Nucl. Phys. **49**, 697 (1989) [Yad. Fiz. **49**, 1125 (1989)].
- [6] V. G. Kartvelishvili, A. K. Likhoded and V. A. Petrov, Phys. Lett. B **78**, 615 (1978).
- [7] V. N. Gribov and L. N. Lipatov, Sov. J. Nucl. Phys. **15** (1972) 675 [Yad. Fiz. **15** (1972) 1218].
- [8] B. V. Geshkenbein, B. L. Ioffe and K. N. Zyablyuk, Phys. Rev. D **64** (2001) 093009.

- [9] B. L. Ioffe, Prog. Part. Nucl. Phys. **56**, 232 (2006).
- [10] T. M. Aliev and V. L. Eletsky, Sov. J. Nucl. Phys. **38**, 936 (1983) [Yad. Fiz. **38**, 1537 (1983)].




Figure 1: The first moment of the heavy quark distribution function in pseudoscalar heavy-light meson plotted as a function of the quark mass  $m$  in the units  $k = m/m_c$ . The lines denoted as "1ord", "2ord" and "total" correspond to the first-, second-order expansion in the inverse quark mass and to the exact answer, respectively. The value of  $\gamma = Q^2/M_D^2 = 2(4)$  is chosen in the left (right) panel.




Figure 2: The same as in Fig. 1 for the second moment