

Frascati Physics Series Vol. XLVI (2007), pp. 000-000
 HADRON07: XII INT. CONF. ON HADRON SPECTROSCOPY – Frascati, October 8-13, 2007
 Parallel Session

Self-consistent coupled-channel approach to D and \bar{D} in hot dense matter

Laura Tolós

FIAS. J.W.Goethe-Universität.

Ruth-Moufang-Str. 1, 60438 Frankfurt (M), Germany

Angels Ramos

Dept. d'Estructura i Constituents de la Matèria. Universitat de Barcelona.

Diagonal 647, 08028 Barcelona, Spain

Tetsuro Mizutani

Department of Physics, Virginia Polytechnic Institute and State University.

Blacksburg, VA 24061, USA

Abstract

A self-consistent coupled-channel approach is used to study the properties of D and \bar{D} mesons in hot dense matter. The starting point is a broken SU(4) s -wave Tomozawa-Weinberg DN ($\bar{D}N$) interaction supplemented by an attractive isoscalar-scalar term. The Pauli blocking effects, baryon mean-field bindings, and π and open-charm meson self-energies are incorporated in dense matter at finite temperature. In the DN sector, the dynamically generated $\tilde{\Lambda}_c$ and $\tilde{\Sigma}_c$ resonances remain close to their free space position while acquiring a remarkable width because of the thermal smearing of Pauli blocking. Therefore, the D meson spectral density shows a single pronounced quasiparticle peak close to the free mass, that broadens with increasing density, and a low energy tail associated to smeared $\tilde{\Lambda}_c N^{-1}$, $\tilde{\Sigma}_c N^{-1}$ configurations. In the $\bar{D}N$ case, the low-density approximation to the repulsive \bar{D} self-energy is found unreliable already at subsaturation densities. From this study we speculate the possible existence of D -mesic nuclei. We also discuss the consequences for J/Ψ suppression at FAIR.

Over the last years there has been a growing interest in the open (D , \bar{D}) and hidden (e.g. J/Ψ) charmed mesons within the context of relativistic nucleus-nucleus collisions. In particular, J/Ψ suppression was predicted as a clear signature of the formation of the quark-gluon plasma (QGP) ¹⁾.

The future CBM (Compressed Baryon Matter) experiment of the FAIR (Facility for Antiproton and Ion Research) project at GSI will investigate, among others, the possible modifications of the properties of open and hidden charmed mesons in a hot and dense baryonic environment. The in-medium modification of the $D(\bar{D})$ mesons may explain the J/Ψ suppression in an hadronic environment, on the basis of a mass reduction of $D(\bar{D})$ in the nuclear medium ^{2, 3)}. Furthermore, this reduction in mass was thought to provoke possible D^0, D^-, \bar{D}^0 bound states in heavy nuclei such as Pb ²⁾.

However, a self-consistent coupled-channel meson-baryon approach in nuclear medium is found essential due to the strong coupling among the DN and other meson-baryon channels ^{4, 5, 6, 7)}, which induces the appearance of dynamically-generated resonances close to threshold.

In the present article, we pursue a coupled-channel study of the spectral properties of D and \bar{D} mesons in nuclear matter at finite temperatures by extending the result of Ref. ⁷⁾ to finite temperature. Then our finding is used to discuss the possibility of D -mesic nuclei as well as to examine the possible implications in the J/Ψ suppression at FAIR.

Self-consistent coupled-channel approach for D and \bar{D} mesons

The D and \bar{D} self-energies at finite temperature are obtained from a self-consistent coupled-channel calculation taking, as bare interaction (V), a type of broken $SU(4)$ s -wave Tomozawa-Weinberg (TW) interaction supplemented by an attractive isoscalar-scalar term (Σ_{DN}). The multi-channel transition matrix T

$$T = V + V G T \quad (1)$$

is solved using a cutoff regularization ⁷⁾, which is fixed by reproducing the position and the width of the $I = 0$ $\Lambda_c(2593)$ resonance. As a result, a new Σ_c resonance in the $I = 1$ channel is generated around 2800 MeV.

The in-medium solution at finite temperature is obtained by incorporating the corresponding medium modifications in the loop function G . We incorpo-

rate Pauli blocking effects, mean-field bindings of baryons via a temperature-dependent $\sigma - \omega$ model, and π and open-charm meson self-energies in the intermediate propagators (see Ref. 8).

The $D(\bar{D})$ self-energy is obtained self-consistently summing the in-medium $T_{D(\bar{D})N}$ amplitudes over the thermal nucleon Fermi distribution, $n(\vec{q}, T)$, as

$$\Pi_{D(\bar{D})}(q_0, \vec{q}, T) = \int \frac{d^3p}{(2\pi)^3} n(\vec{p}, T) [T_{D(\bar{D})N}^{(I=0)}(P_0, \vec{P}, T) + 3T_{D(\bar{D})N}^{(I=1)}(P_0, \vec{P}, T)] , \quad (2)$$

where $P_0 = q_0 + E_N(\vec{p}, T)$ and $\vec{P} = \vec{q} + \vec{p}$ are the total energy and momentum of the $D(\bar{D})N$ pair in the nuclear matter rest frame, and the values (q_0, \vec{q}) and (E_N, \vec{p}) stand for the energy and momentum of the $D(\bar{D})$ meson and nucleon, respectively, also in this frame. The in-medium spectral density then reads

$$S_{D(\bar{D})}(q_0, \vec{q}, T) = -\frac{1}{\pi} \frac{\text{Im} \Pi_{D(\bar{D})}(q_0, \vec{q}, T)}{q_0^2 - \vec{q}^2 - m_{D(\bar{D})}^2 - \Pi_{D(\bar{D})}(q_0, \vec{q}, T)} . \quad (3)$$

Open charm in hot dense matter

Fig. 1 shows the behavior of the in-medium $I = 0$ $\Lambda_c(2593)$ and $I = 1$ $\Sigma_c(2770)$ resonances, denoted as $\tilde{\Lambda}_c$ and $\tilde{\Sigma}_c$, respectively, for three different self-consistent calculations: i) the self-consistent dressing of D mesons only (dotted lines), ii) adding the mean-field binding of baryons (MFB) (dash-dotted lines) and iii) including MFB and the pion self-energy dressing (PD) (solid lines). We consider two models: the thick lines correspond to model A ($\Sigma_{DN} \neq 0$) while the thin-dashed lines refer to case (iii) within model B ($\Sigma_{DN} = 0$).

The inclusion of medium modifications at $T = 0$ lowers the position of the $\tilde{\Lambda}_c$ and $\tilde{\Sigma}_c$ resonances with respect to their free values, in particular with the inclusion of MFB. Their widths, which increase due to $\tilde{Y}_c N \rightarrow \pi N \Lambda_c, \pi N \Sigma_c$ processes, differ according to the phase space available. In contrast to the $\bar{K}N$ results [9, 10], the PD induces a small effect in the resonances because of reduced charm-exchange channel couplings. Still it is seen in the positions and widths through the absorption of these resonances by one and two nucleon processes when the pion self-energy is incorporated. On the other hand, models A and B show qualitatively similar features.

Finite temperature results in the reduction of the Pauli blocking effects due to the smearing of the Fermi surface. Both resonances move up in energy




Figure 1: $I = 0$ $\tilde{\Lambda}_c$ and $I = 1$ $\tilde{\Sigma}_c$ resonances in a hot medium.

closer to their free position while they are smoothen out, as seen previously in 5). At $T = 100$ MeV, $\tilde{\Lambda}_c$ is at 2579 MeV and $\tilde{\Sigma}_c$ at 2767 MeV for model A, while model B generates both resonances at higher energies: $\tilde{\Lambda}_c$ at 2602 MeV and $\tilde{\Sigma}_c$ at 2807 MeV.

We display in Fig. 2 the D meson spectral function at zero momentum obtained in cases (i) to (iii) for model A (thick lines) and in case (iii) for model B (thin line) at saturation density, $\rho_0 = 0.17 \text{ fm}^{-3}$. Two peaks appear in the spectral density at $T = 0$. The lower one corresponds to the $\tilde{\Lambda}_c N^{-1}$ excitations, whereas the higher one is the quasi(D)-particle peak mixed with the $\tilde{\Sigma}_c N^{-1}$ state. The lower energy mode goes up by about 50 MeV relative to (i) when MFB effects are included. The reason is that the meson requires to carry more energy to excite the $\tilde{\Lambda}_c$ in order to compensate for the attraction felt by the nucleon. The same effect is observed for the $\tilde{\Sigma}_c N^{-1}$ configuration that mixes with the quasiparticle peak. As expected, the PD does not alter much the position of $\tilde{\Lambda}_c N^{-1}$ excitation or the quasiparticle peak. For model B (case (iii) only), the absence of the Σ_{DN} term moves the $\tilde{\Lambda}_c N^{-1}$ excitation closer to the quasiparticle peak, while the latter completely mixes with the $\tilde{\Sigma}_c N^{-1}$ excitation.

Finite temperature effects result in the dilution of those structures with increasing temperature while the quasiparticle peak gets closer to its free value




Figure 2: The $q = 0$ D meson spectral function at ρ_0 for $T = 0, 100$ MeV

and it becomes narrower. This is due to the fact that the self-energy receives contributions from DN pairs at higher momentum where the interaction is weaker.

The evolution of the spectral function with temperature is seen in Fig. 3 for two densities, ρ_0 and $2\rho_0$, and two momenta, $q = 0$ MeV/c and $q = 450$ MeV/c, in case (iii) for model A. As in the previous figure, we observe how the $\tilde{\Lambda}_c N^{-1}$ and $\tilde{\Sigma}_c N^{-1}$ structures dissolve with increasing temperature, while the quasiparticle peak becomes narrower and moves closer to its free value position. The widening of the quasiparticle peak for larger nuclear density may be understood as due to enhancement of collision and absorption processes. The $\tilde{\Lambda}_c N^{-1}$ mode moves down in energy with increasing density due to the lowering in the position of the $\tilde{\Lambda}_c$ resonance induced by the more attractive Σ_{DN} term in model A.

With regard to the $\bar{D}N$ sector, we first study the effective $\bar{D}N$ interaction in free space and, in particular, the $\bar{D}N$ scattering lengths. For model A (B) those are $a^{I=0} = 0.61$ (0) fm and $a^{I=1} = -0.26$ (-0.29) fm. The zero value of the $I = 0$ scattering length for model B is due to the vanishing coupling coefficient of the corresponding pure TW $\bar{D}N$ interaction. This is in contrast to the repulsive $I = 0$ scattering length reported in ⁶⁾, while agreement is found in the $I = 1$ contribution. In the case of model A, the non-zero value of the $I = 0$ scattering length is due to the magnitude of the Σ_{DN} term. Our




Figure 3: The evolution of the D meson spectral function with temperature




Figure 4: \bar{D} mass shift and the low-density approximation

results are consistent with those of a recent calculation based on meson- and one-gluon exchanges [11]).

Next, we display the \bar{D} mass shift in cold nuclear matter in Fig. 4. For both A and B models, the mass shift is repulsive due to the $I = 1$ dominant component. However, despite the absence of resonances in the $\bar{D}N$ interaction, the low-density $T\rho$ approximation breaks down at relatively low densities, so it is not applicable at saturation density.

The comparison between D and \bar{D} optical potentials at $q = 0$ MeV/c as




Figure 5: The D and \bar{D} potentials as function of temperature

functions of temperature for two different densities (ρ_0 and $2\rho_0$) is shown in Fig. 5. For model A (B) at $T = 0$ and ρ_0 , the D meson feels an attractive potential of -12 (-18) MeV while the \bar{D} feels a repulsion of 11 (20) MeV. A similar shift for D meson mass is obtained in Ref. ⁵⁾. The temperature dependence of the repulsive real part of the \bar{D} optical potential is very weak, while the imaginary part increases steadily due to the increase in the collisional width. The picture is somewhat different for the D meson due to the overlap of the quasiparticle peak with the $\tilde{\Sigma}_c N^{-1}$ mode. Furthermore, the in-medium behavior of the $\tilde{\Sigma}_c N^{-1}$ mode is determinant for understanding the effect of the Σ_{DN} term on the D meson potential.

Taking into account our results, we might look at the question of possible \bar{D} bound states discussed in ²⁾. While D^- -mesic atoms can always be bound by the Coulomb interaction, no strongly bound nuclear states or even bound \bar{D}^0 nuclear systems are expected due to the repulsive \bar{D} -nucleus optical potential at zero momentum. In the charm $C = 1$ sector, an experimental observation of bound D nuclear states is ruled out by the moderate attraction and large width found for the D meson optical potential.

With respect to the J/Ψ suppression, the in-medium \bar{D} mass is seen to increase by about $10 - 20$ MeV whereas the tail of the quasiparticle peak of the D spectral function extends to lower "mass" due to the thermally spread

$\tilde{Y}_c N^{-1}$. Nevertheless, it is very unlikely that this lower tail extends as far down by 600 MeV with sufficient strength to influence the J/Ψ production. So the only way for the J/Ψ suppression to take place is by cutting its supply from the excited charmonia: $\chi_{c\ell}(1P)$ or Ψ' , which will be strongly absorbed in the medium by multi-nucleon processes.

1 Acknowledgments

This work is partly supported by contracts MRTN-CT-2006-035482 (FLAVIANet, EU), RII3-CT-2004-506078 (HadronPhysics, EU), by FIS2005-03142 (MEC, Spain), 2005SGR-00343 (Generalitat de Catalunya), ANBest-P and BNBest-BMBF 98/NKBF98 (Germany).

References

1. T. Matsui and H. Satz, *Phys. Lett.* **B178**, 416 (1986).
2. K. Tsushima, D. H. Lu, A. W. Thomas, K. Saito and R. H. Landau, *Phys. Rev.* **C59**, 2824 (1999)
3. A. Sibirtsev, K. Tsushima and A. W. Thomas, *Eur. Phys. J.* **A6**, 351 (1999); A. Hayashigaki, *Phys. Lett.* **B487**, 96 (2000); A. Mishra, E. L. Bratkovskaya, J. Schaffner-Bielich, S. Schramm and H. Stöcker, *Phys. Rev.* **C70**, 044904 (2004).
4. L. Tolos, J. Schaffner-Bielich and A. Mishra, *Phys. Rev.* **C70**, 025203 (2004).
5. L. Tolos, J. Schaffner-Bielich and H. Stöcker, *Phys. Lett.* **B635**, 85 (2006).
6. M. F. M. Lutz and C. L. Korpa, *Phys. Lett.* **B633**, 43 (2006).
7. T. Mizutani and A. Ramos, *Phys. Rev* **C74**, 065201 (2006).
8. L. Tolos, A. Ramos and T. Mizutani, *arXiv:nucl-th/0710.2684*.
9. A. Ramos and E. Oset, *Nucl. Phys.* **A671**, 481 (2000)
10. L. Tolos, A. Ramos and A. Polls, *Phys. Rev.* **C65**, 054907 (2002)
11. J. Haidenbauer, G. Krein, U. G. Meissner and A. Sibirtsev, *Eur. Phys. J.* **A33**, 107 (2007).