

# The charged Higgs boson mass in the 2HDM: decoupling and CP violation

Maria Krawczyk<sup>1,2 \*</sup> and Dorota Sokółowska<sup>1</sup>

1- Institute of Theoretical Physics, University of Warsaw  
00-681 Warsaw, ul. Hoża 69, Poland

2- TH-Division, CERN, CH-1211 Genève 23, Switzerland

Mass range of the charged Higgs boson in the 2HDM with explicit and spontaneous CP violation is discussed. Constraints on  $M_{H^\pm}$  in the CP conserving 2HDM(II) are shown.

## 1 The 2HDM potential and spontaneous symmetries breaking

The most general, invariant under gauge group  $SU(2)_L \times U(1)_Y$  and renormalizable potential of the Two Higgs Doublet Model (2HDM) [2, 3, 4] is given by

$$V = \frac{\lambda_1}{2} (\Phi_1^\dagger \Phi_1)^2 + \frac{\lambda_2}{2} (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) + \lambda_4 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) + \left[ \frac{1}{2} \lambda_5 (\Phi_1^\dagger \Phi_2) (\Phi_1^\dagger \Phi_2) + \lambda_6 (\Phi_1^\dagger \Phi_1) (\Phi_1^\dagger \Phi_2) + \lambda_7 (\Phi_2^\dagger \Phi_2) (\Phi_1^\dagger \Phi_2) + h.c. \right] - \frac{1}{2} m_{11}^2 (\Phi_1^\dagger \Phi_1) - \frac{1}{2} m_{22}^2 (\Phi_2^\dagger \Phi_2) - \left[ \frac{1}{2} m_{12}^2 (\Phi_1^\dagger \Phi_2) + h.c. \right], \quad (1)$$

where  $\lambda_{1-4}, m_{11}^2, m_{22}^2 \in \mathbb{R}$  (by the hermicity of the potential), while in general  $\lambda_{5-7}, m_{12}^2 \in \mathbb{C}$ . In the most general CP breaking form it has 14 parameters, however only 11 are independent, see e.g. [5, 6]. In the model there are five Higgs particles: three neutral  $h_1, h_2, h_3$  (for CP conservation - two CP-even  $h, H$  and one CP-odd  $A$ ) and two charged Higgs bosons  $H^\pm$ .

### 1.1 $Z_2$ and CP symmetries

The  $Z_2$  symmetry of the potential (1) is defined as the invariance of  $V$  under the following transformation of doublets:  $\Phi_1 \rightarrow -\Phi_1, \Phi_2 \rightarrow \Phi_2$  or  $\Phi_1 \rightarrow \Phi_1, \Phi_2 \rightarrow -\Phi_2$ . If  $Z_2$  (in either form) is a symmetry of the potential, then  $m_{12}^2 = \lambda_6 = \lambda_7 = 0$ . The  $Z_2$  symmetry is *softly* broken by the terms proportional to  $m_{12}^2$ .

General 2HDM allows for CP violation both explicitly and spontaneously [7, 8, 2]. The CP violation can be naturally suppressed by imposing a  $Z_2$  symmetry on the Higgs potential.

### 1.2 Reparametrization transformation

A global unitary transformation which mix two doublets and change their relative phase does not change the physical content of 2HDM as discussed recently in [9], see also [3, 4, 2]. It is given by

$$\begin{pmatrix} \Phi'_1 \\ \Phi'_2 \end{pmatrix} = \mathcal{F} \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix}, \quad \mathcal{F} = e^{-i\rho_0} \begin{pmatrix} \cos \theta e^{i\rho/2} & \sin \theta e^{i(\tau-\rho/2)} \\ -\sin \theta e^{-i(\tau-\rho/2)} & \cos \theta e^{-i\rho/2} \end{pmatrix}. \quad (2)$$

---

\*Supported in part by EU Marie Curie Research Training Network HEPTOOLS, under contract MRTN-CT-2006-035505 and by FLAVIANet contract No. MRTN-CT-2006-035482.

There are three *reparametrization* parameters -  $\rho, \theta, \tau$ , and in addition  $\rho_0$  parameter as an overall phase. If  $\theta = 0$  there is no mixing of two doublets and the transformation becomes a global transformation of doublets with an independent phase rotations (*rephasing*):

$$k = 1, 2 : \Phi_k \rightarrow e^{-i\rho_k} \Phi_k, \quad \rho_1 = \rho_0 - \frac{\rho}{2}, \quad \rho_2 = \rho_0 + \frac{\rho}{2}, \quad \rho = \rho_2 - \rho_1. \quad (3)$$

The original form of the potential is recovered by the appropriate changes of phases of the following coefficients:

### 1.3 Explicit and spontaneous CP violation in 2HDM

CP violation may occur in 2HDM only if  $Z_2$  symmetry is broken [8, 2, 3, 4, 9]. A necessary condition for an *explicit CP violation* in the Higgs potential  $V$  is an existence of complex parameters. However, if there exists a reparametrization leading to  $V$  with only real parameters (*real basis*), then there is no explicit CP violation in  $V$ . A spontaneous CP breaking, by the vacuum state, is still possible [7, 8, 2].

In the simply analysis [14], which results we present here, only the potential with exact and softly broken  $Z_2$  symmetry was considered, i.e.  $\lambda_{6,7} = 0$ . In studying 2HDM with an *explicit* CP conservation or violation the *real vacuum representation* [4] was applied. A spontaneous CP violation was discussed assuming the *explicitly CP conserving*  $V$ .

### 1.4 Vacuum expectation values

The most general vacuum (extremum) state can be described by [8, 11, 12, 13, 14]

$$\langle \Phi_1 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_1 \end{pmatrix}, \quad \langle \Phi_2 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} u \\ v_2 e^{i\xi} \end{pmatrix}, \quad (4)$$

where  $v_1, v_2, \xi, u \in \mathbb{R}$ . By gauge transformation one can always make  $v_1 > 0$ . Below we will assume that  $v_2 \neq 0$ , with  $v^2 = v_1^2 + v_2^2 = (246 \text{ GeV})^2$ , and  $0 \leq \xi < 2\pi$ .

For vacuum with  $u \neq 0$  the electric charge is not conserved and the photon becomes a massive particle ("charged vacuum"). If  $u = 0$  then a "neutral vacuum" are possible. Depending on the value of  $\xi$  there may or may not be a spontaneous CP violation [8, 3, 12, 13]. The useful quantity is  $\nu = \frac{m_{12}^2}{2v_1 v_2}$  (or  $\nu = \frac{\Re m_{12}^2}{2v_1 v_2}$ ) [4], which here is taken to be positive.

### 1.5 Extremum conditions

For the extremum states (4) the first derivatives of the considered potential lead to the following set of extremum conditions:

$$0 = u [v_1 v_2 \cos \xi (\lambda_4 + \lambda_5) - m_{12}^2], \quad 0 = u [\lambda_2 (u^2 + v_2^2) + \lambda_3 v_1^2 - m_{22}^2] \quad (5)$$

$$0 = v_2 \sin \xi [2\lambda_5 v_1 v_2 \cos \xi - m_{12}^2], \quad 0 = v_2 \sin \xi [v_1^2 (\lambda_3 + \lambda_4 - \lambda_5) + \lambda_2 (u^2 + v_2^2) - m_{22}^2] \quad (6)$$

$$0 = v_1 [v_2^2 (\lambda_5 \cos^2 2\xi + \lambda_4) + \lambda_1 v_1^2 + \lambda_3 (u^2 + v_2^2) - m_{11}^2] - m_{12}^2 v_2 \cos \xi \quad (7)$$

$$0 = uv_1 v_2 \sin \xi (\lambda_4 - \lambda_5), \quad 0 = v_2 \cos \xi [v_1^2 (\lambda_3 + \lambda_4 + \lambda_5) + \lambda_2 (u^2 + v_2^2) - m_{22}^2] - m_{12}^2 v_1 \quad (8)$$

If  $u = 0$  then above conditions are satisfied for an exact  $Z_2$  symmetry ( $m_{12}^2 = 0$ ) when the only possible neutral vacuum state is the one which respects CP, i.e. with  $\sin \xi = 0$ , and for a broken  $Z_2$  symmetry. In the latter case two neutral vacuum states are possible - without

and with CP violation, for  $\sin \xi = 0$  and  $\sin \xi \neq 0$ , respectively. To get a real minimum of the potential the eigenvalues of the squared mass matrix have to be positive. We will assume in addition that positivity constraints hold guaranteeing stability of the vacuum [10].

### 1.6 Physical regions for CP conserving 2HDM

Expressions for masses of  $H^\pm$  and  $A$  for 2HDM with an explicit or a spontaneous CP conservation are as follows.

*$Z_2$  symmetry broken* If  $Z_2$  symmetry is softly broken ( $\nu \neq 0$ ), then the masses squared of  $H^\pm$  and  $A$  are given by:

$$M_{H^\pm}^2 = v^2 \left( \nu - \frac{1}{2}(\lambda_4 + \lambda_5) \right), \quad M_A^2 = v^2 (\nu - \lambda_5). \quad (9)$$

In order to have positive  $M_{H^\pm}^2$  and  $M_A^2$  inequalities  $\lambda_5 + \lambda_4 < 2\nu$  and  $\lambda_5 < \nu$  should hold.

Large masses for  $H^\pm$  and  $A$  (9) can arise from large  $\nu$ . In the limit  $\nu \rightarrow \infty$  the decoupling is realized -  $h$  is like the Higgs boson in the Standard Model, while  $H^\pm, A, H$  are heavy and almost degenerate [3, 4].

*Exact  $Z_2$  symmetry* The results for an exact  $Z_2$  symmetry can be obtained from above expressions in the limit  $\nu \rightarrow 0$ . Then  $\lambda_5 < 0$ . Masses cannot be too large, as here they can arise only due to  $\lambda'$ s. However, large  $\lambda'$ s may violate tree-level unitarity constraints [15].

### 1.7 Physical regions for CP violating 2HDM

As it was mentioned above if the 2HDM potential breaks  $Z_2$  symmetry then CP violation may be realized in the model. Note, that if CP is violated physical neutral Higgs states are  $h_1, h_2, h_3$ , without definite CP properties, while  $h, H, A$  are useful but only auxiliary states.

*Explicit CP violation* If there is explicit CP violation all formulae derived for the CP conservation case (9 and beyond) hold after the replacements:  $\lambda_5 \rightarrow \Re \lambda_5$  and  $m_{12}^2 \rightarrow \Re m_{12}^2$ . Note, that the decoupling can be realized here as well, with large  $M_{H^\pm}^2$  arising from large  $\nu$ .

*Spontaneous CP violation* Spontaneous CP violation may appear if there is a CP breaking phase of the VEV, so  $\sin \xi \neq 0$ . From the extremum condition one gets that:

$$\cos \xi = \frac{m_{12}^2}{\lambda_5 2v_1 v_2} = \frac{\nu}{\lambda_5}, \quad (10)$$

from which it follows that  $|\nu/\lambda_5| < 1$ . The squared masses for  $H^\pm$  and  $A$  are given by the following expressions, see also [13]:

$$M_{H^\pm}^2 = \frac{v^2}{2} (\lambda_5 - \lambda_4), \quad M_A^2 = \frac{v^2}{\lambda_5} (\lambda_5^2 - \nu^2) = v^2 \lambda_5 \sin^2 \xi. \quad (11)$$

We see that they are quite different from the formulae for  $M_{H^\pm}^2$  and  $M_A^2$  discussed above. (Note, that although  $A$  is no longer a physical state, positivity of  $M_A^2$  still provides a good constraint since it gives at the same time a condition for positivity of squared masses of physical particles.) From the last expression for  $M_A^2$  (11) it is easy to see that  $\lambda_5$  have to be positive. Furthermore, squared masses (11) are positive if  $\lambda_5 > \lambda_4$  and  $\lambda_5 > \nu > 0$ .

It is worth mentioning that the squared mass of  $H^\pm$  does not depend on  $\nu$  at all. Therefore,  $M_{H^\pm}$  cannot be too large in 2HDM with CP violated spontaneously, for the same reason as in the discussed above case of exact  $Z_2$  symmetry.

### 1.8 Conclusion on possible vacuum states in 2HDM

Regions where various vacuum states (conserving or spontaneously violating CP) can be realized in 2HDM are mutually exclusive [10, 12, 13, 14]. The mass of charged Higgs boson may serve as a guide over various regimes of the 2HDM. Existence of heavy charged Higgs boson, with mass above 600-700 GeV [4, 14], would be a signal that in 2HDM  $Z_2$  symmetry is violated, and CP can be violated only explicitly.

## 2 Experimental constraints on the 2HDM(II) with CP conservation

Here we consider the CP conserving 2HDM, assuming that  $Z_2$  symmetry is extended also on the Yukawa interaction, which allows to suppress the FCNC [16]. We limit ourself to constraints on the Model II of the Yukawa interaction, as in MSSM, see e.g. [17]. There are 7 parameters for the potential with softly breaking  $Z_2$  symmetry: masses  $M_h, M_H, M_A, M_{H^\pm}$ , mixing angles  $\alpha$  and  $\tan\beta = v_2/v_1$ , and parameter  $\nu$ .

Couplings (relative to the corresponding couplings of the SM Higgs) are as follows:

	$\boxed{h}$	$\boxed{A}$
to W/Z:	$\chi_V = \sin(\beta - \alpha)$	$0$
to down quarks/charged leptons:	$\chi_d = \chi_V - \sqrt{1 - \chi_V^2} \tan\beta$	$-i\gamma_5 \tan\beta$
to up quarks:	$\chi_u = \chi_V + \sqrt{1 - \chi_V^2} / \tan\beta$	$-i\gamma_5 / \tan\beta$

$H$  couples like  $h$  with following replacements:  $\sin(\beta - \alpha) \rightarrow \cos(\beta - \alpha)$  and  $\tan\beta \rightarrow -\tan\beta$ . For large  $\tan\beta$  there are enhanced couplings to  $d$ -type fermions. Note, that coupling  $\chi_{VH^\pm}^h = \cos(\beta - \alpha)$  is complementary to the  $\chi_V^h$ .

Important constraints on mass of charged Higgs boson in 2HDM (II) are coming from the  $b \rightarrow s\gamma$  and  $B \rightarrow \tau\nu$  decays. The rate for the first process calculated at the NNLO accuracy in the SM [18], after a comparison with the precise data from BaBar and Belle, leads to the constraint:  $M_{H^\pm} > 295$  GeV at 95 % CL for  $\tan\beta > 2$ . This limit together with the constraints from the tree-level analysis of  $B \rightarrow \tau\nu$  [19] is presented in Fig.1 (Left).

The 2HDM analysis has been performed at the one-loop level for the leptonic tau decays [20]. The constraints are shown in Fig.1 (Right). Not only lower, but also in the non-decoupling scenario upper limits can be derived here. In contrast to the mentioned results from  $b$  decays here the (one-loop) constraints depend on masses of neutral Higgs bosons.

## 3 Acknowledgment

MK is grateful to Ilya Ginzburg and Rui Santos for important discussions.

## References

- [1] Slides:  
<http://ilcagenda.linearcollider.org/contributionDisplay.py?contribId=454&sessionId=718&confId=1296>
- [2] G. C. Branco, L. Lavoura and J. P. Silva, ‘CP violation,’ *Oxford, UK: Clarendon (1999) 511 p*
- [3] J. F. Gunion and H. E. Haber, Phys. Rev. D **67** (2003) 075019, Phys. Rev. D **72** (2005) 095002.
- [4] I. F. Ginzburg and M. Krawczyk, Phys. Rev. D **72**, 115013 (2005).
- [5] L. Lavoura, Phys. Rev. D **50** (1994) 7089.
- [6] E. Accomando *et al.*, arXiv:hep-ph/0608079.
- [7] T. D. Lee, Phys. Rev. D **8** (1973) 1226.




Figure 1: Left: Constraints from  $B \rightarrow \tau\nu_\tau$  and  $b \rightarrow s\gamma$  data on the charged Higgs boson mass as a function of  $\tan\beta$  in 2HDM (II) [19]; Right: Limits from the leptonic  $\tau$  decay for  $M_h = 20$  GeV and  $\chi_V^h = 0$  in 2HDM(II): tree-level exclusion of a region below the straight line  $M_{H^\pm} \geq 1.71 \tan\beta$  GeV and one-loop exclusion of the region above the curve  $\Delta \sim \tan\beta^2 [\ln \frac{M_h}{M_{H^\pm}} + 1]$ . The excluded region lies on the right on the curves: bold for  $M_A = M_{H^\pm}$ , dotted for  $M_A = 100$  GeV. Exclusion from  $\tau \rightarrow e\nu_\tau\bar{\nu}_e$  is represented by dashed line [20].

- [8] G. C. Branco, Phys. Rev. Lett. **44** (1980) 504. G. C. Branco, Phys. Rev. D **22** (1980) 2901.
- [9] I. P. Ivanov, Phys. Lett. B **632** (2006) 360, Phys. Rev. D **75**, 035001 (2007) [Erratum-ibid. D **76**, 039902 (2007)], arXiv:0710.3490 [hep-ph];  
M. Maniatis, A. von Manteuffel and O. Nachtmann, arXiv:0707.3344 [hep-ph];  
C. C. Nishi, Phys. Rev. D **76**, 055013 (2007), Phys. Rev. D **74**, 036003 (2006).
- [10] N. G. Deshpande and E. Ma, Phys. Rev. D **18**, 2574 (1978).
- [11] J. L. Diaz-Cruz and A. Mendez, Nucl. Phys. B **380**, 39 (1992).
- [12] A. Barroso, P. M. Ferreira and R. Santos, Phys. Lett. B **652**, 181 (2007);  
A. Barroso, P. M. Ferreira, R. Santos and J. P. Silva, Phys. Rev. D **74**, 085016 (2006);  
A. Barroso, P. M. Ferreira and R. Santos, Phys. Lett. B **632**, 684 (2006), Phys. Lett. B **603**, 219 (2004) [Erratum-ibid. B **629**, 114 (2005)];  
J. Velhinho, R. Santos and A. Barroso, Phys. Lett. B **322**, 213 (1994).
- [13] I. F. Ginzburg and K. A. Kanishev, arXiv:0704.3664 [hep-ph].
- [14] M. Krawczyk and D. Sokolowska - in preparation;  
D. Sokolowska, Master Diploma, Dept. of Physics, U. of Warsaw, July 2007
- [15] I. F. Ginzburg and I. P. Ivanov, Phys. Rev. D **72**, 115010 (2005);  
A. G. Akeroyd, A. Arhrib and E. M. Naimi, Phys. Lett. B **490**, 119 (2000)
- [16] S. L. Glashow and S. Weinberg, Phys. Rev. D **15** (1977) 1958;  
E. A. Paschos, Phys. Rev. D **15** (1977) 1966.
- [17] M. Krawczyk, Acta Phys. Polon. B **33**, 2621 (2002).
- [18] M. Misiak *et al.*, Phys. Rev. Lett. **98**, 022002 (2007).
- [19] D. s. Du, arXiv:0709.1315 [hep-ph], M. Nakao, talk given at LP2007, Korea.
- [20] M. Krawczyk and D. Temes, Eur. Phys. J. C **44**, 435 (2005).