arXiv:0712.3566v1 [cond-mat.str-el] 20 Dec 2007

Abstract

Recent achievements in experiments with cold fermionic atoms indicate the potential for developing novel superconducting devices which may be operated in a wide range of regimes, at a level of precision previously not available. Unlike traditional, solid-state superconducting devices, the cold-atom systems allow the fast switching on of the BCS phase, and the observation of non-equilibrium, coherent oscillations of the order parameter. The integrable and non-linear nature of the equations of motions makes this operating regime particularly rich in potential applications, such as quantum modulation and encoding, or nonlinear mixing of quantum coherent oscillations, to name only two. From a mathematical point of view, such systems can be described using the Knizhnik-Zamolodchikov-Bernard equation, or more generally, by the matrix Kadomtsev-Petviashvilii integrable hierarchy. This identification is particularly useful, since it allows a direct application of the known non-linear phenomena described by particular solutions of these equations. Other important features of this formulation, such as the relation to the spin Calogero-Sutherland model, also have relevant physical interpretations. In this work, a complete description of these relationships is presented, along with their potential practical consequences.

Coherent oscillations in superconducting cold Fermi atoms and their applications

Razvan Teodorescu
Theoretical Division and Center for Nonlinear Studies,
Los Alamos National Laboratory

1 Introduction

The operating parameters specific to experiments with cold fermionic gases exhibiting the paired BCS state [31–38] allow the investigation of previously inaccessible regimes, as suggested in [17]. It was shown that for such systems, the time scales of the order parameter $\tau_\Delta \sim |\Delta|^{-1}$, and the quasiparticle energy relaxation time τ_ϵ are both much larger than typical time for switching on the pairing interaction τ_0, essentially given by the variation of external parameters, such as detuning from the Feshbach resonance. It was argued that in this regime, for times $t \ll \tau_\epsilon$, the dynamics of the system is given by non-linear, non-dissipative equations describing the coherent BCS fluctuations for the system out of equilibrium. In this limit, the system is integrable, and features non-perturbative behavior, such as soliton-type solutions.

In the mean-field limit, such non-trivial solutions describing the collective mode of the Anderson spins [39] were derived in [17], for a two-level effective system. This work was generalized [15] in algebro-geometrical terms.

In [40–42], the long-time behavior of the solution has been considered, under various conditions. An issue that remains incompletely addressed so far is the relaxation of the nonlinear oscillatory solution induced by perturbations of the spectral curve, physically justified by coupling to the environment. Several possible kinds of perturbations may be considered, which may lead to different types of relaxation.

In this article, we describe the mathematical structure underlying the integrable quantum pairing problem, and its relations with the theory of isomonodromic deformations of differential equations and other integrable models of strongly interacting particles. We also explain the classical limit of the quantum hamiltonian, and give the explicit construction of the solution in this case. Possible applications due to the integrability of the model are listed in the conclusion of the article.

The paper is organized as follows: in section 2, the Richardson-Gaudin model and its relation to other integrable models are reviewed. Section 3 describes the classical limit and explicit multi-phase solution. In section 4, a few possible applications are briefly discussed.

2 The Richardson-Gaudin Model

As argued for the first time in [17], optically trapped cold Fermi gases in the BCS state are correctly described by the Richardson-Gaudin quantum pairing model. Using this model is justified by the discreteness of the energy spectrum, and by the spatially uniform profile characterizing such systems. This integrable system, introduced by Richardson and Sherman [1–6] in the context of nuclear physics, has received revived interest in recent years, after being applied to metallic superconducting grains [16]. The model is intimately related [18–20] to a class of integrable systems generally referred to as Gaudin magnets [7]. These systems have been studied both at quantum and classical level [21–30], in the elliptic case as well as trigonometric and rational degenerations, using various methods from integrable vertex models to singular limits of Chern-Simons theory.

In this section, we recall the main features of this model, as well as their physical interpretations.

2.1 The quantum pairing hamiltonian

Following [21], we briefly review the Richardson pairing model. It describes a system of n fermions characterized by a set of independent one-particle states of energies ϵ_l, where the label l takes values from a set Λ. The labels may refer, for instance, to orbital angular momentum eigenstates. Each state l has a total degeneracy d_l, and the states within the subspace corresponding to l are further labeled by an internal quantum number s. For instance, if the quantum number l labels orbital momentum eigenstates, then $d_l = 2l + 1$ and $s = -l, \ldots, l$. However, the internal degrees of freedom can be defined independently of l. We will assume that d_l is even, so for every state (ls), there is another one related by time reversal symmetry $(l\bar{s})$. For simplicity, we specialize to the case $d_l = 2, s = \uparrow, \downarrow$. Let $\hat{c}^\dagger_{ls}$ represent the fermionic creation operator for the state (ls). Using the Anderson pseudo-spin operators [39] (quadratic pairing operators), satisfying the $su(2)$ algebra

$$[t_i^3,\, t_j^\pm] = \pm\delta_{ij} t_j^\pm, \quad [t_i^+,\, t_j^-] = 2\delta_{ij} t_j^3, \tag{1}$$

the Richardson pairing hamiltonian is given by

$$H_P = \sum_{l\in\Lambda} 2\epsilon_l t_l^3 - g\sum_{l,l'} t_l^+ t_{l'}^- = \sum_{l\in\Lambda} 2\epsilon_l t_l^3 - g\mathbf{t}^+\cdot\mathbf{t}^-, \tag{2}$$

where $\mathbf{t} = \sum_l \mathbf{t}_l$ is the total spin operator. It maps to the reduced BCS model

$$\hat{H} = \sum_{\mathbf{p},\sigma} \epsilon_{\mathbf{p}} \hat{c}^\dagger_{\mathbf{p},\sigma}\hat{c}_{\mathbf{p},\sigma} - g\sum_{\mathbf{p},\mathbf{k}} \hat{c}^\dagger_{\mathbf{p}\uparrow}\hat{c}^\dagger_{-\mathbf{p}\downarrow}\hat{c}_{-\mathbf{k}\downarrow}\hat{c}_{\mathbf{k}\uparrow} \tag{3}$$

by replacing the translational degrees of freedom by rotational ones, where $l \in \Lambda = \{1, \ldots n\}$ ennumerates the one-particle orbital degrees of freedom, while $s = \uparrow, \downarrow$ indicates the two internal spin states per orbital ($d_l = 2$). The pairing hamiltonian can be decomposed into the linear combination

$$H_P = 2\sum_{l\in\Lambda} \epsilon_l R_l + g\left[\left(\sum_{l\in\Lambda} t_l^3\right)^2 - \frac{1}{4}\sum_{l\in\Lambda}(d_l^2 - 1)\right]. \tag{4}$$

At a fixed value of the component t^3 of the total angular momentum, the last term becomes a constant and is dropped from the hamiltonian. The operators R_l (generalized Gaudin magnets [7]) are given by

$$R_l = t_l^3 - \frac{g}{2}\sum_{l'\neq l}\frac{\mathbf{t}_l\cdot\mathbf{t}_{l'}}{\epsilon_l-\epsilon_{l'}}. \tag{5}$$

These operators solve the Richardson pairing hamiltonian because [18] they are independent, commute with each other, and span all the degrees of freedom of the system. Richardson showed [1, 2] that the exact $N-$pair wavefunction of his hamiltonian is given by application of operators $b_k^\dagger = \sum_l \frac{t_l^\dagger}{2\epsilon_l - e_k}$ to vacuum (zero pairs state). The unnormalized $N-$pair wavefunction reads $\Psi_R(\epsilon_i) = \prod_{k=1}^N b_k^\dagger|0\rangle$. The eigenvalues e_k satisfy the self-consistent algebraic equations

$$\frac{1}{g} = \sum_{p\neq k}\frac{2}{e_k-e_p} + \sum_l \frac{1}{2\epsilon_l - e_k}, \tag{6}$$

which can be given a 2D electrostatic interpretation [21] with energy

$$U(\epsilon_l, e_k) = \frac{2}{g}\left[\sum_{k=1}^N \mathcal{R}e(e_k) - \sum_{l=1}^n \mathcal{R}e(\epsilon_l)\right] + \tag{7}$$

$$2\sum_{l=1}^n\sum_{k=1}^N \log|e_k - 2\epsilon_l| - 4\sum_{k<p}\log|e_k-e_p| - \sum_{i<j}\log|2\epsilon_i - 2\epsilon_j| \tag{8}$$

Equations (6) appear as equilibrium conditions for a set of charges of strength $q=2$ placed at points e_k, in the presence of fixed charges of strength $q=-1$ at points $2\epsilon_l$, and uniform electric field of strength $\frac{1}{g}$, pointing along the real axis. This interpretation proves to be very useful for the conformal field theory (CFT) description of the Richardson problem. The electrostatic energy (7) is minimized for values $\{e_k\}$ corresponding to pair energies. In (7), n, N represent the number of single-particle levels and the number of pair energies, respectively. For large interaction constant g, the equilibrium positions $\{e_k\}$ form a set of complex conjugated pairs defining a curve γ in the complex plane of energies. We note that the eigenvalues r_i of Gaudin hamiltonia are proportional in this language to the values of the electric field at positions $2\epsilon_i$, $2r_i = g\frac{\partial U}{\partial \epsilon_i}$.

For a set of single-particle energies $\{\epsilon_i\}$, the BCS ground state is obtained by minimizing the electrostatic energy (7) with respect to positions of the free charges at $\{e_k\}$. Once found, they also determine exactly the values of the electric field at positions $\{2\epsilon_i\}$ on the real axis, which are proportional to the ground-state eigenvalues $\{r_i^{GS}\}$. For any other values of $\{r_i\}$, the electrostatic energy (7) is not minimized. This indicates that for arbitrary values $r_i \neq r_i^{GS}$, the system is not in equilibrium.

2.2 Richardson-Gaudin model as limit of KZB equations

In this section we recall the relation between the Richardson-Gaudin model and singular $SU(2)$ Chern-Simons and WZW models [25, 26]. These relations stem from the study of quantum states of the Chern-Simons theory with gauge group $SU(2)$ on the

manifold $\mathbb{T}^2 \times \mathbb{R}$ and in the presence of Wilson lines $\{z_i\} \times \mathbb{R}$, $i = 1, \ldots, n$ [25], corresponding to the one-particle spectrum of the Richardson hamiltonian, $z_i = 2\epsilon_i$. The torus $\mathbb{T}^2$ has modular parameter $\tau = \tau_1 + i\tau_2,\ \tau_{1,2} \in \mathbb{R}$. Thus, $\mathbb{T}^2 = \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z}$. The quantum states of this theory are known to satisfy the Knizhnik-Zamolodchikov-Bernard (KZB) equations

$$\nabla \Psi_{CS} = 0, \quad \nabla = (\nabla_\tau, \nabla_{z_i}), \tag{9}$$

where ∇ is the flat KZB connection,

$$\nabla_\tau = \kappa\partial_\tau + H_0(\tau, z_i), \quad \nabla_{z_i} = \kappa\partial_{z_i} + H_i(\tau, z_i), \tag{10}$$

and H_0, H_i are elliptic versions of the Gaudin hamiltonia. The parameter κ is related to the level of representation of $SU(2)$. By a limiting procedure, the system degenerates into the Richardson-Gaudin problem: by taking the limit $\tau \to i\infty$, the torus degenerates into a cylinder, and the elliptic hamiltonia H_i, H_0 take a trigonometric limit. Upon rescaling the cylinder such that the set $\{z_i\}$ is collapsed into the origin, the cylinder becomes the complex plane with punctures at $\{z_i\}$, and the rescaled operators become the rational Gaudin magnets (5).

2.3 Singular representations of Wess-Zumino-Witten theory

Given the well-known relation between $2+1$ Chern-Simons theory with $SU(2)$ gauge group and WZW theory, it is not surprising that a non-Abelian CFT representation for Richardson-Gaudin models is also available. In fact, the relationship between these theories is already transparent by identifying the electrostatic picture of R-G with a Coulomb gas representation of WZW.

In the CFT approach [16], a primary field of spin $1/2$ of the $SU(2)$ representation of WZW is associated to every energy level of the R-G problem. The background charge α_0 and the level of Kac-Moody algebra $SU(2)_k$ are related by $k = \frac{1}{2\alpha_0^2} - 2$. It turns out that the R-G model is retrieved in the singular limit $\alpha_0 \to \infty$, corresponding to $k = -2$. In this limit, the representations are given by *perturbed* WZW blocks (PWZW),

$$\Psi_P(z_i) = \oint_{\mathcal{C}_1} de_1 \oint_{\mathcal{C}_N} de_N e^{-\alpha_0^2 U(z_i, e_k)} \Psi_R(z_i, e_k). \tag{11}$$

In (11), e_k are the eigenvalues of the Richarsdon hamiltonian with N fermion pairs, α_0 is the background charge in the Coulomb gas representation of the WZW model associated with the $2+1$ Chern-Simons theory described above, and $U(z_i, e_k)$ is the holomorphic piece of the 2D electrostatic potential of the Coulomb gas representation (7). The Richardson wave function $\Psi_R(z_i, e_k)$ is obtained by applying Bose fields corresponding to positions z_i, e_k,

$$\Psi_R(z_i, e_k) = \langle \prod_{i=1}^{n} \gamma(z_i) \prod_{k=1}^{N} \beta(e_k) \rangle, \tag{12}$$

and with correlator $\langle \beta(z)\gamma(z') \rangle = \frac{1}{z - z'}$.

The singular limit described above in the language of the Chern-Simons theory corresponds to the limit $\alpha_0 \to \infty$ in the CFT approach. The perturbed WZW blocks

are dominated by the saddle-point configurations, given by the eigenvalue equations $\partial_{e_k} U = 0$, and become proportional to the Richardson wavefunction. In the singular limit, the KZB equations takes the form

$$\left(\frac{1}{2\alpha_0^2} \partial_{z_i} - R_i \right) \Psi_{PWZW} = 0, \tag{13}$$

which shows that the Richardson wavefunction is an eigenstate of the Gaudin hamiltonia, with eigenvalues given by the solutions to the electrostatics problem (7).

Equation (13) is the analog of the singular limit of the KZB equation (9), in the limit $\kappa \to 0$, corresponding to $\alpha_0 \to \infty$. Likewise, there is a dirrect correspondence between the Chern-Simons states and the perturbed WZW blocks. We will see in the following sections that these equations survive in the semiclassical limit and that they allow the identification of the tau function for the corresponding classical integrable system.

2.4 Relation with the KP hierarchy and other integrable models

The zero curvature conditions for the KZB connection can be interpreted as a truncation of the general Kadomtsev-Petviashvilii (KP) integrable hierarchy, at a lelvel set by the number of eigenvalues in the single-particle spectrum. This is also illustrated by the fact that, as shown in the preceding paragraphs, in the genus one case, the Richardson-Gaudin problem is equivalent with the elliptic Calogero model. This model is known to be embedded into the KP hierarchy, and admits simple-pole solutions built from the solution of KP. In fact, all the relationships described so far are consequences of an even more general formulation of the Richardson-Gaudin problem as a hierarchy of isomonodromic deformations with respect to changing the moduli of its spectral curve.

These relationships allow to classify the solutions to the quantum Richardson-Gaudin model using degenerate limits of the solutions for the elliptic Calogero problem. Such solutions are known (for instance, the simple-pole solutions mentioned above) and would therefore allow (in principle) constructing exact solutions for given initial conditions. Even more importantly, there is a large body of knowledge concerning the classes of solutions supported by the KP hierarchy (periodic, algebraic-geometric or rational), which translates into distinct "phases" (albeit we are dealing with non-equilibrium, coherent oscillations) supported by the quantum pairing model, depending on the initial conditions.

3 The mean-field limit of Richardson-Gaudin models

Throughout this section, we consider the classical limit of the quantum pairing model. The main reason for this approximation is the fact that explicit, generic solutions can be obtained in this case.

3.1 General description of the classical model

In the mean-field limit, the spin operators $\mathbf{t}_l$ are replaced by their quantum mechanical averages. Written in terms of the classical vectors $\mathbf{S}_l = 2\langle \mathbf{t}_l \rangle$, the semiclassical




Figure 1: Schematic representation of the Liouville torus for the integrable system (16).

approximation for the pairing hamiltonian becomes

$$H_{MF} = \sum_{l\in\Lambda} \epsilon_l S_l^3 - \frac{g}{4}|J^-|^2, \tag{14}$$

where $\mathbf{J} = \sum_{l\in\Lambda} \mathbf{S}_l$ and the BCS gap function is given by $\Delta = gJ^-/2$. Replacing commutators by canonical Poisson brackets,

$$\{S_i^\alpha,\, S_j^\beta\} = 2\epsilon^{\alpha\beta\gamma} S_i^\gamma \delta_{ij}, \tag{15}$$

variables S_i^α become smooth functions of time. In this limit, the problem can analyzed with tools of classical integrable systems, and the solution is known to be exact as $n \to \infty$.

The Poisson brakets (15) and hamiltonian (14) lead to the equations of motion

$$\dot{\vec{S}}_i = 2(-\vec{\Delta} + \epsilon_i \hat{z}) \times \vec{S}_i, \tag{16}$$

where $2\vec{\Delta} = (gJ_x, gJ_y, 0)$ and $\vec{J}$ is the total spin. The semiclassical limit of Gaudin hamiltonia are independent constants of motion,

$$r_i = \frac{1}{2}\left[S_i^z - \frac{g}{2} \sum_{j\neq i} \frac{\vec{S}_i \cdot \vec{S}_j}{\epsilon_i - \epsilon_j} \right], \quad \dot{r}_i = 0. \tag{17}$$

Equations (16) describe a set of strongly interacting spins and have generic non-linear oscillatory solutions. The exact solution may be obtained through the Abel-Jacobi inverse map, explained in the next section.

3.2 Isospectral case and the Abel-Jacobi inversion problem

In [9, 15], the system (16) with fixed spectral curve was solved through inverse Abel-Jacobi mapping, by using Sklyanin separation of variables techniques [8, 52]. Interesting connections to generalized Neumann systems and Hitchin systems were discovered

in [24]. The solution starts from the Lax operator

$$\mathcal{L}(\lambda) = \frac{2}{g}\sigma_3 + \sum_{i=0}^{n} \frac{\vec{S}_i \cdot \vec{\sigma}}{\lambda - \epsilon_i} = \begin{bmatrix} a(\lambda) & b(\lambda) \\ c(\lambda) & -a(\lambda) \end{bmatrix}, \tag{18}$$

where $\sigma_\alpha,\ \alpha = 1, 2, 3$ are the Pauli matrices, and λ is an additional complex variable, the spectral parameter. Let $u_k,\ k = 1, \ldots, n-1$ be the roots of the coefficient $c(\lambda)$. Poisson brackets for variables S_i^α read

$$\{S_j^\alpha,\ S_k^\beta\} = 2\epsilon_{\alpha\beta\gamma} S_k^\gamma \delta_{jk} \tag{19}$$

The Lax operator (18) defines a Riemann surface (the spectral curve) $\Gamma(y, \lambda)$ of genus $g = n - 1$, through

$$y^2 = Q(\lambda) = \det \mathcal{L}(\lambda) \left[g \frac{P(\lambda)}{2} \right]^2, \tag{20}$$

where $P(\lambda) = \prod_{i=1}^{n} (\lambda - \epsilon_i)$.

The equations of motion for the hamiltonian (14) become

$$\dot{u}_i = \frac{2iy(u_i)}{\prod_{j \neq i}(u_i - u_j)}, \quad i\dot{J}^- = J^- \left[gJ^3 + 2\sum_{k=1}^{n} \epsilon_k - u \right]. \tag{21}$$

In (21), $u = -2\sum_{i=1}^{n-1} u_i, \quad b(u_i) = 0$.

From the equations of motion, it is clear that knowledge of the initial amplitude of J^- and of the roots $\{u_i\}$ is enough to specify the n unit vectors $\{\mathbf{S}_i\}$, for a given set of constants of motion $\{R_l\}$ given by the classical limit of Gaudin hamiltonia. The Dubrovin equations (21) are solved by the inverse of the Abel-Jacobi map, as we explain in the following. We begin by noting that the polynomial $Q(\lambda)$ has degree $2n$, and is positively defined on the real λ axis. Therefore, the curve $\Gamma(y, \lambda)$ has n cuts between the pairs of complex roots $[E_{2i-1},\ E_{2i}], i = 1, 2, \ldots, n$, perpendicular to the real λ axis. The points u_i belong to $n-1$ of these cuts, $u_i \in [E_{2i-1},\ E_{2i}], i = 1, \ldots, n-1$. These $g = n - 1$ cuts allow to define a canonical homology basis of Γ, consisting of cycles $\{\alpha_i, \beta_i\}, i = 1, \ldots, g$. With respect to these cycles, a basis of normalized holomorphic differentials $\{\omega_i\}$ can be defined, through

$$\mu_i = \lambda^{g-i} \frac{d\lambda}{y}, \ \ M_{ij} = \int_{\alpha_j} \mu_i, \ \ \boldsymbol{\omega} = M^{-1}\boldsymbol{\mu}. \tag{22}$$

The period matrix $B_{ij} = \int_{\beta_j} \omega_i$ is symmetric and has positively defined imaginary part. The Riemann θ function is defined with the help of the period matrix as

$$\theta(\boldsymbol{z}|B) = \sum_{\boldsymbol{n} \in \boldsymbol{z}^g} e^{2\pi i(\boldsymbol{n}^t \boldsymbol{z} + \frac{1}{2}\boldsymbol{n}^t B \boldsymbol{n})}. \tag{23}$$

The g vectors $\boldsymbol{B}_k$ consisting of columns of B and the basic vectors $\boldsymbol{e}_k$ define a lattice in $\mathbb{C}^g$. The $Jacobian$ variety of the curve Γ, is then the $g-$dimensional torus defined as the quotien $J(\Gamma) = \mathbb{C}^g/(\mathbb{Z}^g + B\mathbb{Z}^g)$. The Abel-Jacobi map associates to any point P on Γ, a point ($g-$ dimensional complex vector) on the Jacobian variety, through $\boldsymbol{A}(P) = \int_\infty^P \boldsymbol{\omega}$. Considering now a $g-$dimensional complex vector of points

$\{P_k\}, k = 1, \ldots, g$ on Γ, defined up to a permutation, we can associate to it the point on the Jacobian

$$\boldsymbol{z} = \boldsymbol{a}(\boldsymbol{P}) = \sum_{k=1}^{g} \boldsymbol{A}(P_k) + \boldsymbol{K}, \tag{24}$$

where $\boldsymbol{K}$ is the Riemann characteristic vector for Γ.

The map (24) suggests that we now have a way to describe the dynamics on Γ by following the image point on the Jacobian. Given a point on the $g-$dimensional Jacobian $\boldsymbol{z} = (\zeta_1, \ldots, \zeta_{n-1})$, we can find an unique set of points $\{\lambda_k\}, k = 1, \ldots, g$ on Γ, such that $\boldsymbol{z} = \boldsymbol{a}(\boldsymbol{\lambda})$, and $\theta(\boldsymbol{a}(\boldsymbol{P}) - \boldsymbol{z}|B) = 0$. The system evolves in time according to the point $\boldsymbol{z}(t)$

$$\zeta_k = ic_k, \ 1 \leq i \leq g-1, \ \zeta_{n-1} = i(c_{n-1} + t), \tag{25}$$

where $\{c_k\}$ is a set of initial conditions, such that $\boldsymbol{z}_0 = \boldsymbol{z}(t=0) = \boldsymbol{a}(\boldsymbol{c})$, and $\boldsymbol{c}$ is the set of initial conditions for positions of $\boldsymbol{\lambda}$ on Γ. Together with the initial condition which determines the initial amplitude of J^-, this set will determine entirely the evolution of the functions $u_i(t), J^-(t)$.

It is perhaps useful to give a graphical description (Figure 1) of the general solution described above. For a given set of initial conditions for the spins $\{\vec{S}_i\}$, *i.e.* also of the constants of motion $\{r_i\}$, a polynomial $Q(u)$ of degree $2n$ and with n pairs of complex conjugated roots $E_{2k+2} = \overline{E}_{2k+1}$, $k = 0, \ldots, n-1$, is constructed. A schematic representation of these roots is given in Figure 1. Between each pair of roots, we place a simple cut $\mathcal{C}_k = [E_{2k+1}, E_{2k+2}]$ on the complex plane of energies. The surface thus obtained is a representation of a torus of smooth genus $g = n - 1$.

Variables u_k are introduced for $n-1$ of these cuts, with respect to which the equations of motion separate. The variables u_k evolve in time in a complicated fashion, solving a system of nonlinear coupled differential equations (21). Up to a constant, the time dependence of the gap parameter amplitude is given by

$$\log|\Delta(t)| = \mathcal{I}m \int u(t)dt, \quad u(t) = \sum_k u_k(t). \tag{26}$$

The widths of the cuts $\mathcal{C}_k$ and the periods of the non-linear oscillators u_k are determined by the values of constants of motion $\{r_i\}$. For the particular choice $r_i = r_i^{GS}$, all the cuts $\mathcal{C}_k$, $k = 1, \ldots, n-1$ vanish, and the width of the remaining cut $\mathcal{C}_n$ equals the equilibrium value of the gap function: $|E_{2n-1}^{GS} - E_{2n}^{GS}| = 2|\Delta|^{GS}$. In that case, the oscillators $u_k = E_{2k-1} = E_{2k}$ are at rest, and the only time dependence left in the system is the uniform precession of the parallel planar spins S_i^-, with frequency $\omega = 2\sum_{k=1}^{n} \epsilon_k - 2\sum_{p=1}^{n-1} E_{2p-1} - \sum_{i=1}^{n} S_i^z$. In the case of particle-hole symmetry, ω vanishes as well.

4 Conclusion

In the present work, the regime of coherent, non-equilibrium quantum oscillations characterizing the onset of the BCS state in cold Fermi gases was investigated from a mathematical point of view. The integrability of the model and main features of the space of solutions were described, both at the quantum and classical levels.

The most direct application of these coherent quantum oscillations would be in encoding/decoding of information. Although simple in principle, this would in fact require very accurate control over the initial conditions at the onset of the pairing regime, which is unfeasible. However, given the stability of the coherent oscillatory regime, it is clearly easier to first implement the quantum modulation of an arbitrary oscillatory solution, by slowly perturbing the single-particle spectrum. This kind of perturbation is in fact within the same class as the original isomonodromic deformations, and hence does not destroy the integrability of the model.

Another possible application makes use of the non-linear nature of the solution of the Richardson-Gaudin model. Considering only the simplest classical solutions given by the Abel-Jacobi construction, the elliptical solutions, it is possible to devise a non-linear 'mixer' by using the fact that the dynamics is linear on the associated torus (Jacobian). Then, linear operations with respect to the dynamics on the Jacobian would translate into strongly non-linear (but still invertible) effects observed at the level of quantum oscillations.

Acknowledgments

This work was carried out under the auspices of the National Nuclear Security Administration of the U.S. Department of Energy at Los Alamos National Laboratory under Contract No. DE-AC52-06NA25396.

References

[1] R.W. Richardson. *Phys. Lett.*, 3:277, 1963.

[2] R.W. Richardson. *Phys. Lett.*, 5:82, 1963.

[3] R. W. Richardson and N. Sherman. Exact eigenstates of the pairing-force Hamiltonian. *Nuclear Phys.*, 52:221–238, 1964.

[4] R. W. Richardson. Exact eigenstates of the pairing-force Hamiltonian. II. *J. Mathematical Phys.*, 6:1034–1051, 1965.

[5] R.W. Richardson. *Phys. Rev.*, 141:949, 1966.

[6] R.W. Richardson. *J. Math. Phys.*, 9:1327, 1968.

[7] M. Gaudin. Diagonalisation d'une classe d'Hamiltoniens de spin. *J. Physique*, 37(10):1089–1098, 1976.

[8] E. K. Sklyanin. Separation of variables in the Gaudin model. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 164(Differentsialnaya Geom. Gruppy Li i Mekh. IX):151–169, 198, 1987.

[9] A. N. W. Hone, V. B. Kuznetsov, and O. Ragnisco. Bäcklund transformations for the $\mathrm{sl}(2)$ Gaudin magnet. *J. Phys. A*, 34(11):2477–2490, 2001.

[10] V. B. Kuznetsov. Quadrics on Riemannian spaces of constant curvature. Separation of variables and a connection with the Gaudin magnet. *Teoret. Mat. Fiz.*, 91(1):83–111, 1992.

[11] V. Kuznetsov and P. Vanhaecke. Bäcklund transformations for finite-dimensional integrable systems: a geometric approach. *J. Geom. Phys.*, 44(1):1–40, 2002.

[12] E. G. Kalnins, V. B. Kuznetsov, and Willard Miller, Jr. Quadrics on complex Riemannian spaces of constant curvature, separation of variables, and the Gaudin magnet. *J. Math. Phys.*, 35(4):1710–1731, 1994.

[13] V. B. Kuznetsov. Isomorphism of an n-dimensional Neumann system and an n-site Gaudin magnet. *Funktsional. Anal. i Prilozhen.*, 26(4):88–90, 1992.

[14] V. B. Kuznetsov. Equivalence of two graphical calculi. *J. Phys. A*, 25(22):6005–6026, 1992.

[15] E. A. Yuzbashyan, B. L. Altshuler, V. B. Kuznetsov, and V. Z. Enolskii. Nonequilibrium Cooper Pairing in the Non-adiabatic Regime. *ArXiv Condensed Matter e-prints*, May 2005.

[16] G. Sierra. Integrability and conformal symmetry in the BCS model. In *Statistical field theories (Como, 2001)*, volume 73 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 317–328. Kluwer Acad. Publ., Dordrecht, 2002.

[17] R. A. Barankov, L. S. Levitov, and B. Z. Spivak. Collective Rabi Oscillations and Solitons in a Time-Dependent BCS Pairing Problem. *Phys. Rev. Lett.*, 93(16):160401, October 2004.

[18] M. C. Cambiaggio, A. M. F. Rivas, and M. Saraceno. Integrability of the pairing hamiltonian. *Nuclear Physics A*, 624:157–167, February 1997.

[19] L. Amico, A. di Lorenzo, and A. Osterloh. Integrable Model for Interacting Electrons in Metallic Grains. *Phys. Rev. Lett.*, 86:5759–5762, June 2001.

[20] J. Dukelsky, C. Esebbag, and P. Schuck. Class of Exactly Solvable Pairing Models. *Phys. Rev. Lett.*, 87(6):066403, August 2001.

[21] J. Dukelsky, S. Pittel, and G. Sierra. Colloquium: Exactly solvable Richardson-Gaudin models for many-body quantum systems. *Rev. Modern Phys.*, 76(3):643–662, 2004.

[22] M. Asorey, F. Falceto, and G. Sierra. Chern-Simons theory and BCS superconductivity. *Nuclear Phys. B*, 622(3):593–614, 2002.

[23] N. Manojlović and H. Samtleben. Schlesinger transformations and quantum R-matrices. *Comm. Math. Phys.*, 230(3):517–537, 2002.

[24] K. Hikami. Separation of variables in the BC-type Gaudin magnet. *J. Phys. A*, 28(14):4053–4061, 1995.

[25] F. Falceto and K. Gawędzki. Unitarity of the Knizhnik-Zamolodchikov-Bernard connection and the Bethe ansatz for the elliptic Hitchin systems. *Comm. Math. Phys.*, 183(2):267–290, 1997.

[26] F. Falceto and K. Gawędzki. Chern-Simons states at genus one. *Comm. Math. Phys.*, 159(3):549–579, 1994.

[27] G. Ortiz, R. Somma, J. Dukelsky, and S. Rombouts. Exactly-solvable models derived from a generalized Gaudin algebra. *Nuclear Phys. B*, 707(3):421–457, 2005.

[28] D. Korotkin and H. Samtleben. On the quantization of isomonodromic deformations on the torus. *Internat. J. Modern Phys. A*, 12(11):2013–2029, 1997.

[29] J. Harnad. Quantum isomonodromic deformations and the Knizhnik-Zamolodchikov equations. In *Symmetries and integrability of difference equations (Estérel, PQ, 1994)*, volume 9 of *CRM Proc. Lecture Notes*, pages 155–161. Amer. Math. Soc., Providence, RI, 1996.

[30] A. Di Lorenzo, L. Amico, K. Hikami, A. Osterloh, and G. Giaquinta. Quasi-classical descendants of disordered vertex models with boundaries. *Nuclear Phys. B*, 644(3):409–432, 2002.

[31] B. Demarco and D. S. Jin. *Science*, 285:1703, 1999.

[32] B. Demarco, J. L. Bohn, J. P. Burke, M. Holland, and D. S. Jin. Measurement of p-Wave Threshold Law Using Evaporatively Cooled Fermionic Atoms. *Phys. Rev. Lett.*, 82:4208–4211, May 1999.

[33] B. Demarco, S. B. Papp, and D. S. Jin. Pauli Blocking of Collisions in a Quantum Degenerate Atomic Fermi Gas. *Phys. Rev. Lett.*, 86:5409–5412, June 2001.

[34] B. Demarco and D. S. Jin. Spin Excitations in a Fermi Gas of Atoms. *Phys. Rev. Lett.*, 88(4):040405, January 2002.

[35] A. Truscott, K. Strecker, G. Partridge, R. Hulet, and R. Hulet. *Science*, 291:2570, 2001.

[36] T. Loftus, C. A. Regal, C. Ticknor, J. L. Bohn, and D. S. Jin. Resonant Control of Elastic Collisions in an Optically Trapped Fermi Gas of Atoms. *Phys. Rev. Lett.*, 88(17):173201, April 2002.

[37] K. M. O'Hara, S. L. Hemmer, M. E. Gehm, S. R. Granade, and J. E. Thomas. Observation of a Strongly Interacting Degenerate Fermi Gas of Atoms. *Science*, 298:2179–2182, December 2002.

[38] K. M. O'Hara, S. L. Hemmer, S. R. Granade, M. E. Gehm, J. E. Thomas, V. Venturi, E. Tiesinga, and C. J. Williams. Measurement of the zero crossing in a Feshbach resonance of fermionic Li. *Phys. Rev. A*, 66(4):041401, October 2002.

[39] P. W. Anderson. Random-Phase Approximation in the Theory of Superconductivity. *Physical Review*, 112:1900–1916, December 1958.

[40] A. V. Andreev, V. Gurarie, and L. Radzihovsky. Nonequilibrium Dynamics and Thermodynamics of a Degenerate Fermi Gas Across a Feshbach Resonance. *Phys. Rev. Lett.*, 93(13):130402, September 2004.

[41] G. L. Warner and A. J. Leggett. Quench dynamics of a superfluid Fermi gas. *Phys. Rev. B*, 71(13):134514, April 2005.

[42] E. A. Yuzbashyan, O. Tsyplyatyev, and B. L. Altshuler. Relaxation and persistent oscillations of the order parameter in the non-stationary BCS theory. *ArXiv Condensed Matter e-prints*, November 2005.

[43] I. Krichever. Isomonodromy equations on algebraic curves, canonical transformations and Whitham equations. *Mosc. Math. J.*, 2(4):717–752, 806, 2002. Dedicated to Yuri I. Manin on the occasion of his 65th birthday.

[44] I. Krichever. Vector bundles and Lax equations on algebraic curves. *Comm. Math. Phys.*, 229(2):229–269, 2002.

[45] L. A. Dickey. *Soliton equations and Hamiltonian systems*, volume 26 of *Advanced Series in Mathematical Physics*. World Scientific Publishing Co. Inc., River Edge, NJ, 2003.

[46] D. Mumford. *Tata lectures on theta. II*, volume 43 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1984.

[47] K. Takasaki. Spectral curves and Whitham equations in isomonodromic problems of Schlesinger type. *Asian J. Math.*, 2(4):1049–1078, 1998.

[48] G. B. Whitham. *Linear and nonlinear waves*. Pure and Applied Mathematics (New York). John Wiley & Sons Inc., New York, 1999.

[49] A. Beauville. Jacobiennes des courbes spectrales et systèmes hamiltoniens complètement intégrables. *Acta Math.*, 164(3-4):211–235, 1990.

[50] E.D. Belokolos, A.I. Bobenko, V.Z. Enol'skii, A.R. Its, and V.B. Matveev. *Algebro-Geometric Approach to Nonlinear Integrable Equations*, volume 11 of *Springer Series in Nonlinear Dynamics*. Springer-Verlag, Berlin, 1994.

[51] A. R. Its. The Painlevé transcendents as nonlinear special functions. In *Painlevé transcendents (Sainte-Adèle, PQ, 1990)*, volume 278 of *NATO Adv. Sci. Inst. Ser. B Phys.*, pages 49–59. Plenum, New York, 1992.

[52] A. Nakayashiki and F. A. Smirnov. Cohomologies of affine hyperelliptic Jacobi varieties and integrable systems. *Comm. Math. Phys.*, 217(3):623–652, 2001.