# Statistics of harm onic measure and winding of critical curves from conformal eld theory

A. Belikov<sup>1</sup>, I. A. Gruzberg<sup>1</sup>, I. Rushkin<sup>2</sup> <sup>1</sup>Jam es Franck Institute, University of Chicago, Chicago IL 60637 <sup>2</sup>School of Physics, Georgia Institute of Technology, Atlanta, GA 30332

M ay 29, 2008

#### A bstract

Fractal geom etry of random curves appearing in the scaling limit of critical two-dimensional statistical systems is characterized by their harm onic measure and winding angle. The former is the measure of the jaggedness of the curves while the latter quantiles their tendency to form logarithmic spirals. We show how these characteristics are related to local operators of conformal eld theory and how they can be computed using conform allinvariance of critical systems with central charge c 6 1.

# 1 Introduction

G com etric properties of critical two-dimensional systems of statistical medhanics have attracted considerable interest of both physicists and mathematicians in recent years. On the one hand, methods of quantum gravity have been successfully applied by D uplantier to obtain the multifractal spectrum of harm onic measure on critical cluster boundaries [1, 2]. On the other hand, the invention of the Schram m-Loew ner evolution (SLE) by Schram m [3] has made it possible to study conform ally-invariant critical curves rigorously (several reviews of SLE are available by now; see, for example, Refs. [4, 5, 6, 7, 8]).

Conform al eld theory (CFT ) has been a natural and traditional language for description of critical statistical systems in two dimensions. It is important to understand how fractal properties of critical curves can be obtained from CFT.A part of the emerging picture is that the stochastic geom etry of critical curves can be studied by traditional methods of CFT.As any local eld theory, CFT focuses on correlation functions of local operators. It turns out that the stochastic geom etry of critical curves is related to correlators of certain prim ary elds in CFT. This connection, rst discussed in Ref. [9], has been further developed in our papers [10, 11] where we have identi ed the curve-creating prim ary operators and considered their correlators with other prim aries serving as \probes" of harm onic m easure. A long this way we have reproduced D uplantier's results for multifractal exponents associated with harm onic m easure.

In addition to harm onic m easure, com plicated planar dom ains and their boundaries are characterized by winding or rotation. The notion of rotation spectrum for planar dom ains has been introduced by B inder in Ref. [12]. Then the mixed rotation and harm onic measure spectrum has been found exactly for critical curves by D uplantier and B inder [13]. In that paper the authors com bined an earlier C oulom b gas approach to the distribution of winding angles of critical curves [14] with the quantum gravity m ethods of Refs. [1, 2].

In this paper we extend our previous analysis [10, 11] to include the winding (rotation) of critical curves. As it happens, to account for rotation of critical curves, we need to consider CFT prim ary elds with com plex weights and charges. The procedure involves an analytic continuation of chiral correlation functions and then gluing the chiral sectors to obtain realbut angular dependent correlators. The paper is organized as follows. In Section 2 we form ulate the problem and give heuristic de nitions of various objects of interest including harm onic measure and rotation of critical curves. In Section 3 we consider a special case of a single probe of harm onic measure and rotation near a star con guration of several critical curves. Then we go on to the general case, where several probes can be placed in the vicinity of a critical star con guration. We conclude in Section 5 with a discussion and com parison with other works.

# 2 Description of the problem

C onsider a two-dimensional critical statistical system. Such systems can be formulated as critical points in an ensemble of curves on a lattice, e.g. the bops of the O (n) model or the high-T expansion bops of the Q -state Potts model. W hen the system is critical, these curves are called critical curves. In the continuum limit, statistical systems are invariant under conform al transform ations and can be described by a CFT with a certain central charge c. In this case the critical curves are uctuating fractal curves. Any conform al transform ation in two dimensions locally looks like a combination of rotation and a scale transform ation (dilatation). For a given composition of dilatation and rotation in the plane there exist curves which remain invariant. They are logarithm ic spirals = ln r. The quantity characterizes the winding of a spiral and is called the rate of rotation. For a pure dilatation the spirals degenerate into straight lines (no rotation: = 0) which may intersect and form corners with arbitrary opening angles.

Since any conform altransform ation is locally a composition of dilatation and rotation, a conform ally invariant critical curve can be thought of as an assembly of elementary corners and logarithm ic spirals at all scales. The so called multifractal analysis aim s at determ ining the fractal dimensions of the subsets of a critical curve with given opening angle and winding. These fractal dimensions form a continuous family of multifractal exponents.

Harm onic measure is another way of characterizing the complicated fractal geometry of a critical curve. In a simple electrostatic analogy we can imagine that a critical cluster is charged with a unit charge that all goes to the boundary of the cluster. Charge density (which is the density of harm onic measure) on the boundary is very uneven and lum py, and can be characterized by its moments. Therefore, in the ensemble of critical curves it is natural to characterize their fractal geometry by the statistics of local winding and harm onic measure near some point on the curve.

Locally, near a charged corner the scaling of the charge density or, equivalently, of the magnitude of the electric eld E is determined by the opening angle at the corner: rE (r) r = .W hat will be important for us in the following is that this power-law behavior is the same along all straight lines that converge at the apex of the corner. All these lines intersect the equipotentials at constant angles , and only one of them is an electric eld line (corresponding to = = 2). In mathematics such lines are sometimes called \slanted G reen lines", while the electric eld lines are called \G reen lines". In a general situation we will denote the slanted G reen line that goes from the origin and forms the angle with equipotentials by .

Sim ilar statements can be made for a logarithmic spiral. The information about the rate of rotation of a spiral is contained in the winding of any of the slanted G reen lines that emanate from the spiral's origin. It is easy to see that the line is the rotated logarithmic spiral =  $2 + \ln r$  (with = =2 again corresponding to the unique electric eld line going from the origin), and we can follow any of them to \m easure".

The information on the scaling of the magnitude of the electric eld and the winding of a slanted G reen line near a certain point on the critical curve is contained in the value of the derivative of a uniform izing conformal m ap w (z) that m aps the exterior of the critical cluster onto a standard sim ple dom ain. In these term s, the electrostatic potential of the curve is

log jw (z) j the magnitude of the eld near the curve is  $E = jw^{0}(z) j$  and the direction of the eld is related to  $\arg w^{0}(z)$ . (In terms of the complex potential (z) =  $\log w(z) we have^{-0}(z) = E_{x} + iE_{y}$ .) We will use this idea extensively in what follows. It is convenient for our purposes to restrict the statistical ensemble so that a critical curve always passes through a xed point, and work in the vicinity of this point.

In this paper we consider only dilute critical system s, which means that critical curves generically do not intersect (see, however, the discussion of exceptional star con gurations below). This choice is dictated by the nature of our problem. In the dense phase where critical curves have multiple points, the harm onic measure is only supported on the external perimeters of critical clusters (think of electric charge spreading on the surface of a conductor). The external perimeters are always simple critical curves described by a model in the dilute phase, and it is su cient to consider dilute system s in our problem. The prime example of this phenom enon is provided by critical percolation: percolation hulls intersect them selves at all scales, while the external perimeters of percolation clusters are simple curves in the universality class of self-avoiding random walks.

In the absence of boundaries in the system critical curves are closed loops. If the system has a boundary and the boundary condition changes at certain boundary points, it gives rise to critical curves which start and end at these points. A hole in the system is a component of the boundary. If the boundary condition changes on it, we will observe critical curves emanating from the hole. On scales much larger than the size of the hole the latter becomes a puncture. In this case there are critical curves coming out of a single point in the bulk of the system. Inserting a puncture at some point in the bulk we can ensure that in each realization of the statistical ensemble there is a  $star" \mid a$  xed number k of curves starting from this point.

Generally speaking, a critical system need not contain punctures or be restricted in any way by insertion of curve-creating operators. These are merely arti cial devices which we will use in our calculations. The meaning of k = 2 stars is obvious: any point on a curve is such a star. In fact, stars with k > 2 can also be meaningful even in the absence of punctures. A star with an even number k > 4 of legs is a point of self-intersection of a critical curve. Since in the dilute phase the latter are simple curves with probability one, stars with k > 4 can only appear as exceptional con gurations in rare realizations of the statistical ensemble, all such realizations having the total weight (probability) zero. In fact, we can quantify this discussion by

assigning a negative fractal dimension  $d_k$  (see Eq. (50) below) to the set of stars with k > 4 legs. The meaning of a negative fractal dimension is as follows. If in a system of size L several curves approach each other within a distance l, then on the scales much larger than l the con guration of the curves will look like a star. However, in a typical realization of the statistical ensemble, the number of points where such approach happens, scales as  $(L=1)^{d_k}$ , and goes to zero in the therm odynam ic lim it.

In the rest of the paper we apply methods of CFT to the problem of the mixed multifractal spectrum of harmonic measure and winding of critical curves. In this approach a k-legged star of critical curves is created by an insertion of a speci c curve-creating operator  $_{0;k=2}$  [9,10,11]. This operator can be realized as a vertex operator in the C oulom b gas form ulation of CFT, and has known holom orphic charge  $_{0;k=2}$  and weight  $h_{0;k=2}$ , see Eqs. (4, 5, 6) below. O ther prim ary operators inserted close to the curve-creating one can then serve as probes for the harmonic measure. This has been explained and used in our previous works [10, 11]. In the present paper we will see that to account for winding of critical curves, the probesm ust have com plex charges and conform all weights.

W e want to point out here that while harm onic measure can behave nontrivially and be probed both in the bulk and on a boundary of a critical system, the mixed spectrum that takes into account winding is only de ned in the bulk. Indeed, there is no way a critical curve can rotate around a point on a boundary. Therefore, our present analysis only deals with the bulk curve creating operators.

# 3 The case of a single point

For simplicity we start with the case of the electric eld (both the magnitude and the direction) measured at a single point and then generalize the method to multiple points.

### 3.1 De nition of multifractal exponents

Consider a k-legged star | k critical curves emanating from the origin in the bulk of the system . We are interested in the properties of the starm uch closer to the origin than the system boundaries. The shape of the legs of the star far from the origin is unimportant, which allow sus to trace each leg only up to a certain distance from the origin. Let us describe the shape of such a truncated star by an analytic function w (z) which maps the exterior S of the star onto the exterior of the unit disk C nD. The statistical ensemble



Figure 1: The conform altransform ation blows up the star into the shape of a disk, like a pu er sh. The origin of the star is mapped to several points  $u_i$ . The point z is chosen to lie on the dotted circle of a xed radius and so that w (z) lies on the same radial line as  $u_1$ .

of the shapes of the star de nest the statistical ensemble of w (z). C lose to the origin the k curves of the star divide the plane into k sectors. Approached from di erent sectors, the origin is mapped onto k points  $u_i$  on the unit circle:  $\lim_{z \to 0} w(z) = u_i$  for z in the i-th sector. To make w (z) unique we specify w (1) = 1 and w<sup>0</sup>(1) = <sup>1</sup> > 0, where is the conform all radius of the star. Its uctuations are unimportant. For each realization of the star we can choose the truncation radius so that = 1.

As we have explained above, the magnitude and the direction of the electric eld at a point z are determined by  $jw^{0}(z)j$  and  $argw^{0}(z)$  respectively. Both these quantities are random. Conform al invariance suggests that locally  $jw^{0}(z)j$  scales as a power of jzj and  $argw^{0}(z)$  behaves as  $\ln jzj$ . Therefore, we can study the joint moments of  $jw^{0}(z)j$  and  $e^{argw^{0}(z)}$  expecting them to scale as

$$E \dot{w}^{0}(z) f^{e^{p \operatorname{arg} w^{0}(z)}} \dot{z} j^{k} (n P^{p}); \qquad (1)$$

where E denotes average over the uctuating geometry of the star. This equation de nes the multifractal exponents  $_k$  (n;p), and our goal below is to obtain them using methods of CFT.

Let us now discuss a subtle point about Eq. (1). We can treat the average over the star shapes in two ways. The rst way is to specify the position of point z in such a way that for any star S this point is on the slanted G reen line in, say, the rst sector (the choice of the sector is arbitrary and imm aterial). Let us denote this choice by z (im plicitly keeping in m ind that the choice depends on the star shape). Then the average in the left hand side of Eq. (1) should be understood as

$$E \dot{w}^{0}(z) f^{e^{pargw^{0}(z)}} \dot{z} \dot{z} \dot{z}^{k}(n,p); \qquad (2)$$

(only the argument of z uctuates here, while  $j_z$  j is xed). The simplest such choice is to place z on the electric eld line  $_{=2}$ . Then in the w plane the image w (z) lies on a radial straight line:

$$\arg w (z_{=2}) = \arg u_1; \tag{3}$$

see Fig. 1 for illustration.

On the other hand, as we have argued in the previous section, the scaling of  $jv^{0}(z)j$  and  $e^{\arg w^{0}(z)}$  should be the sam e along any line . Therefore, we can average (integrate) over (and sum over the di erent sectors) in Eq. (2) while keeping jz jconstant. This procedure produces a manifestly rotationally invariant average. In this average we can distribute the contributions that come from star shapes which di er by a global rotation around the origin into sets, and then sum up all such sets. Due to isotropy of our critical system, all terms in a given set have the same statistical weight. Therefore, to calculate the average, we can choose any single term from each set and sum these up. We can choose these terms so that arg z is the same a for all of them, which gives the left hand side in Eq. (1) with a xed point z. This is the second way to interpret the average over the star shapes. A swe see, the two ways are equivalent and produce the same scaling behavior with the multifractal exponents  $_{k}(n;p)$ .

#### 3.2 CFT calculation of multifractal exponents

A critical system is characterized by the central charge cwhich we parametrize, as is usual in CFT, by the so called background charge  $_0$ :

$$c = 1 \quad 24 \quad {}_{0}^{2}$$
: (4)

A swe have shown in Ref. [11], in the C oulom b gas description of the dilute phase we have to impose the D irichlet conditions on the system boundaries. A consistent way of doing it also implies that the background charge  $_0$  in Eq. (4) is non-positive:  $_0$  6 0.

W ithin the CFT description (see [11, 16] for introduction to the m athem attical apparatus of CFT used below ) a k-legged star centered at the origin

is created by inserting the curve-creating operator  $_{0;k=2}$  (0) [10, 11]. This operator has holom orphic charge

$$_{0;k=2} = _{0} \frac{k}{4} _{0} \frac{p}{_{0}^{2}+1};$$
 (5)

and its antiholom orphic charge is  $_{0,k=2} = _{0,k=2}$ . It is spinless with real conform alweights:

$$h = h = h_{0;k=2} = 0;k=2 (0;k=2 2 0):$$
(6)

To calculate the m ixed multifractal exponents  $_k$  (n;p) by m eans of CFT we consider a prim ary operator O<sub>h;h</sub> whose holom orphic and antiholom orphic weights h;h are complex and conjugate to each other:

$$h = h$$
 : (7)

(Notice that we use asterisk to denote complex conjugation, while the bar over a weight or charge simply indicates the antiholom orphic sector.) Its holom orphic and antiholom orphic charges are related to the weights by the equations [11]

$$h = (2_0); \qquad h = (+2_0):$$
 (8)

(The introduction of the charges is convenient because they make fusion of primary operators look simple: the charges of the fused elds simply add up.) Note the sign di erence in the formula for h compared to that in Ref. [16]. The di erence is necessitated by our convention ( $_0 6 0$ ) for the dilute phase. The solutions (h), (h) are chosen so that (0) = (0) = 0:

$$= _{0} + \sqrt{\frac{2}{0} + h}; \qquad = _{0} \sqrt{\frac{2}{0} + h}: \qquad (9)$$

The two charges are related by = . If we denote the real and in aginary parts of the holom orphic charge as  $^{0}$  and  $^{0}$ , the real and in aginary parts of the holom orphic weight h can be written as

Reh = 
$${}^{02}$$
  ${}^{002}$  2  ${}_{0}$   ${}^{0}$ ; Im h = 2  ${}^{00}$  (  ${}^{0}$   ${}_{0}$ ): (10)

The operator  $O_{h;h}$  plays the role of a \probe" of harm onic measure and rotation of critical curves when we place it at a point z near the origin of a k-star. The star is created by inserting the curve-creating operator  $_{0;k=2}(0)$ . Thus we propose to consider the following correlation function:

$$D \qquad E \qquad 0_{jk=2} (0) O_{h,jh} (z) (1) / z^{2} O_{jk=2} ^{2} = jz \int_{0}^{4} O_{jk=2} ^{0} e^{4} O_{jk=2} ^{0} arg z : (11)$$

Here and in what follows these angular brackets denote a CFT average and

stands for all operators far from the origin and is necessary to make the correlation function non-zero. The right hand side in Eq. (11) was obtained in a standard way by fusion of prim ary operators (expressed as vertex operators in the C oulom b gas form ulation of C FT). Notice that the presence of complex weights and charges leads to an expression that is explicitly angular dependent and multivalued. Thus, this correlator does not correspond to any physical quantity, and we need to consider

$$C = e^{4} e^{4} e^{2} e^{0} arg z} e^{0} e^{2} e^{0} e^{2} e^{0} e^{2} e^{-0} e^$$

In this formulation the position of the point z is xed, and the extra exponential factor is a constant that can be pulled out of the average h:::i.

The same correlation function C, being an average over all degrees of freedom in the system, can be evaluated in another way, using the so-called two-step averaging procedure [7, 10, 11, 15]. We can rst x the shape of the star and sum over the rest of the degrees of freedom. Then we average over all possible shapes of the star:

$$C = E e^{4} e^{0;k=2} O_{h;h}(z) (1) = (13)$$

The subscript of a correlation function refers to the domain in which it is computed, in this case the exterior of the star S.

The point z is still xed in Eq. (13), therefore, for each shape S this point lies on a di erent slanted G reen line . However, as in Section 3.1, we can argue that the scaling with  $j_2 j_3$  is the same (independent of ) for all the term s with di erent values of . Then to nd the exponents  $_k (n;p)$  it is su cient to retain only term s with a xed (in each star con guration) value of and consider a di erent quantity

$$C = E e^{4_{0;k=2}} O_{h;h}(z) (1)_{S} :$$
(14)

There is a parallel between the correlator C and the expression in Eq. (1), and the correlator C and the expression in Eq. (2): though C and C are form ally very di erent objects, they scale with jzj in the same way. The crucial di erence between C and C is that in the latter object the argument of z is uctuating (it depends on the star con guration) and the factor  $e^{4}_{0,k=2}$  must be kept under the average E [:::].

The next step in transform ing the expression (14) is to map the exterior of the star S onto the exterior of the unit disk by w(z). Under this map

the operator O<sub>h,h</sub> transform s according to the de nition of a CFT primary eld, and operators at in nity do not transform due to the normalization of w (z). Then conform all covariance of correlators of CFT implies

$$C = E e^{4} e^{(a_{k=2})^{(0)} arg z} w^{(0)} (z)^{h} w^{(0)} (z)^{h} O_{h;h} (w (z)) (1) C_{nD} : (15)$$

In this correlator the value of the angle is imm aterial. For simplicity of notation let us choose it to be = =2, which means that the point  $z_{=2}$  is chosen according to Eq. (3) and lies on the unique electric eld line connected to the origin of the star. We will drop the subscript =2 in the subsequent equations.

In the w plane, argw (z) is a xed number no greater than 2. At the same time, when w (z) approaches  $u_1$ , its pre-image z moves along the electric eld line and exhibits a lot of winding. Since argw (z) argw<sup>0</sup>(z) + arg z, we can write

$$\arg z \qquad \arg w^{0}(z);$$
 (16)

because both angles here are large. Thus, we can replace arg z by  $\arg w^{0}(z)$  in Eq. (15). Using Eq. (10) we can also rewrite

$$w^{0}(z)^{h}w^{0}(z)^{h} = w^{0}(z)^{h^{2}} = jw^{0}(z)^{j^{2}Reh}e^{2\operatorname{Im} hargw^{0}(z)}$$
$$= jw^{0}(z)^{j^{(m^{2}-m^{2}2-0)}}e^{4-m^{(m^{2}-m^{2}2-0)}argw^{0}(z)}: (17)$$

The remaining CFT correlation function in the exterior of the disk is computed by fusion of the primary eld with the boundary (which means the fusion of the holom orphic part O<sub>h</sub> (w (z)) with its image across the boundary of the unit disk). A coording to Eq. (3), both w (z) and its image in the unit circle 1=w (z) have the same (constant) argument  $\arg u_1$ . Then we nd the following scaling behavior:

$$O_{h,h}(w(z))(1)_{CnD} / w(z) \frac{1}{w(z)}^{2} jw(z) u_{1}j^{2j}^{2}$$

$$jz j^{(w+wz)} jw^{0}(z) j^{(w+wz)}: (18)$$

At this stage the correlator C evaluated in two steps becomes

$$C \dot{z}^{2(0^{2}+0^{2})} E \dot{y}^{0}(z)^{4(0^{0}-0)} e^{4(0^{0}-0^{+}-0)} e^{2\operatorname{arg} w^{0}(z)} : (19)$$

Since, as we have argued, both C and C should scale in the same way with  $j_{2}j_{j}$  comparison with Eq. (12) gives

$$\begin{array}{c} h \\ E j w^{0}(z) j^{4} j^{0}(0) e^{4} e^{2(z)} e^{-(z)} e^$$

The last equation is identical to the de nition (1) provided we denote

$$n = 4 \ {}^{0}(\ {}^{0} \ {}_{0}); \qquad p = 4 \ {}^{00}(\ {}^{0} \ {}_{0} + {}_{0;k=2}); \qquad (21)$$

$$_{k}(n;p) = 2(^{(2)} + ^{(2)}) + 4^{(0)}_{0;k=2}$$
: (22)

As usual, we need to choose the solutions of Eqs. (21) that vanish for n = 0; p = 0:

$$2^{0} = {}_{0} + {}^{p} \frac{2}{0} + n; \qquad {}^{0} = \frac{p}{4({}^{0} {}_{0} + {}_{0;k=2})}: \qquad (23)$$

Substituting these values into Eq. (22) we not the mixed multificated exponents k(n;p). With the shorthand

$$p_n = 0 + \frac{p_{-2}}{0} + n$$
 (24)

the answer can be written as

$$_{k}(n;p) = 2_{0;k=2} n \frac{1}{2} \frac{2}{n} \frac{1}{8} \frac{p^{2}}{\frac{1}{2} n 0 + 0;k=2}^{2}$$
: (25)

W e can also rewrite this result as

$$_{k}(n;p) = _{k}(n) \frac{1}{4} \frac{p^{2}}{_{k}(n) + n + 2(_{0;k=2} _{0})^{2}};$$
 (26)

where

$$_{k}(n) = _{k}(n;0) = 2_{0;k=2} n \frac{1}{2} \frac{2}{n}$$
: (27)

# 4 The general case

To de neamulti-point generalization of the mixed spectrum (1) we consider the following average:

$$\begin{array}{cccc} h & & i \\ E & e^{p \arg w^{0}(z_{1})} & & j w^{0}(z_{1}) \\ & & j w^{0}(z_{1}) \\ & & i = 1 \end{array}$$
 (28)

Here all the points  $z_i$  have the sam e distance to the star origin:  $\dot{z}_i j = r$ , and no two of them lie in the sam e sector. Because the curves do not intersect, all the winding angles  $\arg w^0(z_i)$  di er by no more than 2, and since they are all very large, they must all scale in the sam e way:  $\arg w^0(z_i)$  arg  $w^0(z_1)$ .

Thus, the topology of the star leads to only one parameter p describing the rotation, while we have k parameters  $n_i$  for harm onic measure in each of the sectors between the star's legs.

Similar to the discussion in Section 3.1 the points  $z_i$  can be either xed or can be chosen to lie on speci c slanted G reen lines. These choices do not a ect the scaling with r in Eq. (28). For example, we can choose  $\arg z_i$  by the requirement

$$argw(z_i) = argu_i$$
: (29)

Then in all realizations of the star each  $z_i$  lies on the unique G reen line in the i-th sector that starts at the star's origin.

The calculation of  $_{k}$  (fn<sub>i</sub>g;p) is a straightforward generalization of the calculation of the previous section. We introduce k primary operators  $O_{h_{i},h_{i}}(z_{i})$  each with complex conjugate weights:  $h_{i} = h_{i}$  and holom orphic charges  $_{i} = _{i}^{0} + i _{i}^{0}$ . In close parallel to Eq. (12) we consider a CFT correlation function:

$$C_{k} = e^{4} e^{(0)(k+2)\sum_{i} (0)^{m} \arg z_{i} + 4\sum_{i < j} (0)^{m} \exp (z_{i} z_{j})} E_{0;k=2}(0) O_{h_{i};h_{i}}(z_{i}) (1) :$$
(30)

This correlation function can be computed by fusion of primary elds:

D

$$D \qquad Y^{k} \qquad E \qquad Y \qquad z_{1}^{2} \qquad z_{1}^{0;k=2 \ i \ 2} \qquad (z_{1} \qquad z_{j})^{2} \qquad (z_{1} \qquad (z_{1} \qquad z_{j})^{2} \qquad (z_{1} \qquad (z_{1} \qquad z_{j})^{2} \qquad (z_{1} \qquad (z_$$

The exponential factors in the de nition of  $C_k$  are chosen to cancel the angular dependence in the last equation. Since in our setup  $\dot{z}_i j = r$  and  $\dot{z}_i$   $z_j j$  r for all i and j, the correlation function  $C_k$  scales as

$$C_{k} = r^{4} {}_{0;k=2} \sum_{i} {}_{i}^{0} + 4 \sum_{i < j} ( {}_{i}^{0} {}_{j}^{0} {}_{i}^{0} {}_{j}^{0} ) ; \qquad (32)$$

On the other hand, we can consider the sim ilar correlator  $C_k$ ; where the points  $z_i$  are chosen according to Eq. (29) and, therefore, uctuate form con guration to con guration. The correlators  $C_k$  and  $C_k$ ; should scale as

the same powers of r. We can evaluate the correlation function  $C_k$ ; in two steps: rst we x the shape of the star and sum over the rest of the degrees of freedom. Near a xed star all the arguments  $\arg z_i$  and  $\arg (z_i \quad z_j)$  are large and dier by nomore than 2. Therefore we can replace them all by one of them, say  $\arg z_1$ . Then we average over all possible shapes of the star:

$$C_{k}; \quad E e^{4 \left( \begin{array}{c} 0; k=2 \\ \end{array} \right) \sum_{i} \begin{array}{c} 0 \\ i \\ \end{array} \right) + \sum_{i \in j} \begin{array}{c} 0 \\ i \\ \end{array} \right) \arg z_{1} \\ O_{h_{i}; h_{i}} (z_{i}) \\ S \end{array} (1) \\ S \qquad (33)$$

Now we map the exterior of the star S onto the exterior of the disc CnD and note that  $\arg z_1$  arg w<sup>0</sup>( $z_1$ ) (com pare with Eqs. (15, 16)):

The remaining correlator in C nD is computed by fusion of primary elds with the boundary (compare with Eq. (18)):

There are other factors in this correlator that are coming from the fusion of di erent bulk elds O  $_{h_i,h_i}$  (w (z<sub>i</sub>)) with each other. These cross-terms look like (w (z<sub>i</sub>) w (z<sub>j</sub>))<sup>2 i j<sup>2</sup></sup>, and do not contribute to the short-distance behavior, since all z<sub>i</sub> lie in di erent sectors of the star, and the di erence w (z<sub>i</sub>) w (z<sub>j</sub>) stays nite in the lim it z<sub>i</sub>! 0. As a result,

$$C_{k}; \quad r^{2\sum_{i} (\frac{\omega}{i} + \frac{\omega}{i})} E^{h} e^{p \arg w^{0}(z_{1})} \overset{Y}{j} \dot{w}^{0}(z_{i}) \overset{i}{J^{i}}; \quad (36)$$

where

We solve these equations for  ${}^{0}_{i}$  and  ${}^{P}_{i}$   ${}^{0}_{i}$  and choose the solutions which vanish at  $n_{i} = p = 0$ . Then Eqs. (32, 36) im ply Eq. (28) with

$${}_{k} (fn_{i}g;p) = \begin{pmatrix} X \\ 2 \\ 0;k=2 \\ n_{i} \end{pmatrix} \begin{pmatrix} 2 \\ n_{$$

where  $n_i = 0 + \frac{p_i}{0} + n_i$ . Similarly to Eq. (26), this result can be rewritten as

$${}_{k}(fn_{i}g;p) = {}_{k}(fn_{i}g) - \frac{1}{4} - \frac{p}{k} \frac{p^{2}}{(fn_{i}g) + \frac{p}{i}n_{i} + 2(0,k=2)};$$
(39)

If all  $n_i$  vanish except one, the form ulas (38, 39) reduce to (25, 26).

## 5 D iscussion

The exponents  $_{k}$  (n;p) and  $_{k}$  (fng;p) that we have obtained in Eqs. (25, 38) are most natural from the point of view of CFT and correlation functions of prim ary operators. They are easily related to otherm ultifractal exponents de ned directly in terms of harm onic measure and winding. To exhibit this relation we start with a special case.

Consider a closed critical curve. Let us cover it by circles of radius r with centers at  $_i$ . Let us denote the harm onic measure within each circle by H ( $_i$ ;r), and the winding angle of the G reen line ending at  $_i$  at distance r away from  $_i$  by ( $_i$ ;r). Then we can de ne a sort of \global" mixed spectrum (n;p) by (see Eq. (2) in Ref. [13])

$$\overset{h\chi}{\underset{i}{\overset{}}} H^{n}(_{i};r) e^{p_{(i};r)} / r^{(n_{f}p)}:$$
(41)

If we set n = p = 0 in the above equation, we simply get the number of circles of radius r needed to cover the curve. This number should scale as  $r^{d_f}$ , where  $d_f$  is the fractal dimension of the critical curve. Therefore,  $(0;0) = d_f$ .

It is natural to assume some sort of ergodicity, so that the average of the sum in Eq. (41) can be replaced by the local average E [H  $^{n}e^{p}$  ] multiplied by the number of terms in the sum. If the local average scales as  $r^{\sim (n,p)}$ , then the global and the local exponents are related by

$$(n;p) = \sim (n;p) \quad d_{\pm}:$$
 (42)

The harm onic measure H ( $_i$ ;r) by electrostatic analogy should locally scale as rjw<sup>0</sup>( $z_i$ ) j where  $z_i$  is a point that lies not on the curve at distance r away from  $_i$ , compare with Fig. 1. In the situation we describe here each point on the curve can be viewed as the origin of a 2-legged star. Then

we can relate our exponent  $_2(n;p)$  to the local and the global exponents  $\sim (n;p)$ , (n;p):

$$\sim (n;p) = _2(n;p) + n;$$
  $(n;p) = _2(n;p) + n d_f:$  (43)

The fractal dimension  $d_f$  of the curve is related to the dimension of the curve creating operator  $_{0;1}$  [17]:

$$d_f = 2 \quad 2h_{0;1}$$
: (44)

F inally, we get the following relation

$$(n;p) = _{2}(n;p) + n + 2h_{0;1}$$
 2: (45)

Substituting here the expressions (6, 26) with k = 2 we get

$$(n;p) = (n) \frac{1}{4} \frac{p^2}{(n) + b};$$
 (46)

where

$$(n) = {}_{2}(n) + n + 2h_{0,1} + \frac{2}{2} = \frac{1}{2} p_{-\frac{2}{0}+n} + p_{-\frac{2}{0}+1} + \frac{2}{2} ({}_{0}^{2}+1) = \frac{n}{2} + \frac{25}{24} r_{-\frac{24n+1}{25}c} + \frac{24n+1}{25} r_{-\frac{25}{25}c} + \frac{1}{2} (47)$$

is the multifractal spectrum of the harm onic measure (see Eq. (6.32) in Ref. [2]), and

b = 2(
$$_{0;1}$$
  $_{0}$ )<sup>2</sup> + 2 2h<sub>0;1</sub> = 2( $_{0}^{2}$  + 1) =  $\frac{25 \text{ c}}{12}$ : (48)

Form ula (46) is exactly the same as Eq. (14) for (n;p) in Ref. [13]. Generalized exponents  $_{k}$  (fn<sub>i</sub>g;p) can be de ned sim ilarly to Eq. (41):

$$\begin{array}{cccc}
h & \chi & \gamma^{k} & i \\
E & e^{p & (z_{1};r)} & H^{n_{i}}(z_{i};r) & / r^{k} & (fn_{i}g;p) \\
k & stars & i=1
\end{array}$$
(49)

In this de nition the points  $z_i$  are the same as in Eq. (28), and H ( $z_i$ ;r) denotes harm onic measure on the portion of the star inside the i-th sector up to radius r. As we have argued in Section 2, k-legged stars with k > 4 m ay appear spontaneously in a critical system, even though in a subset of the realization of the statistical ensemble with the total measure zero. The

origins of the k-legged stars have a certain fractal dimension  $d_k$ . In full analogy with Eq. (44), the fractal dimension  $d_k$  is related to the scaling dimension  $2h_{0,k=2}$  of the operator  $_{0,k=2}$ :

$$d_k = 2 \quad 2h_{0;k=2}$$
: (50)

Notice that, consistent with the discussion in Section 2,  $d_k$  is negative for  $k > 4\,.$ 

A ssum ing ergodicity and the scaling H  $(z_i;r)$   $rjv^0(z_i)j$  as before, we can relate the generalized exponents  $_k$  (fn<sub>i</sub>g;p) to  $_k$  (fn<sub>i</sub>g;p):

$$_{k}(fn_{i}g;p) = _{k}(fn_{i}g;p) + _{n_{i}}^{X} n_{i} + 2h_{0;k=2} 2:$$
 (51)

U sing Eqs. (6, 39) we can express this result as

$$_{k}(fn_{i}g;p) = _{k}(fn_{i}g) - \frac{1}{4} \frac{p^{2}}{_{k}(fn_{i}g) + b};$$
 (52)

where

are the \higher multifractal exponents" of Ref. [2], Section 7.3.1. (To convince oneself of the equivalence, one needs to substitute 2  $_0 = (1 )^{1=2}$ , where 6 0 is the so-called string susceptibility exponent.)

Expressions (46, 47, 52, 53) and their meaning and consequences are analyzed in detail in Ref. [2]. Here we only mention that the Legendre transform softhese exponents (the so called singularity spectra) have a direct geom etrical meaning of fractal dimensions of subsets of points with a given local scaling of harm onic measure and winding.

In conclusion, we have shown how a detailed description of stochastic geom etry of critical conform ally-invariant curves in terms of their harm onic m easure and winding can be obtained by m eans of CFT, from correlation functions of certain prim ary operators. The inclusion of winding (rotations) m akes it necessary to consider operators with complex weights.

## 6 A cknow ledgem ents

This research was supported in part by NSF MRSEC Program under DMR-0213745, NSF Career Award under DMR-0448820, NSF grant under DMR-0645461 (AB), and the U S.D epartm ent of Energy grant DE-FG 02-ER 46311 (IR). We wish to acknow ledge helpful discussions with I. Binder and P. Wiegmann. While preparing this paper for publication, we have learned about a related work by B.D uplantier and I.Binder.

## References

- B. Duplantier, Conformally invariant fractals and potential theory, Phys. Rev. Lett. 84, 1363 (2000).
- [2] B. Duplantier, Conform al fractal geometry and boundary quantum gravity, in Fractal geometry and applications: a jubile of Benot M andelbrot, Part 2, 365, Proc. Sym pos. Pure M ath., 72, Part 2, AM S, 2004; arX iv: m ath-ph/0303034.
- [3] O. Schramm, Scaling limits of loop-erased random walks and uniform spanning trees, Israel J. Math. 118, 221 (2000); arXiv: mathPR/9904022.
- [4] G.F.Lawler, Conform ally invariant processes in the plane. M athem atical Surveys and M onographs, 114. Am erican M athem atical Society, P rovidence, R I, 2005.
- [5] W .Kager, B.Nienhuis, A guide to stochastic loew ner evolution and its applications, J.Stat. Phys. 115, 1149 (2004); arX iv: m ath-ph/0312056.
- [6] J. Cardy, SLE for theoretical physicists, Ann. Phys. 318, 81 (2005); arX iv: cond-m at/0503313.
- [7] M. Bauer and D. Bernard, 2D growth processes: SLE and Loewner chains, Phys. Rep. 432, 115 (2006); arX iv: m ath-ph/0602049.
- [8] I. A. Gruzberg, Stochastic geometry of critical curves, Schramm-Loewner evolutions, and conform al eld theory, J.Phys.A:Math.Gen. 39, 12601 (2006); arXiv:math-ph/0607046.
- [9] M. Bauer and D. Bernard, Conformal eld theories of stochastic Loewner evolutions, Commun. Math. Phys. 239, 493 (2003); arX iv: hep-th/0210015.

- [10] E.Bettelheim, I.Rushkin, I.A.Gruzberg, and P.W iegm ann, On harmonic measure of critical curves, Phys. Rev. Lett. 95, 170602 (2005); arXiv: hep-th/0507115.
- [11] I.Rushkin, E.Bettelheim, I.A.G nuzberg, P.W iegm ann, Critical curves in conform ally invariant statistical system s, J. Phys. A: M ath. Theor. 40, 2165 (2007); arX iv: cond-m at/0610550.
- [12] I.A.Binder, PhD.thesis, Caltech, 1998. Also see \Ham onic measure and rotation of simply connected planar domains", preprint available from

http://www.math.toronto.edu/ilia/Research/index.html

- [13] B. Duplantier and I. A. Binder, Harmonic measure and winding of conform ally invariant curves, Phys. Rev. Lett. 89, 264101 (2002); arX iv: cond-m at/0208045.
- [14] B. Duplantier and H. Saleur, W inding-angle distributions of twodimensional self-avoiding walks from conformal invariance, Phys. Rev. Lett. 60, 2343 (1988).
- [15] M. Bauer, D. Bernard, and J. Houdayer, D ipolar stochastic Loewner evolutions, J. Stat. Mech. P 03001 (2005); arX iv: m ath-ph/0411038.
- [16] P.DiFrancesco, P.M athieu, and D. Senechal, Conformal eld theory (Springer, 1999).
- [17] M.Bauer and D.Bernard, SLE, CFT and zig-zag probabilities, arX iv: m ath-ph/0401019v1.