

Particle multiplicities in nuclear collisions at the LHC from saturation physics

Javier L. Albacete

The Ohio State University. Department of Physics.
191 W. Woodruff Ave, Columbus OH-43210, USA

E-mail: albacete@mps.ohio-state.edu

Abstract. The inclusion of running coupling effects in the BK-JIMWLK evolution equations considerably reduces the energy dependence of the saturation scale with respect to previous estimates based on fits to HERA data. We discuss how such slowdown affects the expectations for particle multiplicities in Pb-Pb collisions at LHC energies. Our prediction is based on the use of k_t -factorization and on the use of unintegrated gluon distributions taken from the numerical solutions of the BK equation with running coupling. We obtain a central value $dN^{ch}/d\eta(\sqrt{s} = 5.5 \text{ TeV})|_{\eta=0} \approx 1390$.

PACS numbers: 21.65 Qr, 12.38 Mh

The experimental results from RHIC [1] strongly suggest that heavy ion collisions at high energies probe QCD in the non-linear regime characterized by strong coherent fields and gluon saturation, the Color Glass Condensate (CGC) [2]. Thus, many bulk features of multiparticle production in RHIC collisions, such as the energy, rapidity and centrality dependence of multiparticle production, are successfully described by models based in CGC physics [4, 5]. With collision energies of up to 5.5 TeV, the upcoming program in lead-lead collisions at the CERN Large Hadron Collider (LHC) is expected to provide confirmation for the tentative conclusions reached at RHIC and to discriminate between the different physical mechanisms proposed to explain particle production in high energy nuclear reactions (a review of predictions for heavy ion collisions at the LHC based on alternative approaches can be found in [3]).

The phenomenological models in [4, 5] rely on the assumption that the saturation scale Q_{sA} that governs the onset of non-linear effects in the wave function of the colliding nuclei is perturbatively large $\sim 1 \text{ GeV}$ at the highest RHIC energies. Next, gluon production is calculated via the convolution of the nuclear unintegrated gluon distributions (ugd's) according to k_t -factorization [6]. Under the additional assumption of local parton-hadron duality, the pseudorapidity density of charged particles produced in a nucleus-nucleus collisions can be written as follows

$$\frac{dN_{ch}}{dy d^2b} = C \frac{4\pi N_c}{N_c^2 - 1} \int \frac{d^2p_t}{p_t^2} \int d^2k_t \alpha_s(Q) \varphi\left(x_1, \frac{|k_t + p_t|}{2}\right) \varphi\left(x_2, \frac{|k_t - p_t|}{2}\right), \quad (1)$$

where φ denote the ugd's of projectile and target respectively, p_t and y are the transverse momentum and rapidity of the produced particle, $x_{1,2} = (p_t/\sqrt{s}) e^{\pm y}$, $Q =$

$0.5 \max\{|p_t \pm k_t|\}$ and b the impact parameter of the collision. The lack of impact parameter integration in this calculation and the gluon to charged hadron ratio are accounted for by the constant C , which sets the normalization.

The second basic ingredient of CGC models is the one of saturation of the nuclear ugd's entering Eq. (1). Using a relatively simple ansatz for the nuclear ugd's, and for symmetric collisions, the midrapidity multiplicity rises proportional to the nuclear saturation scale, which is assumed to grow as a power of the collision energy, $\sqrt{s}$:

$$\left. \frac{dN_{ch}}{dy d^2b} \right|_{y=0} \propto Q_{sA}^2 \approx const \cdot \sqrt{s}^\lambda. \quad (2)$$

So far, the energy dependence of the saturation scale has been adjusted to the empirical value extracted from fits to DIS HERA data of saturation-based models [7], which yields $\lambda \approx 0.288$. This has been largely motivated by the inability of the leading-log BK-JIMWLK evolution equations to reproduce experimental data. Here discuss how the recent advances in the calculation of running coupling corrections to the BK-JIMWLK evolution equations [8] allow to compute the energy evolution of the ugd's from first principles, getting a good agreement with experimental data.

We start by solving the non-linear small- x evolution equation for the dipole-nucleus scattering matrix, $S(Y, r)$, including running coupling corrections [8–10]:

$$\frac{\partial S(Y, r)}{\partial Y} = \mathcal{R}[S] - \mathcal{S}[S], \quad (3)$$

where r is the dipole size and $Y = \ln(x_0/x)$. Explicit expressions for the kernel terms in Eq. (3) as well as a detailed explanation of the numerical method used to solve Eq. (3) are given in [9]. The initial conditions for the evolution are taken from the McLerran-Venugopalan model [11]. We classify them according to their initial value of the saturation scale, Q_0 . Importantly, the saturation scale extracted from solutions of Eq. (3) grows with energy significantly more slowly than the value extracted from empirical parametrizations of DIS data, as shown in Fig. 1. On the other hand, the evolution speed $\lambda = d \ln Q_s^2 / dY$ turns out to be a function of Y , rather than a constant.

We now use the exact solutions of Eq. (3) to calculate particle multiplicities according to Eq. (1). Thus, the nuclear ugd's are now given in terms of the dipole scattering matrix evolved according to Eq. (3):

$$\varphi(Y, k) = \int \frac{d^2r}{2\pi r^2} \exp\{i \underline{r} \cdot \underline{k}\} (1 - S(Y, r)), \quad (4)$$

The relation between the evolution variable in Eq. (3) and Feynman- x of the produced particle is taken to be $Y = \ln(0.05/x_{1,2}) + \Delta Y_{ev}$. Since the relevant values of Bjorken- x probed at mid-rapidities and $\sqrt{s_{NN}} = 130$ GeV at RHIC are estimated to be $\sim 0.1 \div 0.01$, the free parameter ΔY_{ev} controls the extent of evolution undergone by the nuclear gluon densities resulting of Eq. (3) prior to comparison with RHIC data. Similar to [4], large- x effects have been modelled by replacing $\varphi(x, k) \rightarrow \varphi(x, k)(1 - x)^4$. The running of the strong coupling in Eq. (1), evaluated according to the one loop QCD expression, is




Figure 1. Left: Saturation scale as a function of rapidity corresponding to solutions of Eq. (3) (solid line) and setting $\lambda = 0.288$ (dashed line). Right: Evolution speed, $\lambda = \frac{d \ln Q_s^2(Y)}{dY}$, corresponding to the plot in the left. In both cases the initial saturation scale is $Q_0 = 1$ GeV.

regularized in the infrared by freezing it to a constant value $\alpha_{fr} = 1$ at small momenta. Finally, in order to compare Eq. (1) with experimental data it is necessary to correct the difference between rapidity, y , and the experimentally measured pseudo-rapidity, η . This is achieved by introducing an average hadron mass, m . Remarkably, the optimal value found in comparison with data, $m \sim 0.25$ GeV is in good quantitative agreement with the hadrochemical composition of particle production at RHIC.

With this set up we find a remarkably good agreement with the pseudo-rapidity densities of charged particles measured in 0–6% central Au+Au collisions at collision energies $\sqrt{s_{NN}} = 130$ and 200 GeV, as shown in Fig. 2. The comparison with data [1] constrains the free parameters of the calculation to the ranges: $Q_0 \sim 0.75 \div 1.25$ GeV, $m \sim 0.25$ GeV and $3 \gtrsim \Delta Y_{ev} \gtrsim 0.5$. Finally, the normalization constant is $C \sim \mathcal{O}(1)$ in all cases. The smallness of ΔY_{ev} indicates that the nuclear udg's probed at RHIC are in the *pre-asymptotic* regime [12]. With all the free parameters now constrained by comparison to RHIC data, the extrapolation to central Pb-Pb collisions at LHC energies, $\sqrt{s} = 5.5$ TeV per nucleon, is straightforward and completely driven by the non-linear small-x dynamics. We get

$$\frac{dN_{ch}^{Pb-Pb}}{d\eta}(\sqrt{s_{NN}} = 5.5 \text{ TeV}, \eta = 0) \sim 1290 \div 1480, \quad (5)$$

with a central value corresponding to the best fits to RHIC data ~ 1390 . These values are significantly smaller than those of other saturation based calculations [4, 5, 13], $\sim 1700 \div 2500$, and compatible with the ones based on studies of the fragmentation region [14], which yield $\approx 1000 \div 1400$.

This research is sponsored in part by the U.S. Department of Energy under Grant




Figure 2. Pseudo-rapidity density of charged particles produced in Au-Au 0-6% central collisions at $\sqrt{s_{NN}} = 130$ and 200 GeV and for Pb-Pb central collisions at $\sqrt{s_{NN}} = 5.5$ TeV. Data taken from [1]. The upper, central (solid lines) and lower limits of the theoretical uncertainty band correspond to $(Q_0 = 1 \text{ GeV}, \Delta Y = 1)$, $(Q_0 = 0.75 \text{ GeV}, \Delta Y = 3)$ and $(Q_0 = 1.25 \text{ GeV}, \Delta Y = 0.5)$ respectively, with $m = 0.25 \text{ GeV}$ in all cases.

No. DE-FG02-05ER41377 and by an allocation of computing time from the Ohio Supercomputer Center.

References

- [1] I. Arsene *et al.* [BRAHMS Coll.], Nucl. Phys. A **757**, 1 (2005); J. Adams *et al.* [STAR Coll.], Nucl. Phys. A **757**, 102 (2005); K. Adcox *et al.* [PHENIX Coll.], Nucl. Phys. A **757**, 184 (2005);
- [2] H. Weigert, Prog. Part. Nucl. Phys. **55** (2005) 461. E. Iancu and R. Venugopalan, arXiv:hep-ph/0303204.
- [3] N. Armesto *et al.*, J. Phys. G **35**, 054001 (2008) [arXiv:0711.0974 [hep-ph]].
- [4] D. Kharzeev, E. Levin and M. Nardi, Nucl. Phys. A **747**, 609 (2005).
- [5] N. Armesto, C. A. Salgado and U. A. Wiedemann, Phys. Rev. Lett. **94**, 022002 (2005).
- [6] L. V. Gribov, E. M. Levin and M. G. Ryskin, Phys. Rept. **100**, 1 (1983).
- [7] K. Golec-Biernat and M. Wüsthoff, Phys. Rev. D **59**, 014017 (1998).
- [8] I. I. Balitsky, Phys. Rev. D **75** (2007) 014001. Y. Kovchegov and H. Weigert, Nucl. Phys. A **784** (2007) 188–226. E. Gardi, J. Kuokkanen, K. Rummukainen, and H. Weigert, Nucl. Phys. A **784** (2007) 282–340.
- [9] J. L. Albacete and Y. V. Kovchegov, Phys. Rev. D **75**, 125021 (2007).
- [10] J. L. Albacete, Phys. Rev. Lett. **99**, 262301 (2007) [arXiv:0707.2545 [hep-ph]].
- [11] L. D. McLerran and R. Venugopalan, Phys. Rev. D **49**, 2233 (1994).
- [12] J. L. Albacete, N. Armesto, J. G. Milhano, C. A. Salgado and U. A. Wiedemann, Phys. Rev. D **71**, 014003 (2005).
- [13] K. J. Eskola, K. Kajantie, P. V. Ruuskanen and K. Tuominen, Nucl. Phys. B **570**, 379 (2000).
- [14] F. Gelis, A. M. Stasto and R. Venugopalan, Eur. Phys. J. C **48**, 489 (2006).