

Generalized Elliott-Yafet theory of electron spin relaxation in metals: the origin of the anomalous electron spin life-time in MgB_2

F. Simon,¹ B. Dora,^{1,2} F. Murany,¹ A. Janossy,¹ S. Garaj,³
L. Forró,³ S. Bud'ko,⁴ C. Petrovic,⁴ and P. C. Canfield⁴

¹Budapest University of Technology and Economics,

Institute of Physics and Condensed Matter Research Group of the Hungarian Academy of Sciences, H-1521 Budapest, Hungary

²Max-Planck-Institut für Physik Komplexer Systeme, Nothnitzer Str. 38, 01187 Dresden, Germany

³Institute of Physics of Complex Matter, FBS Swiss Federal Institute of Technology (EPFL), CH-1015 Lausanne, Switzerland

⁴Ames Laboratory, U.S. Department of Energy and Department of Physics and Astronomy, Iowa State University, Ames, Iowa 50011, USA

The temperature dependence of the electron spin relaxation time in MgB_2 is anomalous as it does not follow the temperature dependence of the resistivity above 150 K, it has a maximum around 400 K, and it decreases for higher temperatures. This violates the well established Elliott-Yafet theory of electron spin relaxation in metals. We show that the anomaly occurs when the quasi-particle scattering rate (in energy units) becomes comparable to the energy difference between the conduction- and a neighboring band. We find that the anomalous behavior is related to the unique band structure of MgB_2 and the large electron-phonon coupling. The saturating spin-lattice relaxation can be regarded as the spin transport analogue of the Loewen-Rohlfing criterion of electron transport.

PACS numbers: 74.70.Ad, 74.25.Nf, 76.30.Pk, 74.25.Ha

Knowledge of the electron spin-lattice relaxation time, T_1 , of conduction electrons plays a central role in assessing the applicability of metals for information processing using electron spins, spintronics [1]. T_1 is the time it takes for the conduction electron spin ensemble to relax to its thermalequilibrium magnetization after a non-equilibrium magnetization has been induced e.g. by conduction electron-spin resonance (CESR) excitation [2] or by a spin-polarized current [1]. The Elliott-Yafet (EY) theory of T_1 in metals [3, 4] has been well established in the past 50 years on various systems such as elemental metals [5], strongly correlated one-dimensional [6], and some of the alkali fulleride salt [7] metals. It is based on the fact that the spin part of the conduction electron wave functions is not a pure Zeeman state but is an admixture of the spin up and down states due to spin-orbit (SO) coupling. As a result, momentum scattering due to phonons or impurities induces electron spin-flip, which leads to spin relaxation. Typically every millionth momentum scattering is accompanied by the electron spin-flip due to the relative weakness of the SO coupling. Thus, T_1 (being the momentum relaxation time) which explains the motivation behind the efforts devoted to the spintronics applications of metals.

A consequence of the EY theory is the so-called Elliott-relation, i.e. a proportionality between T_1 and β :

$$\frac{1}{T_1} = \frac{L}{E}^2 \frac{1}{\beta} \quad (1)$$

Here β is a band structure dependent constant and for most elemental metals $\beta \approx 1:10$ (Ref. [5]). L is the SO splitting for spin up and down electrons in a valence

(or unoccupied) band near the conduction band with an energy separation of E . E.g. in sodium, the conduction band is 3s derived and the relevant SO state is the 2p with $E = 30.6$ eV and $L = 0.16$ eV giving $(L/E)^2 = 2.7 \cdot 10^{-4}$ [4].

The Elliott-relation shows that the temperature dependent resistivity and CESR line-width are proportional, the two being proportional to the inverse of β and T_1 , respectively. This enabled to test experimentally its validity for the above mentioned range of metals. Much as the Elliott-relation has been confirmed, it is violated in MgB_2 as therein the CESR line-width and the resistivity are not proportional above 150 K [8].

Here, we study this anomaly using MgB_2 samples with different B isotopes and impurity concentrations and we show that the anomalous effect is indeed intrinsic to MgB_2 . We explain the anomaly with an exact treatment of the SO scattering of conduction electrons in the presence of a nearby band with energy separation E , by extending the Mori-Kawasaki formula developed for localized spins to itinerant electrons. The result shows that the Elliott-relation breaks down when E is comparable to $\sim \hbar \omega_D$. Adrian deduced a similar result with a qualitative argument [9].

The role of E is disregarded in the EY theory since typical values are $E \approx 10$ eV and $\hbar \omega_D \approx 2 k_B T \approx 6$ meV at $T = 100$ K and $\lambda = 0.1$ electron-phonon coupling. We show that the occurrence of the anomaly in MgB_2 is related to the unique features in its band structure and the large electron-phonon coupling.

We performed CESR measurements on three kinds of ne powder MgB_2 with isotope pure ^{10}B , ^{11}B , and natural boron (20 % ^{10}B and 80 % ^{11}B). The samples have




FIG. 1: Comparison of the temperature dependent CEsR line-width ($\circ$: Mg^{11}B_2 , $\bullet$: MgB_2 of natural boron) and the resistance (solid curve) for Mg^{11}B_2 . The two types of data overlap in the 40-150 K temperature range. A representative error bar is shown.

slightly different impurity content, shown by the varying residual CEsR line-width, B_0 . The temperature dependent T_1 and the CEsR line-width, B , are related:

$B = B_0 + 1 = T_1$, where $\gamma = 28 \text{ GHz/K}$ is the electron gyromagnetic factor. ESR spectroscopy was done on a Bruker X-band spectrometer (center field 0.33 T) in the 4-700 K temperature range on samples sealed under He in quartz tubes. The most important result of the current report, the anomalous temperature dependence of B or T_1 , is independent of sample morphology, isotope content, or thermal history. B is also independent of the magnetic field, apart from a small change in B_0 [10]. Resistance on pellet samples and SQUID magnetometry were studied on the same batch as those used for ESR. The $\text{RRR} > 20$ and the sharp ($< 0.5 \text{ K}$) superconducting transition attest the high quality of the samples. Heating the samples in the ESR measurement (about 1 h duration) to 700 K does not affect the superconducting properties as shown by magnetization measurements.

We reported previously the anomalous temperature dependence of the CEsR line-width in Mg^{11}B_2 : although the line-width follows the resistance for the 40-150 K temperature range, it deviates above 150 K and saturates above 400 K [8]. This was confirmed independently [11, 12]. To our knowledge, this is the only known metal where such phenomenon is observed. We extended the previous measurement to 700 K and the result is shown in Fig. 1. Interestingly, the CEsR line-width does not just saturate at high temperatures, as found previously, but decreases slightly above 500 K. The result is reversible upon cooling with no dependence on the thermal treatment protocol. The phenomenon is reproduced on several

samples of different purity and boron isotopes, thus it is intrinsic to MgB_2 .

We explain the anomalous temperature dependence of T_1 in general before including the specifics of MgB_2 . The Elliott-Yafet theory disregards the magnitude of γ and takes lifetime effects only to lowest order into account [3, 4]. The extended description involves the Kubo-formalism and is based on a two-band model Hamiltonian, $H = H_0 + H_{SO}$, where:

$$H_0 = \sum_{k; s} [\epsilon(k) + \frac{1}{2} B s] c_{k; s}^\dagger c_{k; s} + H_{\text{scatt}}; \quad (2)$$

$$H_{SO} = \sum_{k; s; s^0} L_{s; s^0}(k) c_{k; s}^\dagger c_{k; s^0}$$

Here $s, s^0 = 1$ or 2 are the band, s, s^0 are spin indices, $L_{s; s^0}$ is the SO coupling, and B is the magnetic field along the z direction. H_{scatt} is responsible for the finite T_1 . The SO coupling does not split spin up and down states in the same band for a crystal with inversion symmetry, however it mixes different spin states in the two bands [1]. The Hamiltonian in Eq. 2 is essentially the same as that considered by Elliott [3]. However, instead of a time-dependent perturbation treatment, we calculate T_1 from the Mori-Kawasaki formula [13, 14]:

$$\frac{1}{T_1} = \frac{1}{2 \chi_0 B} \text{Im} G_{PP}^R(\omega_L); \quad (3)$$

where χ_0 is the static magnetic susceptibility, $\omega_L = B$ is the Larmor frequency, and $G_{PP}^R(\omega)$ is the Fourier transform of

$$G_{PP}^R(t) = -i \langle [P(t) - \langle P \rangle, P(0)] \rangle_{H_0}; \quad (4)$$

$$P = [H_{SO}; S^z];$$

The expectation value in Eq. 4. is evaluated with the unperturbed Hamiltonian, H_0 .

Assuming that the two bands are separated by $E(k) = \epsilon_1(k) - \epsilon_2(k) = \gamma(k)$, a standard calculation yields [15]:

$$\frac{1}{T_1} = \frac{L_z^2(k_F) + 2 \langle J_{\#} \rangle^2}{\gamma^2} \frac{1}{1 + (\gamma(k_F)/\gamma)^2}; \quad (5)$$

where the $\langle \cdot \rangle$ means Fermi surface averaging, $L_z(k) = L_{\#}(k) - L_{\#}(k)$, and we neglected L_z , which is small compared to $\gamma(k_F)$. Eq. 5. was previously deduced by Adrian using a qualitative argument, which involved an effective magnetic field, $L = \gamma$, fluctuating with correlation time due to the SO coupling [9].

We approximate Eq. 5 using effective values for the band-band energy separation and the SO coupling:




FIG. 2: (Color online) Schematics of the spin-lattice relaxation in MgB_2 in the two-band model framework. The arrow thicknesses represent the relaxation rates (not to scale). Note that the inter-band momentum scattering rate is larger than the spin-lattice relaxation rates, therefore there is a spin transfer between the two types of bands.

$$\frac{1}{T_1} = \frac{L_e^2}{\sim^2} \frac{1}{1 + \frac{1}{2} \frac{1}{\tau_e^2}}; \quad (6)$$

This result returns the Elliott-relation when $\frac{1}{\tau_e} \rightarrow 1$ and gives a decreasing spin relaxation rate with increasing $\frac{1}{\tau_e}$ when $\frac{1}{\tau_e} \rightarrow 1$, thus it can be regarded as a generalization of the Elliott-Yafet theory. In the following, we show that it describes the spin relaxation in MgB_2 .

Electronic properties of MgB_2 are described by the so-called two-band model meaning that the conduction bands related to the boron π bands have different electron-phonon couplings, different affinity to defects, and that the inter-band momentum scattering is weaker than the intra-band ones [16]. As a result, the conductivity is given by a parallel resistor formula [16], i.e. the band with longer dominates the transport. In contrast, the CESR spin relaxation is dominated by the band with shorter T_1 . Although the inter-band momentum scattering time, $\tau_{\sigma\pi}$, is longer than the intra-band momentum scattering times, τ_σ and τ_π , it is still much shorter than T_1 . Thus an electron with a given spin state is scattered back and forth between the two types of bands several times before flipping its spin, which is depicted in Fig. 2. The overall $1/T_1$ is the average of the spin-lattice relaxation rates weighted by the relative DOS on the σ and π bands, $N_\sigma = 0.56$ and $N_\pi = 0.44$ [17]:

$$\frac{1}{T_1} = \frac{N_\sigma}{T_{1,\sigma}} + \frac{N_\pi}{T_{1,\pi}}; \quad (7)$$

In Fig. 3, we show the band structure of MgB_2 from Refs. [18, 19] near the Fermi energy. Two boron π and




FIG. 3: (Color online). Band structure of MgB_2 near the Fermi energy after Refs. [18, 19]. Two of the bands (black) cross the Fermi surface near each other in the vicinity of the Γ and A points, whereas π bands (red) are separated from other bands with a larger optical gap at the crossing. We also show the dispersion with 8 times larger wave-vector resolution around the Γ points with vertical arrows for possible E values.

two π bands cross the Fermi energy such that the π bands are well separated from other bands with $E \sim 2$ eV whereas the two σ bands are close to each other and $E \sim 0.2$ eV. Based on the above theory and Eq. 6, we conclude that T_1 follows the EY mechanism for the π bands, whereas it is described by the novel mechanism for the σ bands. With this in mind and the two band model result of Eq. 7, we describe the CESR linewidth with:

$$B = B_0 + \frac{1}{\sim^2} \frac{N_\pi L_e^2}{\frac{1}{2} \frac{1}{\tau_\pi^2}} + \frac{N_\sigma L_e^2}{1 + \frac{1}{2} \frac{1}{\tau_\sigma^2}}; \quad (8)$$

where we introduced the band index for the parameters. The momentum relaxation times are calculated using the Debye-model assuming clean samples, i.e. zero residual scattering:

$$\frac{1}{\tau_n} = \frac{2 k_B T}{\sim} \frac{Z}{\omega_D} \frac{d}{d\omega} \frac{\omega^4}{\sinh \frac{\omega}{2k_B T}}; \quad (9)$$

where $n = \sigma, \pi$; ω_D is the Debye frequency, and τ_{trm} are the transport electron-phonon couplings from Ref. [16], which contain both intra- and inter-band scattering.




FIG. 4: Measured (symbols) and calculated (solid curves) CESR line-width in MgB_2 with ^{11}B and ^{10}B . Note the larger residual line-width in the latter sample. Dashed curves show the contributions to the line-width from the σ and π bands for the ^{11}B . Dotted curve shows a calculation for the ^{11}B sample assuming $l=T_1$ is due to π bands only.

In Fig. 4., we show the CESR line-width for Mg^{11}B_2 and Mg^{10}B_2 between 40 and 700 K and the calculated line-width using Eq. 8. with parameters in Table I. obtained from a fit. Results on the natural boron sample are identical to the data on the Mg^{11}B_2 within experimental error and are not shown. The larger residual line-width in the ^{10}B ($B_0 = 2 \text{ mT}$) than in the ^{11}B sample ($B_0 = 1 \text{ mT}$) is related to a larger defect concentration in the starting boron, the preparation method and the starting Mg being identical. Apart from this, the only difference between the two samples are the different Debye temperature, Θ_D . The calculated CESR line-width (solid curves) reproduces well the experimental data with the parameters in Table I. The dotted curve in Fig. 4. is a calculation assuming that relaxation is given by the π bands alone, which accounts relatively well for the data with three free parameters (L_e , E_e , σ , and B_0). However, it fails to reproduce the slope of B at higher temperatures, which shows the need to include relaxation due to the σ bands.

The determination of E_e ; 0.2 eV is robust as it is given by the temperature where the maximal B is attained and its value is close to values expected from the band structure (arrows in Fig. 3.). Knowledge of E_e ; allows to determine the SO splitting independently, L_e ; $= 0.64 \text{ meV}$, as usually only the $L=E$ ratio is known. The SO splitting for the atomic boron $2p$ orbital is $L = 0.23 \text{ meV}$ (Ref. [4]), which is in a rea-

TABLE I: Parameters used to calculate the CESR line-width in MgB_2 . The given standard deviations indicate the free parameters of the fit.

τ [16]		L_e (meV)		E_e (eV)		Θ_D (K)	
						^{11}B	^{10}B
1.09	0.46	0.64(2)	2.8(1)	0.194(5)	2	535(15)	555(15)

sonable agreement with the experimental value. E_e was fixed to 2 eV which affects L_e ; as these are not independent.

The isotope effect on p_D is $\frac{p_D^{10}}{p_D^{11}} = 1.04$, that is close to the expected $\frac{11}{10}$ ratio. The Θ_D values are in agreement with the $440..1050 \text{ K}$ values in the literature, which scatter depending on the experimental method [20, 21]. We note that the model could be improved by including the Einstein model of phonons or by an exact treatment of the band structure dependent SO coupling [22], and band-band separation.

Finally, we note that the maximum of $l=T_1$ occurs when $\sigma \approx 1$. This coincides with the Ioffe-Regel criterion for the electron transport [23] when the band-band separation is comparable to the bandwidth, w , e.g. in narrow band metals. For MgB_2 , $w \approx 10 \text{ eV}$ [8] therefore saturation of the CESR line-width is not accompanied by a saturation of electrical resistivity.

In conclusion, we explained the anomalous spin-lattice relaxation in MgB_2 by extending the Elliott-Yafet theory to the case of rapid momentum scattering and nearby bands. The anomaly does not occur in conventional metals, which have small electron-phonon coupling and well separated bands. A similar phenomenon, the so-called Dyakonov-Perel relaxation [1], occurs for semiconductors without inversion symmetry, although its physical origin is different. The band structure of some of the other diborides in e.g. BeB_2 and CaB_2 predicts [19] similar phenomena but conventional spin relaxation in AlB_2 , ScB_2 , and YB_2 . We also predict that the described effect is sensitive to pressure since this shifts the bands [24].

We are grateful to J. Fabian and A. Virosztek for enlightening discussions. FS and FM acknowledge the Bolyai programme of the Hungarian Academy of Sciences and the Humboldt Foundation for support. Work supported by the Hungarian State Grants (OTKA) No. F61733, K72613, and NK60984. Ames Laboratory is operated for the U.S. Department of Energy by Iowa State University under Contract No. W-7405-Eng-82.

Corresponding author: simon@esr.phy.bme.hu

^Y Present address: IFW Dresden, Institute for Solid State Research, P.O. Box 270116, D-01171 Dresden, Germany

^Z Present address: Condensed Matter Physics and Materials Science Department, Brookhaven National Laboratory, Upton, New York 11973-5000, USA

Electronic address: simon@esrphy.bme.hu

- [1] I. Zutic, J. Fabian, and S. D. Samra, *Rev. Mod. Phys.* **76**, 323 (2004).
- [2] G. Feher and A. F. Kip, *Physical Review* **98**, 337 (1955).
- [3] R. J. Elliott, *Phys. Rev.* **96**, 266 (1954).
- [4] Y. Yafet, *Solid State Phys.* **14**, 1 (1963).
- [5] F. Beuneu and P. Monod, *Phys. Rev. B* **18**, 2422 (1978).
- [6] L. Forro and F. Beuneu, *Solid State Commun.* **44**, 623 (1982).
- [7] P. Petit, J. Robert, T. Yildirim, and J. E. Fischer, *Phys. Rev. B* **54**, 3764(R) (1996).
- [8] F. Simon, A. Janossy, T. Feher, F. Muranyi, S. Garaj, L. Forro, C. Petrovic, S. L. Bud'ko, G. Lapertot, V. Kogan, et al., *Phys. Rev. Lett.* **87**, 047002 (2001).
- [9] F. J. Adrian, *Phys. Rev. B* **53**, 2206 (1996).
- [10] F. Simon, F. Muranyi, T. Feher, A. Janossy, L. Forro, C. Petrovic, S. L. Bud'ko, and P. C. Canfield, *Phys. Rev. B* **75**, 024519 (2007).
- [11] P. Monod and M. Elsen, private communication.
- [12] R. R. Urbano, P. G. Pagliuso, C. Rettori, Y. Kopelevich, N. O. Moreno, and J. L. Sarrao, *Phys. Rev. Lett.* **89**, 087602 (2002).
- [13] H. Mori and K. Kawasaki, *Prog. Theor. Phys.* **27**, 529 (1962).
- [14] M. Oshikawa and I. A. A. Eck, *Phys. Rev. B* **65**, 134410 (2002).
- [15] A. A. Abrikosov, L. P. Gor'kov, and I. E. Dzyaloshinski, *Methods of Quantum Field Theory in Statistical Physics* (Dover Publications, New York, 1963).
- [16] I. I. Mazin, O. K. Andersen, O. Jepsen, O. V. Dolgov, J. Kortus, A. A. Golubov, A. B. Kuzmenko, and D. van der Marel, *Phys. Rev. Lett.* **89**, 107002 (2002).
- [17] H. J. Choi, D. Roundy, H. Sun, M. L. Cohen, and S. G. Louie, *Nature* **418**, 758 (2002).
- [18] J. Kortus, I. I. Mazin, K. D. Belashchenko, V. P. Antropov, and L. L. Boyer, *Phys. Rev. Lett.* **86**, 4656 (2001).
- [19] N. I. Medvedeva, A. L. Ivanovskii, J. E. Medvedeva, and A. J. Freeman, *Phys. Rev. B* **64**, 020502(R) (2001).
- [20] V. A. Gasparov, N. S. Sidorov, and M. P. Zver'kova, I. I. Kulakov, *JETP Lett.* **73**, 532 (2001).
- [21] M. Putti, E. G. d'Agliano, D. Marre, F. Napoli, M. Tassisto, P. Manfrinetti, A. Palenzona, C. Rizzuto, and S. Massidda, *Eur. Phys. J. B* **25**, 439 (2002).
- [22] J. Fabian and S. Das Samra, *Phys. Rev. Lett.* **83**, 1211 (1999).
- [23] O. Gunnarsson, *Rev. Mod. Phys.* **75**, 1085 (2003).
- [24] K. Kobayashi and K. Yamamoto, *J. Phys. Soc. Jap.* **70**, 1861 (2001).