

# A fusion of the LAPW and the LMTO methods: the augmented plane wave plus mu n-tin orbital (PMT) method

Takao Kotani and Mark van Schilfjaarde  
 School of Materials, Arizona State University, Tempe, AZ, 85284  
 (dated: April 2, 2024)

We present a new full-potential method to solve the one-body problem, for example, in the local density approximation. The method uses the augmented plane waves (APW s) and the generalized mu n-tin orbitals (MTO s) together as basis sets to represent the eigenfunctions. Since the MTO s can efficiently describe localized orbitals, e.g, transition metal 3d orbitals, the total energy convergence with basis size is drastically improved in comparison with the linearized APW method. Required parameters to specify MTO s are given by atomic calculations in advance. Thus the robustness, reliability, easy-of-use, and efficiency at this method can be superior to the linearized APW and MTO methods. We show how it works in typical examples, Cu, Fe, Li, SrTiO<sub>3</sub>, and GaAs.

PACS numbers: 71.15.Ap, 71.15.Fv 71.15.m

There are several kinds of all-electron full potential (FP) methods to obtain numerically-accurate solutions in the local density approximation to the density functional theory [1]. Among such FP methods, most popular ones are the linearized augmented plane wave (LAPW) method, and the projector augmented-wave (PAW) method [2, 3, 4, 5]. They both use plane waves (PW s) to expand the eigenfunctions in the interstitial regions. However, PW s do not efficiently describe the localized character of eigenfunctions just around the mu n-tins (MT s). For example, oxygen 2p (denoted as O(2p) below) and transition metal's 3d orbitals are well localized and atom-like even in solids, thus we need to superpose many PW s to represent these orbitals. For example, as shown in Ref.6 (and below), the energy cutoff for the augmented PW (APW)  $E_{MAX}^{APW} > 15Ry$  is required in fcc Cu to obtain  $\sim 1mRy$  convergence for total energy in LAPW. In contrast, such orbitals can be rather effectively represented by localized basis in realspace. In fact, it is already accomplished in the linearized mu n-tin orbital (LMTO) method, which differs from the LAPW method in that envelope functions consists of atom-like localized orbitals [7, 8] instead of PW s. Such a localized augmented waves are called as MT orbital (MTO).

To circumvent the inefficiency in the LAPW method, we have implemented a new method named as linearized augmented plane wave plus mu n-tin orbital (PMT) method. The PMT method includes not only the APW s but also MTO s in its basis set. Our implementation becomes LAPW in the no MTO limit. As we show later, we can very effectively reduce the number of basis set by including MTO s; we see the rapid convergence of the total energy as a function of the number of APW s (or energy cutoff  $E_{MAX}^{APW}$ ). As far as we tested, APW s with  $E_{MAX}^{APW} \sim 5Ry$  in addition to minimum MTO s will be reasonably good enough for usual applications; e.g. for  $< 1mRy$  convergence of total energy for Cu. Even in comparison with the LMTO method of Ref.[8], the PMT method is quite advantageous in its simplicity. The parameters to specify these minimum MTO s ( $E_H$  and  $R_H$  for each l channel. See next paragraph.) are automatically deter-

mined by atomic calculations in advance. This is a great advantage practically because optimization of these parameters is a highly non-linear problem [8] which makes the LMTO difficult to use. Thus the PMT method can satisfy the requirements for latest first-principle methods, reliability, speed, easy-of-use, and robustness very well. In what follows, we explain points in our method, and then we show how it works.

We adapted a variant of the LMTO method developed by Methfessel, van Schilfjaarde, and their collaborators [7, 8]. This method uses the "smooth Hankel function" invented by Methfessel as a modification of the usual Hankel functions so as to mimic the atomic orbitals [8, 9]. It contains an additional parameter called as the smoothing radius  $R_H$  in addition to the usual energy parameter  $E_H$  which specifies the damping behavior. By choosing  $R_H$ , we can control bending of the function just outside of MT. By using this degree of freedom, we can reproduce eigenvalues of atoms very well even if we substitute the eigenfunction outside of MT with such a smooth Hankel function. This is an important feature of the function in comparison with others like Gaussians, which are not directly fit to the atomic orbitals. Analytic properties of the smooth Hankel are analyzed in detail by Bott et al, and all the required operations, e.g, Bloch sum, to perform electronic structure calculations are well established [9]. The augmentation procedure requires the one-center expansion of the envelope functions. In our method, the one-center expansion is given as the linear combinations of the generalized Gaussian (Gaussians polynomials), where the expansion coefficients are determined by a projection procedure as described in Sec.XII in Ref.9. Then the generalized Gaussians in each angular momentum l-channel are replaced by the linear combinations of a radial function  $\chi_l$  and its energy derivative  $\chi_l'$  in the usual manner of augmentation [8]. When we use high energy cutoff  $E_{MAX}^{APW} \sim 15Ry$ , we needed to use 15 generalized Gaussians for the one-center expansion; however,  $\sim 5$  is good enough for practical usage with  $E_{MAX}^{APW} = 5Ry$ .

Another key point in our method is the smoothed




FIG. 1: (color online) Total energy is plotted as a function of (number of basis at  $\Gamma$  point  $N_b$ )<sup>-1</sup>. Lower panels for each material zoom up upper panels in energy (in SrTiO<sub>3</sub>, data points in LAPW are out of the scale in lower panel). We also plot spin moment ( $\mu_B$ ) for Fe, and the band gaps for GaAs and SrTiO<sub>3</sub> by dotted lines, referring to right y-axis. The absolute values of energies are ambiguous (See text in Table I).  $E_{MAX}^{APW}$  (Ry) are shown next to each data point. "4d lo" means we treat 4d as local orbital. PMT denotes our new method with APW + MTO, where we use minimum MTO basis; number of MTOs (including lo) are 14 (Cu), 5 (Li), 9 (Fe), 18 (GaAs), and 33 (SrTiO<sub>3</sub>). Number of k point in 1st Brillouin zone (BZ) are 12<sup>3</sup> for Cu and Li, 20<sup>3</sup> for Fe, and 6<sup>3</sup> for GaAs and SrTiO<sub>3</sub>. Lattice constants are not necessarily at their total energy minimum (e.g., 6.8a.u for Cu). See Table I for exchange-correlation used.

charge density treatment introduced by Soler and Williams [3] to the LAPW method. The charge density is divided into the smooth part, the true density on MTOs, and the counterpart on MTOs. The smooth part is tabulated on regular uniform mesh in real space, and the others are tabulated on radial mesh in the spherical harmonics expansion in each MTO sphere. The PAW methods [4, 5] also use this treatment. It enables low-angular momentum cutoff for augmentation and makes the calculation very efficient. As for the regular mesh in real space, we usually need to use the spatial resolution corresponding to the cutoff energy  $E_{MAX}^{mesh} = 10-15$  Ry, which is determined so as to reproduce the smooth Hankel function well in real space. Note that we still have some inefficiency, e.g. an atom in a large supercell requires a fine mesh everywhere only in order to describe the density around the atom. This problem is common to any method which uses a uniform mesh for density.

Though the LMTO formalism shown in [7, 8] is intended for such MTOs constructed from the smooth Hankel functions, it is essentially applicable to any kind of envelope functions. As we explained above, our formalism is not so different from LAPW/PAW formalisms shown in [4, 5] except in the augmentation (projection) proce-

dure. The atomic forces are calculated [8, 10] in the same manner as in PAW [5]. For deep cores, we usually use a frozen core approximation which allows the extension of core densities outside of MTO (but no augmentation) [7]. Further, we use the local orbital (lo) method [11] in some cases; for example, to treat 3d semicore for Ga (denoted as Ga(3d lo) below), or to reproduce high-lying bands for Cu by Cu(4d lo).

Results: In Fig.1, we plot the total energies as functions of the inverse of the number of basis at the  $\Gamma$  point ( $N_b$ )<sup>-1</sup> in order to observe its convergence as  $N_b \rightarrow 1$  (the number of basis is controlled by  $E_{MAX}^{APW}$  as shown on Fig.1 together). Here we include minimum MTOs whose parameters  $E_H$  and  $R_H$  are fixed by the atomic calculations (here "minimum" means only the atomic bound states). The lo's are included as explained in Fig.1.

There is a problem of linear dependency in the basis set when we use large  $E_{MAX}^{APW}$ . For example, in Li, we could not include MTOs of Li(2s2p) (Li(2s) and Li(2p)) as basis at  $E_{MAX}^{APW} > 6$ , because then they are well expanded by PWs. This occurs also for other cases; when we use  $E_{MAX}^{APW}$  high enough to expand a MTO, the rank of the overlap matrix of basis set is reduced by one. Thus possible ways to use large  $E_{MAX}^{APW}$  are: (1) keep only well

localized MTOs which are not yet expanded by given  $E_{MAX}^{APW}$ , or (2) remove a subspace of basis through the diagonalization procedure of the overlap matrix before solving the secular equation. For (2), we need to introduce some threshold to judge the linear dependency. This can cause an artificial discontinuity when changing lattice constants and so on. Thus (1) should be safer, but here we use the procedure (2) with careful check so that such discontinuity do not occur. We use the number of basis after the procedure (2) for  $N_b$  to plot Fig.1. Even for LAPW cases, we applied the procedure (2), e.g. to the case for  $E_{MAX}^{APW} = 20Ry$  in SrTiO<sub>3</sub>; then we reduce the dimension of Hamiltonian from 606 to  $N_b = 550$  (data point at the left end of SrTiO<sub>3</sub> in Fig.1).

As is seen in all the cases, the PMT method shows smooth and rapid convergence for the total energy at  $(N_b)^{-1} \rightarrow 0$ . On the other hand, the convergence in LAPW (no MTO limit in our PMT implementation) shows a characteristic feature; it is way off until it reaches to some  $E_{MAX}^{APW}$ , e.g. 15 Ry in Cu. This is consistent with previous calculations [6].  $E_{MAX}^{APW} = 15$  is required to reproduce the 3d localized orbitals. We also show the convergence behaviors for band gap and magnetic moments by dotted lines (right y-axis); they are quite satisfactory.




FIG. 2: Energy bands for Cu up to 100 eV above the Fermi energy. Top-left is for APW only without local orbitals. Top-right: APW + 4d[lo]. Bottom-left: APW plus MTO (9 basis) + 4d[lo]. Bottom-right: MTO (34 basis) + 4d[lo]. All are converged for  $E_{MAX}^{APW}$ .

Fig.2 shows the energy bands up to 100eV above the Fermi energy. Though the Cu panel in Fig.1 shows the little effects of 4d local orbital to the total energy, it affects energy bands above 30eV. Calculations including 4d[lo] gives good agreements each other. This means




FIG. 3: Total energy v.s. lattice constant. Labels  $E_{MAX}$  means  $E_{MAX}^{APW}$ . MTO (high) for Cu is explained in text.

that we have no artificial bands (ghost bands). The MTO (high) panel is by the pure LMTO method where we use 34 + 5 basis (spdfg + spd + 4d[lo]). With some careful choice of  $E_H$  and  $R_H$ , the LMTO method can be very efficient and accurate.




FIG. 4: Total energy for different type of MTO with changing the number of APWs  $n(APW)$  for SrTiO<sub>3</sub>. Each curves (A, B, C, D, E, F) corresponds to a type of MTO sets (see texts). For example, curve C uses 36 MTOs (including local orbital). Thus curve C at  $n(APW) = 80$  means a calculation with  $n(MTO) + n(APW) = 36 + 40 = 76$  basis. The converged energy  $-2.76$  Ry is a little different from the Fig1 because this calculation is performed with coarse conditions for test purpose.

Fig.3 shows the total energy as function of the lattice constant. All lines look very parallel. This shows that PMT do not include some systematic errors. This is true

TABLE I: Calculated lattice constant, bulk modulus, and cohesive energies. Values are by the PMT method with  $E_{MAX}^{APW} = 15Ry$  ( $E_{MAX}^{APW} = 6Ry$  for Li). Values in parenthesis are with  $E_{MAX}^{APW} = 5Ry$ . We show values from other literatures together. We used the Barth-Hedin exchange correlation [12] except Fe where we use the VWN type [13]. Since we have not determined the total energies of each atoms exactly (we assume spherical atoms and assume some electron configurations), the cohesive energies (constant in energy axis in Fig. 1 and Fig. 3) are somewhat ambiguous; especially for cases including transition metals. For comparison with other calculations, it will be better to use total energies; add  $-3304.4345Ry$  (Cu),  $-14.7106Ry$  (Li),  $-2540.4767Ry$  (Fe),  $-8397.5970Ry$  (GaAs), and  $-8502.4637Ry$  (SrTiO<sub>3</sub>) to the cohesive energies.

	Lattice constant. (a.u.)	Bulk Modulus (GPa)	Cohesive energy (Ry)
fcc Cu	6.650 (6.649)	188 (187)	-332 (-331)
LAPW	6.65	192	
expt. <sup>a</sup>	6.81	131	
bcc Fe	5.209 (5.208)	258 (259)	-.667 (-.665)
PAW <sup>b</sup>	5.20	247	
LAPW	5.210	245	
expt. <sup>c</sup>	5.417	172	
bcc Li	6.347 (6.341)	15.3 (15.5)	-.124 (-.123) <sup>g</sup>
PAW <sup>b</sup>	6.355	15.0	-.149
SrTiO <sub>3</sub>	7.367 (7.360)	220 (225)	-2.731 (-2.709)
PP <sup>d</sup>	7.31	203	
expt. <sup>e</sup>	7.39	184	
GaAs	10.61 (10.61)	74.9 (75.0)	-.576 (-.574)
LAPW	10.62	74	-.587
expt. <sup>e</sup>	10.68	76	-.479

<sup>a</sup>Ref. [14]

<sup>b</sup>Ref. [5]

<sup>c</sup>Ref. [15]

<sup>d</sup>Ref. [16]. PP denotes a pseudopotential method.

<sup>e</sup>Ref. [17]

<sup>g</sup>The difference from PAW may be because it uses the non-polarized Li atom as reference. If we do so, we have  $-.150$  ( $-.150$ ).

also for other materials (not shown). Table I shows the obtained lattice constants and related parameters. We also showed values for  $E_{MAX}^{APW} = 5Ry$ . Even though we still need other extensive tests, we think that such low cutoff is reliable enough for practical applications. For  $E_{MAX}^{APW} = 5Ry$ , we only need 25 basis (when we do not include b) for Cu.

Fig. 4 shows the total energies with different MTO sets combined with different numbers of APW s. Curve "A" includes just MTO s for O (2s2p), sufficient for a crude representation of the valence bands. Also included are Sr(4p [b]) and Ti(3p [b]). A large number of APW s is needed to get a good total energy: 150 APW s are needed to converge energy to within 50 mRy of the converged result. "B" corresponds to an extreme tight-binding basis, consisting of Sr(5s5p) and Ti(4s4p4d) in addition to "A". (Note that the conduction band is mainly Ti(3d), and O (2s2p)). The total energy of the

MTO basis alone (no APW s) is rather crude | more than 200 mRy underbound. However, only 25 orbitals (plus 6 for b's) are included in this basis. The energy drops rapidly as low-energy APW s are included: adding 40 APW s is sufficient to converge energy to 50 mRy. As more APW s are added, the gain in energy becomes more gradual; convergence is very slow for large  $E_{MAX}^{APW}$ . "C" differs from "B" only in that Sr(3d) orbital was added. With the addition of these 5 orbitals, the MTO-only basis is already rather reasonable. This would be the smallest acceptable MTO-basis. As in the "B" basis, there is initially a rapid gain in energy as the first few APW s are added, followed by a progressively slower gain in energy as more APW s are added. "D" is a standard LMTO minimum basis: spd orbitals on all atoms. Comparing "C" or "E" to "A" shows that the MTO basis is vastly more efficient than the APW basis in converging the total energy. This is true until a minimum basis is reached. Beyond this point, the gain APW s and more MTO s improve the total energy with approximately the same efficiency, as the next tests show. "E" is a standard LMTO larger basis: spd + spd orbitals on Sr and Ti, and spd + sp on O. Comparing "D" and "F" shows that the efficiency of any one orbital added to the standard MTO minimum basis is rather similar in the APW and MTO cases. Thus, increasing the MTO basis from 51 to 81 orbitals in the MTO basis lowers the energy by 33 mRy; adding 33 APW s to the minimum basis ("D") lowers the energy by 36 mRy. "F" enlarges the MTO basis still more, with Sr: spd + spd, Ti: spd + spd, O: spd + spd. Also local orbitals are used to represent the high-lying states Ti(4d [b]) and O (3s [b], 3p [b]). For occupied states, these orbitals have little effect for total energy as in the case of Cu.

In conclusion, we have implemented the PMT method whose basis set consists of the APW s together with the MTO s which are localized in real space. This idea is consistent with the nature of the eigenfunctions in solids; they can be localized as atoms or extended as PW. This method combines advantages of LAPW /PAW and LMTO. We have implemented force calculations, but no results are shown here. One of the advantages is its flexibility. As an example, in order to treat Cu impurity in Si, it will be possible to choose very low  $E_{MAX}^{APW}$  just responsible for empty regions because MTO s for Cu and MTO s for Si span its neighbors already very well. Convergence is easily checked by changing  $E_{MAX}^{APW}$  (much simpler than LMTO). In future, we can use the PMT method to make a natural division of the Kohn-Sham Hamiltonian into localized blocks and extended blocks, instead of the energy window method for the maximally localized Wannier functions [18]. The problem of large  $E_{MAX}^{mesh}$  should be solved to make PMT more efficient.

This work was supported by ONR contract N00014-7-1-0479. We are also indebted to the Ira A. Fulton High Performance Computing Initiative.

- 
- [1] W. Kohn and L. J. Sham, *Phys. Rev.* **140**, A1133 (1965).
- [2] O. K. Andersen, *Phys. Rev. B* **12**, 3060 (1975).
- [3] J. M. Soler and A. R. Williams, *Phys. Rev. B* **40**, 1560 (1989).
- [4] P. E. Blochl, *Phys. Rev. B* **50**, 17953 (1994).
- [5] G. Kresse and D. Joubert, *Phys. Rev. B* **59**, 1758 (1999).
- [6] A. Y. Liu, D. J. Singh, and H. Krakauer, *Phys. Rev. B* **49**, 17424 (1994).
- [7] M. Methfessel and M. van Schilfjaar, 'NFP manual 1.01 Oct 10, 1997'. NFP is previous to the current LMTO package maintained by M. van Schilfjaar.
- [8] M. Methfessel, M. van Schilfjaar, and R. A. Casali, in *Lecture Notes in Physics*, edited by H. Dreyse (Springer-Verlag, Berlin, 2000), vol. 535.
- [9] E. Bott, M. Methfessel, W. Krabs, and P. C. Schmidt, *J. Math. Phys.* **39**, 3393 (1998).
- [10] M. Methfessel and M. van Schilfjaar, *Phys. Rev. B* **48**, 4937 (1993).
- [11] D. J. S. E. Sjøstedt, L. Nordström, *Solid State Communications* **114**, 15 (2000).
- [12] U. von Barth and L. Hedin, *J. Phys. C* **5**, 1692 (1972).
- [13] L. W. S. H. Vosko and M. Nusair, *Can. J. Phys.* **58**, 1200 (1980).
- [14] A. Klein, D. J. Singh, and C. J. Umrigar, *Phys. Rev. B* **51**, 4105 (1995).
- [15] L. Stixrude, R. E. Cohen, and D. J. Singh, *Phys. Rev. B* **50**, 6442 (1994).
- [16] L. M. Liborio, C. G. Sanchez, A. T. Paxton, and M. W. Finnis, *Journal of Physics: Condensed Matter* **17**, L223 (2005), URL <http://stacks.iop.org/0953-8984/17/L223>.
- [17] C. Filippi, D. J. Singh, and C. J. Umrigar, *Phys. Rev. B* **50**, 14947 (1994).
- [18] I. Souza, N. Marzari, and D. Vanderbilt, *Phys. Rev. B* **65**, 035109 (2001).