

New Bound States of several Top-quarks bound by Higgs Exchange

C. D. Froggatt
Glasgow University

H. B. Nielsen
Niels Bohr Institute, Blegdamsvej 17 -21, DK 2100 Copenhagen

It is suggested that there is a fine tuning principle remarkably arranging the top-quark Yukawa coupling to be just so as to make the mass m_{12} of a bound state of 6 top + 6 anti-top quarks become very small. The value of the top Yukawa needed for Higgs and gluon etc. exchanges just binding the 12 quarks to make this mass zero is $g_t|_{m_{12}=0} = 1.01 \pm 0.15$, while the experimental running top Yukawa coupling is $g_t = .94$.

1. INTRODUCTION

Because of the top mass being of the order of e.g. the W-masses, the top Yukawa coupling is of order unity. The relevant quantity for estimating when the coupling is sufficiently strong to lead to non-perturbative effects is rather $\frac{g_t^2/2}{4\pi}$. So it is only if we could get of the order of 4π particles to function together that severe strong interaction effects would show up, but there are just 12 top quarks and anti-top quarks in the 1s shell.

2. HIGGS EXCHANGE

Like other even order tensor field exchanges - such as the gravitational field - the scalar Higgs field causes attraction between all combinations of say top quarks and anti-top quarks. So it is just the number of tops and anti-tops together that count for the strength of binding - the binding energy for a bound state of such quarks. In first approximation, analogous to the Bohr atom, one finds [1] that the binding energy goes as $E_{binding} \propto (g_t^2 Z)^2 m_t$, where Z is the number of particles considered to lie in the centre and attract the $(Z + 1)$ 'th particle.

For a full 1s shell of $6t + 6\bar{t}$ in the Bohr model approximation we find for t - and u -channel Higgs exchange:

$$E_{binding} = \frac{11g_t^4}{2\pi^2} m_t \quad (1)$$

3. REMARKABLE COINCIDENCE

We imagine calculating in the infinite momentum frame so as to make the in principle relativistic calculation look more non-relativistic. After including gluon exchange and allowing for the left-handed top quarks to be replaced by left-handed b -quarks, we arrive at the result that the binding energy could be so big as to compensate for the mass of the 6 top and 6 anti-top quarks within the uncertainty in our calculation. Actually we formulate our calculation so as to evaluate the value $g_t|_{m_{12}=0}$ of the top Yukawa coupling, which would correspond to the exact cancellation of the binding energy with the energy of the masses, so as to make the 12 quark bound state exactly massless $m_{12} = 0$. The resulting value turns out to look like a remarkable coincidence. We namely included a long series of corrections in estimating this top Yukawa coupling value and finally found

$$g_t|_{m_{12}=0} = 1.01 \pm 0.15 \quad (2)$$

in very good agreement with the experimental value of the running top Yukawa coupling - which is the one that should be compared to the value (2) we calculated:

$$g_t^{exp} = .935 \pm .080. \quad (3)$$

4. FINE TUNING CALLS FOR MPP

This remarkable coincidence presents us with a fine tuning problem one could say, and thus calls for some principle or law about the values of coupling constants, such as in the present case the top-Yukawa coupling. We have worked for some time on the idea of postulating such a principle, the so-called “Multiple Point Principle” (MPP), which postulates [2] that there shall be many vacua/phases with the same energy density. The approximate masslessness of a particle can in fact be a consequence of this “Multiple Point Principle”, when there are two degenerate phases or vacua just deviating by the particle in question being condensed in one of the phases while it is not condensed in the other one. Such an at least near degeneracy of a condensed and an uncondensed phase of the bound states we have discussed is anyway unavoidable, if the mass squared of the bound state is a smooth function of the coupling constants and in particular of the top Yukawa coupling in our case. In fact if the coupling is a bit stronger than our $g_t|_{m_{12}=0}$ it means, in the infinite momentum frame language, that the bound state becomes a tachyon and that is well-known to imply that there should be a condensate, i.e. a non-zero expectation value for an effective field describing the bound state. On the other hand if the coupling is weaker than our phase border value $g_t|_{m_{12}=0}$, then the bound state has positive mass squared and there will be no condensate, i.e. the expectation value of the field will be zero. The above means that, taking the coincidence as a fine tuning problem that has to be explained, then our “multiple point principle” at least in this case is strongly called for.

5. SPECTRUM OF BOUND STATES

For bound states of just Z top or anti-top particles, the binding energy per particle is proportional (approximately) to Z^2 and so we expect the mass squared m_Z^2 of the bound state of Z such particles to be of the form $m_t^2 Z^2(1 - aZ^2)$ where a is a constant. Assuming, as expected from our infinite momentum frame thinking, that it is m_Z^2 that shall be analytic, and using that from our fine tuning assumption the particle with $Z = 12$ shall be massless, we get, keeping the lowest order Taylor expansion terms, that the bound state with Z constituents has the mass squared:

$$m_Z^2 = m_t^2 Z^2(1 - (Z/12)^2). \quad (4)$$

Since this formula has the character of an interpolation, we think it is rather trustable and we obtain for instance for the bound states with $Z = 11$ a mass of $m_{11} = 760$ GeV and for the one with $Z = 10$ constituents $m_{10} = 950$ GeV. This means that we expect a spectrum of related particles of the following form.




Figure 1: Mass spectrum of bound states.

We can even imagine a W boson being attached to a system of say $Z = 12$ quarks. Then the bound system is in a superposition of being indeed a system with a W added and one in which, instead, we have the system that would be obtained by having replaced a left-handed top quark by a left-handed b -quark in the 12-constituent state. We expect such a bound state with an extra W would have mass closer to that of the $Z = 11$ bound state than to the one with $Z = 12$. Either there would be an extra W that would have to be strongly localized - of the order of the radius - or there would be only 11 top or anti-top quarks although then together with a b -quark too.

6. PRODUCTION AT LHC OR TEVATRON

Both the colour neutral $6t + 6\bar{t}$ and the related resonances with 11 constituents are strongly bound states and have small radii $r_0 \sim 1/m_t$ and their production rate may be estimated as if they were “fundamental particles” just with some form factor corrections. The bound states with 12 constituents cannot, however, be produced by gluon collision directly, because they are colour singlets, but can be produced as decay products from the heavier $6t + 5\bar{t}$ colour triplet bound states, whose pair production cross section should be similar to that for a 4th generation quark.




Figure 2: Feynman diagram for pair production of the $6t + 5\bar{t}$ colour triplet bound state and subsequent decay.

If indeed the 12-constituent bound state is very light it would decay into a few gluon jets, or if very light might look effectively like one jet at the high energies needed. This light bound state would be co-produced with a top quark or anti-top quark resulting from the decay of the 11-state. These 11-states would only be produced in pairs and so we should see a $t\bar{t}$ pair being produced together with two of the light bound states.

7. OTHER MPP PREDICTIONS

The existence of a Planck scale degenerate vacuum in the Standard Model would lead to a Higgs mass prediction:

$$m_h = \begin{cases} 132 \pm 5 \text{ GeV} & \text{exact degeneracy version [3]} \\ 118 \pm 5 \text{ GeV} & \text{metastable version [4]} \end{cases}$$

The fine tuning needed in this paper to ensure the existence of a degenerate phase containing a 12 quark bound state condensate, together with the existence of a Planck scale vacuum, would solve the hierarchy problem [1] in the Standard Model. The values of the fine structure constants can be explained using MPP in the family replicated gauge group model [1, 2]. The MPP assumption has been adapted to SUGRA models in order to explain the tiny value of the cosmological constant [5]. The application of MPP to the general 2 Higgs doublet model leads to an approximate custodial symmetry explaining the absence of flavour changing neutral currents [6].

References

- [1] C.D. Froggatt and H.B. Nielsen, *Surv. High Energy Phys.* **18** (2003) 55 [arXiv:hep-ph/0308144];
C.D. Froggatt, L.V. Iperashvili and H.B. Nielsen, *Int. J. Mod. Phys.* **A20** (2005) 1268 [arXiv:hep-ph/0406110].
- [2] D.L. Bennett, C.D. Froggatt and H.B. Nielsen, *Perspectives in Particle Physics '94*, p. 255, ed. D. Klabučar, I. Picek and D. Tadić (World Scientific, 1995) [arXiv:hep-ph/9504294].
- [3] C.D. Froggatt and H.B. Nielsen, *Phys. Lett.* **B368** (1996) 96 [arXiv:hep-ph/951137].
- [4] C.D. Froggatt, H.B. Nielsen and Y. Takahashi, *Phys. Rev.* **D64** (2001) 113014 [arXiv:hep-ph/0104161].
- [5] C.D. Froggatt, R. Nevzorov and H.B. Nielsen, *Nucl. Phys.* **B473** (2006) 133 [arXiv:hep-ph/0512061].
- [6] C.D. Froggatt, R. Nevzorov and H.B. Nielsen, arXiv:0806.3190[hep-ph].