

S_4 as a natural flavor symmetry for lepton mixing

Federica Bazzocchi and Stefano Morisi

*AHEP Group, Institut de Física Corpuscular – C.S.I.C./Universitat de València
Edificio Institutos de Paterna, Apt 22085, E-46071 Valencia, Spain*

(Dated: February 3, 2022)

Group theoretical arguments seem to indicate the discrete symmetry S_4 as the minimal flavour symmetry compatible with tribimaximal neutrino mixing. We prove in a model independent way that indeed S_4 can realize exact tribimaximal mixing through different symmetry breaking patterns. We present two models in which lepton tribimaximal mixing is realized in different ways and for each one we discuss the superpotential that leads to the correct breaking of the flavor symmetry.

PACS numbers: 11.30.Hv, 14.60.-z, 14.60.Pq, 14.80.Cp

I. INTRODUCTION

Harrison, Perkins and Scott (HPS) [1] proposed the so called tribimaximal mixing matrix

$$U_{TB} = \begin{pmatrix} \sqrt{2/3} & 1/\sqrt{3} & 0 \\ -1/\sqrt{6} & 1/\sqrt{3} & -1/\sqrt{2} \\ -1/\sqrt{6} & 1/\sqrt{3} & 1/\sqrt{2} \end{pmatrix}. \quad (1)$$

This matrix keeps in surprising agreement with experimental data [2]. Lot of theoretical models has been done to explain the mixing matrix of eq. (1) by means of non abelian flavor symmetry, such as S_3 [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13], A_4 [14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29], T' [30, 31, 32, 33, 34], S_4 [35, 36, 37, 38, 39] and $\Delta(27)$ [40, 41, 42, 43]. The non abelian discrete groups have irreducible representations of dimension bigger than one [44]. The most interesting case arises when the group contains a triplet as irreducible representation, allowing to embed the observed three generations of fermions.

When a non abelian discrete group G is broken to one of its subgroup G' the transformation $U_{G'}$ that decomposes the representations of G according to G' can be fixed and are completely model independent. This is the case for example of A_4 broken to Z_3 : the triplet representation of A_4 is sent to the one-dimensional representations of Z_3 , $1, 1', 1''$, through the matrix U_ω defined as

$$U_\omega = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}, \quad (2)$$

while the one dimensional representations of A_4 coincide with the corresponding ones of Z_3 . A good candidate to give TBM is a discrete group G that has a triplet representation, at least two subgroups, G' that decompose according to $U_{G'}$ and G'' that decompose according to $U_{G''}$. It is necessary having at least two different subgroups of G to obtain a lepton mixing matrix different to the identity: if G were broken to the same subgroup G' both in the charged lepton and in the neutrino sector the lepton mixing matrix would be given by $U_{lep} = U_{G'}^\dagger, U_{G'} = \mathcal{I}$.

A priori A_4 seems to be a good candidate because it is the smallest discrete group that contains a triplet as irreducible representation. Furthermore it has two different subgroups, Z_3 and Z_2 . However, while the transformation associated to Z_3 is given by U_ω the one associated to Z_2 is model dependent. This analysis has been already performed in [45] (see eq. A4). A similar analysis done with the discrete symmetry T' lead to the same conclusion (see eq. (8) of Ref.[30]). This means that A_4 and T' yield exact or approximate TBM only assuming a fine tuning in the parameters of the Yukawa lagrangian or a particular model realization. We mention that by assuming further constraints, also models based on S_3 can yield an approximate TBM, although its largest irreducible representation is a doublet and not a triplet.

It has been recently claimed [46] that the minimal flavor symmetry naturally related to the tribimaximal mixing is S_4 , the permutation symmetry of four objects. The author of [46] proved this through group theoretical arguments without entering into the details of a concrete model realization. In this paper we provide a concrete model realization

of these general arguments reconsidered S_4 and its subgroups. We have found that S_4 is able to reproduce TBM following two different symmetry breaking patterns. We have built two different models that realize TBM through the two patterns dictated by the group analysis considerations and finally we discuss the possible superpotential that can break S_4 in the correct way.

II. THE DISCRETE SYMMETRY GROUP S_4 AS THE ORIGIN OF TBM

A. The group S_4

The discrete group S_4 is given by the permutations of four objects and it is composed by 24 elements. It can be defined by two generators S and T that satisfy

$$S^4 = T^3 = 1, \quad ST^2S = T. \quad (3)$$

The 24 elements of S_4 belong to five classes

$$\begin{aligned} \mathcal{C}_1 &: I; \\ \mathcal{C}_2 &: S^2, TS^2T^2, S^2TS^2T^2; \\ \mathcal{C}_3 &: T, T^2, S^2T, S^2T^2, STST^2, STS, S^2TS^2, S^3TS; \\ \mathcal{C}_4 &: ST^2, T^2S, TST, TSTS^2, STS^2, S^2TS; \\ \mathcal{C}_5 &: S, TST^2, ST, TS, S^3, S^3T^2. \end{aligned} \quad (4)$$

The elements of $\mathcal{C}_{2,4}$ define two different sets of Z_2 subgroups of S_4 , that ones of the class $\mathcal{C}_4$ a set of Z_3 abelian discrete symmetries and those belonging to $\mathcal{C}_5$ a set of Z_4 abelian discrete symmetries. The S_4 irreducible representations are two singlets, $1_1, 1_2$, one doublet, 2 , and two triplets, 3_1 and 3_2 . We adopt the following basis

$$S = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \quad T = -\frac{1}{2} \begin{pmatrix} 1 & \sqrt{3} \\ -\sqrt{3} & 1 \end{pmatrix}, \quad (5)$$

for the doublet representation and

$$S_{+,-} = \pm \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \quad T = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad (6)$$

for the triplet representations. Clearly the generators (S_+, T) and (S_-, T) define the two triplet representations $3_1, 3_2$ respectively. All the product rules can be straightforwardly derived. We remind the reader to the product rules reported in [36].

B. S_4 symmetry breaking patterns

We have seen in the introduction that given a discrete non abelian group G a predictive lepton mixing matrix may be obtained if G is broken to one of its subgroups, with the subgroup preserved in the charged lepton sector different from the subgroup preserved in the neutrino sector.

We disregard therefore the case when S_4 is completely broken in one of the two sectors. At the same time, if the left handed leptons transform non-trivially under S_4 , the case of S_4 unbroken in one sector is ruled out since it leads to a diagonal mass matrix with at least two degenerate states. Therefore if S_4 is broken to one of its subgroups G' in the charged lepton sector, in the neutrino sector it has to be broken to another subgroup $G'' \neq G'$. The couple (G', G'') identifies a possible symmetry breaking pattern. In this notation the lepton mixing matrix is given by

$$U_{lep} = U_l^\dagger U_\nu = U_{G'}^\dagger U_{G''}, \quad (7)$$

being $U_{G'}, U_{G''}$ the transformations that decompose the representations of S_4 into the representations of G', G'' respectively.

S_4 contains a non abelian subgroup S_3 , the permutation group of three objects composed by six elements. The elements of S_4 that belong to S_3 correspond to C_1 , T and T^2 of C_3 and $TSTS^2, STS^2, S^2TS$ of C_4 . Furthermore S_4 contains the abelian subgroups Z_2 , Z_3 , Z_4 corresponding to the elements of the classes $\mathcal{C}_{2,4}, \mathcal{C}_3$ and $\mathcal{C}_5$ respectively. The only representation that can break S_4 to S_3 is the triplet 3_1 . The reason is that when a triplet $\phi_1 \sim 3_1$ develops vev as $(1,1,1)$ the six elements that define S_3 belonging to S_4 — $I, T, T^2, TSTS^2, STS^2, S^2TS$ built with the basis reported in eq. (6)—are preserved. On the contrary, when a triplet $\phi_2 \sim 3_2$ develops vev as $(1,1,1)$, only the three elements that define Z_3 are preserved— I, T, T^2 —while $TSTS^2, STS^2, S^2TS$ built according eq. (6) are broken.

The representations of S_3 are two singlet, 1_1 and 1_2 , and a doublet, 2 . In general if S_4 is broken to S_3 the representations of S_4 would transform under S_3 according to

$$3_1 \rightarrow 1_1 + 2, \quad 3_2 \rightarrow 1_2 + 2, \quad 2 \rightarrow 2, \quad 1_1 \rightarrow 1_1, \quad 1_2 \rightarrow 1_2. \quad (8)$$

Therefore if S_4 is broken to S_3 , a triplet of S_4 , $F \sim (F_1, F_2, F_3) \sim 3_1$, will decompose under S_3 as $F(3_1) \rightarrow \psi_0(1_+) + \psi(2_-)$ with

$$\psi_0 = \frac{1}{\sqrt{3}}(F_1 + F_2 + F_3), \quad \psi = \begin{pmatrix} (F_2 - F_3)/\sqrt{2} \\ (-2F_1 + F_2 + F_3)/\sqrt{6} \end{pmatrix}. \quad (9)$$

The new eigenstates S_3 (ψ_0, ψ) are defined by

$$\begin{pmatrix} \psi_0 \\ \psi_1 \\ \psi_2 \end{pmatrix} = U_{S_3} \begin{pmatrix} F_1 \\ F_2 \\ F_3 \end{pmatrix} \quad \text{with} \quad U_{S_3} = P \cdot U_{TBM}^T \quad \text{with} \quad P = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (10)$$

We now assume that $F \sim L$ being L the left handed lepton doublets and for the moment we leave undetermined the transformation properties under S_4 of the electroweak $SU(2)$ singlets.

The first case we consider is the symmetry breaking pattern (S_3, G''), that means that we break S_4 into S_3 in the charged lepton sector while we still not know which is its corresponding S_4 subgroup in the neutrino sector. Assuming that the LR charged lepton mass matrix M_l is obtained once S_4 is broken to S_3 , we can write $M_l M_l^\dagger$ in the new basis defined by eq. (10)

$$M_l M_l^\dagger \rightarrow P U_{TBM}^T M_l M_l^\dagger U_{TBM} P = \tilde{M}_l \tilde{M}_l^\dagger. \quad (11)$$

Since the residual symmetry is S_3 , $\tilde{M}_l \tilde{M}_l^\dagger$ has to be invariant under this symmetry. Once we impose this condition we discover that $\tilde{M}_l \tilde{M}_l^\dagger = M_{diag}^l M_{diag}^{l\dagger}$, with 2 degenerate masses. Neglecting for the moment this phenomenological inconsistency, we have seen that the breaking $S_4 \rightarrow S_3$ in the charged lepton sector has lead to $U_l = U_{TBM} P$. If the neutrino mass matrix were diagonal $U_{lep} = U_l^\dagger U_\nu$ would lead to the wrong conclusion $U_{lep} = U_{TBM}^T$. To cure this problem we have two options. On one hand, we could require that the neutrino mass matrix were diagonalized by $U_{TBM} U_{TBM}$ in order to reproduce the TBM through $U_{lep} = U_{TBM}^T U_{TBM} U_{TBM} = U_{TBM}$. However there is no G'' subgroup of S_4 that yields $U_{G''} = U_{TBM} U_{TBM}$ and therefore exact TBM cannot be obtained according to eq. (7). On the other hand we could require to break the surviving S_3 in the charged lepton sector into Z_2 in such a way to produce a $U_l \neq U_{TBM} P$. Even in this case there is no corresponding G'' in the neutrino sector that allows to obtain exact TBM. As consequence the symmetry breaking pattern with S_4 broken into S_3 in the charged lepton sector is ruled out.

We now analyze what happens considering the breaking pattern (Z_3, G''). As in the previous case the subgroup G'' , corresponding to the neutrino sector, is undetermined. We expect that if we break S_4 into Z_3 in the charged lepton sector—we have already said that in S_4 the breaking into Z_3 is realized when a triplet 3_2 develops a vev in the direction $(1,1,1)$ —the charged lepton mixing matrix will send the S_4 triplet (L_1, L_2, L_3) in the Z_3 eigenstates, $1, 1', 1''$. Indeed the mixing matrix responsible of this rotation is the U_ω defined in eq. (2). Given U_ω the correct TBM can be reproduced if the $U_{G''}$ of eq. (7) is given by

$$U_\nu = \begin{pmatrix} 0 & 1 & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{i}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & 0 & -\frac{i}{\sqrt{2}} \end{pmatrix}, \quad (12)$$

or in other words if the neutrino mass matrix m^ν is diagonalized by U_ν and it has the following form

$$m^\nu = \begin{pmatrix} a & 0 & 0 \\ 0 & c & b \\ 0 & b & c \end{pmatrix}. \quad (13)$$

The matrix form of eq. (13) is recovered by requiring the invariance of m^ν under the $G'' = Z_2$ subgroup of S_4 associated to the element TST of the class $\mathcal{C}_4$. This breaking pattern is the usual one used in models based on A_4 . However we stress that in the context of S_4 we have obtained TBM only according to group theory considerations.

If we consider now the case (Z_2, G'') we discover that S_4 behaves exactly as A_4 and exact TBM cannot be recovered. For a detailed analysis we remand the reader to the Appendix of [45].

In the case (Z_4, G'') we discover that the charged lepton mass matrix $M_l M_l^\dagger$ is diagonal with two states that are degenerate. Since Z_4 is abelian this degeneration can be broken only by completely breaking Z_4 . In this case $U_{G'}$ of eq. (7) completely arbitrary and exact TBM cannot be obtained.

So far we have considered all the possible cases in which the subgroup fixed in the charged lepton sector gives rise to a non diagonal structure to the charged lepton mass matrix M_l . The last case involving Z_4 gives rise to a diagonal $M_l M_l^\dagger$ but with two degenerate states. We could ask if there is any way to realize a diagonal M_l with three different mass eigenvalues. Indeed this is easily realized breaking S_4 to $Z_2 \times Z_2$ corresponding to the elements S^2 and $T^2 S^2 T$ of the class $\mathcal{C}_2$. If the charged lepton mass matrix is diagonal all the mixing structure arise by the neutrino sector. Therefore the last symmetry breaking pattern we are going to consider is $(Z_2 \times Z_2, S_3)$.

In this last case we break S_4 into S_3 in the neutrino sector. Following the same analysis that brought to eq. (10) we have

$$U_{TBM}^T m^\nu U_{TBM} = m_{S_3}^\nu, \quad (14)$$

that means $U_{lep} = U_{TBM}$ being the charged lepton mass matrix diagonal. At this point we have to face off a further problem: when S_4 is broken to S_3 the triplet L splits in a singlet plus a doublet. If S_3 is unbroken the two states in the doublet are degenerate in contrast with experimental data. Therefore we should identify a way of breaking S_3 without affecting the mixing rotation of the neutrino mass matrix. To keep us as general as possible, consider $m_{S_3}^\nu$ obtained once $S_4 \rightarrow S_3$. If S_3 is unbroken we have $m_{S_3}^\nu = \text{Diag}(m_1, m_0, m_0)$.

Suppose now that the singlet and the doublet with respect to S_3 behave as two independent sectors in such a way that S_3 is preserved in the singlet sector while is broken in the doublet one¹. By imposing these conditions we discover that $m_{S_3 \text{ broken}}^\nu$ has the following expression

$$m_{S_3 \text{ broken}}^\nu = \begin{pmatrix} m_1 & 0 & 0 \\ 0 & b_1 & b_2 \\ 0 & b_2 & b_3 \end{pmatrix}. \quad (15)$$

Finally let us impose that S_3 is not completely broken in the doublet sector but it is broken to its subgroup Z_2 identified by the S_3 generator S . This generator coincides with the S generator of the doublet representation of S_4 given in eq. (5). In this case it is possible to show that $m_{S_3 \text{ broken}}^\nu = \text{Diag}(m_1, m_2, m_3)$ and the lepton mixing matrix is still given by U_{TBM} .

We have seen that on the basis of theoretical considerations based on the subgroups of S_4 , the flavor symmetry S_4 has two symmetry breaking patterns giving exact TBM in the lepton sector. In the next section we will present a model realization for each breaking pattern. In the last section we build the corresponding supepotential responsible for the correct S_4 symmetry breaking patterns.

¹ From the point of view of model realization this assumption is not different by assuming that S_4 is broken to different subgroups in the charged lepton sector and in the neutrino one. Indeed we will see in sec. III B how singlet and doublet sectors can be easily separated.

III. MODEL REALIZATION

A. Model I : $S_4 \rightarrow Z_3$ & $S_4 \rightarrow Z_2$

The first model we consider reproduces TBM through the breaking of S_4 into Z_3 and Z_2 in the charged lepton and neutrino sector respectively. We assume our model to be supersymmetric. Matter and scalar supermultiplets are reported in tab. I. The scalar supermultiplets charged under S_4 , that in the following we will identify as flavons, are electroweak $SU(2) \times U(1)$ singlets. Therefore the Yukawa superpotential $\mathcal{W}_Y$ of eq. (16) includes effective operators of order 4. Λ is the cutoff of the model and an extra Z_5 symmetry has been introduced to separate the charged lepton sector from the neutrino one. In tab. I we have omitted the supermultiplets $\hat{H}^u$ and $\hat{\Phi}$, doublet and triplet of $SU(2)$ respectively, necessary to give mass to the up-quarks and to cancel anomalies in a realistic model.

	$\hat{L}$	$\hat{E}^c$	$\hat{H}^d$	$\hat{\Phi}$	$\hat{\sigma}$	$\hat{\phi}_1$	$\hat{\phi}_2$	$\hat{\Delta}$
$SU(2)$	2	1	2	3	1	1	1	1
S_4	3 ₁	3 ₁	1	1	1	3 ₁	3 ₂	3 ₁
Z_5	1	ω_5^4	1	1	ω_5	ω_5	ω_5	1

TABLE I: Matter and scalar content of model I. The lepton mixing matrix is TB.

The full leading order $S_4 \times Z_5$ Yukawa superpotential $\mathcal{W}_Y$ is given by

$$\mathcal{W}_Y = \frac{1}{\Lambda} y_0 (\hat{L} \hat{E}^c)_1 \hat{\sigma} \hat{H}^d + \frac{1}{\Lambda} y_s (\hat{L} \hat{E}^c)_{3_1} \hat{\phi}_1 \hat{H}^d + \frac{1}{\Lambda} y_a (\hat{L} \hat{E}^c)_{3_2} \hat{\phi}_2 \hat{H}^d + y'_1 (\hat{L} \hat{L})_1 \hat{\Phi} + \frac{1}{\Lambda} y'_2 (\hat{L} \hat{L})_{3_1} \hat{\Delta} \hat{\Phi}. \quad (16)$$

When the S_4 triplet and doublet flavons align as

$$\langle \phi_1 \rangle \sim \langle \phi_2 \rangle \sim (1, 1, 1) \quad \langle \Delta \rangle \sim (1, 0, 0), \quad (17)$$

the charged lepton and neutrino mass matrices present the usual forms

$$M_l = \begin{pmatrix} h_0 & h_1 & h_2 \\ h_2 & h_0 & h_1 \\ h_1 & h_2 & h_0 \end{pmatrix} \quad m^\nu = \begin{pmatrix} a & 0 & 0 \\ 0 & a & b \\ 0 & b & a \end{pmatrix} \quad (18)$$

that satisfy

$$U_\omega M_l U_\omega^\dagger = M_l^{diag}, \quad U_\nu^T m_\nu U_\nu = m_\nu^{diag}, \quad (19)$$

with U_ω and U_ν given in eq. (2) and eq. (12) respectively. TBM is obtained as usual by $U_{TB} = U_\omega U_\nu$. The mass eigenvalues for the charged lepton are given by

$$m_e = h_0 + h_1 + h_2, \quad m_\mu = h_0 + h_1 \omega^2 + h_2 \omega, \quad m_\tau = h_0 + h_1 \omega + h_2 \omega^2, \quad (20)$$

and for the neutrino by $(a + b, a, b - a)$. By assuming that the flavon vevs are of order $\sim \lambda^2 \Lambda$ with λ the Cabibbo angle, the deviations from TBM induced by the next to leading order corrections to the Yukawa superpotential slightly modify lepton mixing keeping it still in agreement with neutrino data. Notice that the vev alignments

$$\langle \phi_1 \rangle \sim \langle \phi_2 \rangle \sim (1, 1, 1) \quad (21)$$

preserves the Z_3 subgroup of S_4 associated to the element T , while the vev alignments

$$\langle \varphi \rangle \sim (0, 1) \quad \langle \Delta \rangle \sim (1, 0, 0), \quad (22)$$

preserves the Z_2 associated to the element TST that in the doublet and triplet representation reads respectively as

$$TST = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad TST = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix} \quad (23)$$

B. Model II : $S_4 \rightarrow S_3$

The second model we describe realizes TBM through the sequential breaking of S_4 into S_3 and then into Z_2 in the neutrino sector and the breaking of S_4 into two different $Z_2 \times Z_2$ in the charged lepton sector. The step through S_3 is crucial : if we broke S_4 directly into Z_2 in the neutrino sector we would find a generic neutrino mass matrix $\mu - \tau$ invariant not diagonalized by TB. On the contrary, in the model that we present the step through S_3 leads to a neutrino mass matrix m^ν which is $\mu - \tau$ invariant and satisfy the relation $m_{11}^\nu = m_{22}^\nu + m_{33}^\nu - m_{13}^\nu$ that ensures TB diagonalization. We will see that the key ingredient in building the correct m^ν is the introduction of the right handed neutrinos transforming as a doublet of S_4 . As in the case of the model presented in sec. III A we assume our model be supersymmetric and the flavon supermultiplets electroweak singlets. Matter and scalar supermultiplets are reported in tab. II. As done in sec. III A we have omitted the supermultiplet $\hat{\Phi}$, triplet of $SU(2)$, necessary to cancel anomalies. Two extra discrete abelian symmetries, Z_3 and Z_5 , have been introduced in order to avoid interferences between the sectors.

	$\hat{L}$	$\hat{l}^c$	$\hat{N}^c$	$\hat{H}^u$	$\hat{H}^d$	$\hat{\Phi}$	$\hat{\Delta}$	$\hat{\sigma}$	$\hat{\phi}$	$\hat{\varphi}$
$SU(2)$	2	1	1	2	2	3	1	1	1	1
S_4	3_1	3_1	2	1	1	1	3_1	1	2	2
Z_3	ω^2	1	1	1	ω^2	ω	ω	1	1	1
Z_5	1	ω_5^3	1	1	1	1	1	ω_5^2	ω_5^2	1

TABLE II: Matter and scalar content of model II. The lepton mixing matrix is TB.

The full leading order $S_4 \times Z_3 \times Z_5$ invariant Yukawa superpotential is given by

$$\mathcal{W}_Y = \frac{1}{\Lambda} y_s (\hat{L} \hat{l}^c)_1 \hat{\sigma} \hat{H}^d + \frac{1}{\Lambda} y_d (\hat{L} \hat{l}^c)_2 \hat{\phi} \hat{H}^d + y_1 (\hat{L} \hat{L})_1 \hat{\Phi} + \frac{1}{\Lambda} y_2 (\hat{L} \hat{\Delta})_2 \hat{N}^c \hat{H}^u + M_d \hat{N}^c \hat{N}^c + \tilde{y}_N \hat{\varphi} \hat{N}^c \hat{N}^c, \quad (24)$$

where as usual Λ is the cutoff of the model and all the Yukawa terms are of order 4 with the exception of the ones involving right handed neutrinos. We assume that the flavons Δ and φ , triplet and doublet under S_4 respectively, align as

$$\langle \Delta \rangle \sim (1, 1, 1) \quad \langle \varphi \rangle \sim (0, 1). \quad (25)$$

The vev $\langle \Delta \rangle$ preserves S_3 as has been already discussed in sec. IIB. The vev $\langle \varphi \rangle$ preserves the S generators of S_3 that coincides with the S generator of S_4 of the doublet representation—eq. (5).

The doublet ϕ does not align and develops vev as $\langle \phi \rangle \sim (v_1, v_2)$ —this means that S_4 is broken to $Z_2 \times Z_2$ corresponding to the elements S^2 and $T S^2 T^2$ of $\mathcal{C}_2$ that in the 3_1 triplet representation read as

$$S^2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad T S^2 T^2 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}. \quad (26)$$

For the charged lepton sector we have

$$M_l = \frac{1}{\Lambda} v^d \begin{pmatrix} y'_s v_\sigma - 2y''_d v_2^\phi & 0 & 0 \\ 0 & y'_s v_\sigma + y'_d v_1^\phi + y''_d v_2^\phi & 0 \\ 0 & 0 & y'_s v_\sigma - y'_d v_1^\phi + y''_d v_2^\phi \end{pmatrix} \quad (27)$$

with $v_\sigma = \langle \sigma \rangle$ $v_{1,2}^\phi = \langle \phi_{1,2} \rangle$ $v^d = \langle H_0^d \rangle$, and the product factors absorbed in y'_s and y'_d, y''_d . The neutrino mass matrix gets contributions both from type I and type II see-saw

$$m^\nu = m_{LL} - m_D \cdot \frac{1}{M_N} \cdot m_D^T, \quad (28)$$

where $m_{LL} = y_1 v_\Phi \cdot \mathcal{I}$ with $v_\Phi = \langle \Phi \rangle$ and

$$m_D = y_2 \frac{v^\Delta}{\Lambda} v^u \begin{pmatrix} 0 & -2\sqrt{6} \\ 1/\sqrt{2} & 1/\sqrt{6} \\ -1/\sqrt{2} & 1/\sqrt{6} \end{pmatrix}, \quad M_N = \begin{pmatrix} M_d + V_\varphi & 0 \\ 0 & M_d - V_\varphi \end{pmatrix}, \quad (29)$$

with $v^u = \langle H_0^u \rangle$, $v_{1,2,3}^\Delta = v^\Delta$ and $V_\varphi = \tilde{g}_N \langle \varphi_2 \rangle / \sqrt{2}$. After the usual see-saw mechanism the majorana neutrino mass matrix is given by

$$m^\nu = \begin{pmatrix} a + \frac{2}{3}b & -\frac{1}{3}b & -\frac{1}{3}b \\ -\frac{1}{3}b & a + \frac{1}{6}b + \frac{1}{2}c & \frac{1}{6}b - \frac{1}{2}c \\ -\frac{1}{6}b & \frac{1}{6}b - \frac{1}{2}c & a + \frac{1}{6}b + \frac{1}{2}c \end{pmatrix}, \quad (30)$$

with

$$a = y_1 v_\Phi, \quad b = -y_2^2 \left(\frac{v^\Delta}{\Lambda} \right)^2 \frac{(v^u)^2}{M_d - V_\varphi}, \quad c = -y_2^2 \left(\frac{v^\Delta}{\Lambda} \right)^2 \frac{(v^u)^2}{M_d + V_\varphi}. \quad (31)$$

The neutrino mass matrix m^ν is diagonalized by TBM and its eigenvalues are $(a+b, a, a+c)$ that can accommodate experimental neutrino mass splitting data being expressed in terms of three independent combinations of the parameters of the model. As in the model discussed in sec. III A by assuming the flavon vevs of order $\sim \lambda^2 \Lambda$ next to leading order corrections to the Yukawa superpotential produce small deviations from TBM that are still compatible with neutrino data.

IV. REALIZING THE CORRECT VACUUM CONFIGURATIONS IN S_4

In the context of flavor model based on non abelian discrete symmetry the lepton TBM is obtained thanks to specific alignments of the flavons. The so-called alignment problem in A_4 and T' has been extensively discussed in [18, 21, 25]. Different strategies have been used: the introduction of soft breaking term of the flavor symmetry [25], the use of a continuous $U(1)_R$ symmetry [21] preserved by the scalar potential and the promotion of the model to a fifth dimension [18]. In the context of S_4 in [39] the flavon superpotential was softly broken to guarantee the desired vacuum configuration.

In S_4 as well as in A_4 and T' it is impossible to build a flavon superpotential that guarantees the alignments needed. In the next sections we will show that the extra discrete abelian symmetries introduced in sec. III to separate the two lepton sectors are sufficient to give the correct vacuum configurations.

A. Model I : minimization of the potential

	$\hat{\sigma}$	$\hat{\phi}_1$	$\hat{\phi}_2$	$\hat{\Delta}$	$\hat{\varphi}$	$\hat{\xi}$	$\hat{\eta}$
$SU(2)$	1	1	1	1	1	1	1
S_4	1	3_1	3_2	3_1	2	2	2
Z_5	ω_5	ω_5	ω_5	1	1	ω_5^3	ω_5^2

TABLE III: Scalar content of model I including the flavons that contribute to the mass matrix structures and the ones the drive the correct vacuum alignments, the driving fields.

The flavon potential is obtained by the following part of the full $S_4 \times Z_5$ superpotential

$$\begin{aligned} \mathcal{W}_Y = & M_{\xi\eta} \hat{\xi} \hat{\eta} + \lambda_{\xi\eta} \hat{\xi} \hat{\eta} \hat{\varphi} + \lambda_{\sigma\eta} \hat{\sigma} \hat{\eta} \hat{\eta} + \lambda_{\xi\phi_1} \hat{\xi} \hat{\phi}_1 \hat{\phi}_1 + \lambda_{\xi\phi_2} \hat{\xi} \hat{\phi}_2 \hat{\phi}_2 + \lambda_{\xi\phi_{12}} \hat{\xi} \hat{\phi}_1 \hat{\phi}_2 \\ & + M_\Delta \hat{\Delta} \hat{\Delta} + M_\varphi \hat{\varphi} \hat{\varphi} + \lambda_{\varphi\Delta} \hat{\Delta} \hat{\Delta} \hat{\varphi} + \lambda_\varphi \hat{\varphi} \hat{\varphi} \hat{\varphi} + \lambda_\Delta \hat{\Delta} \hat{\Delta} \hat{\Delta}. \end{aligned} \quad (32)$$

We assume that the flavor symmetry is broken in the SUSY limit and therefore the vacuum configuration is obtained solving the system $\partial \mathcal{W}_Y / \partial f_i = 0$, where f_i are the f components of the supermultiplets entering in eq. (32) and i runs on all the supermultiplets. By assuming the general vacuum configuration

$$\langle \Delta \rangle = (v_1^\Delta, v_2^\Delta, v_3^\Delta), \quad \langle \varphi \rangle = (v_1^\varphi, v_2^\varphi), \quad \langle \phi_1 \rangle = (v_1^\phi, v_2^\phi, v_3^\phi), \quad \langle \phi_2 \rangle = (u_1^\phi, u_2^\phi, u_3^\phi), \quad \langle \xi \rangle = (u_1^\xi, u_2^\xi), \quad \langle \eta \rangle = (z^\eta, z^\eta) \quad \langle \sigma \rangle = v_\sigma, \quad (33)$$

the set of equations is given by

$$\begin{aligned}
a) \quad \partial W / \partial f_1^\Delta &= \frac{2}{\sqrt{3}} M_\Delta v_1^\Delta - \frac{2}{\sqrt{3}} \lambda_{\Delta\varphi} v_1^\Delta v_2^\varphi + 2\lambda_\Delta v_2^\Delta v_3^\Delta = 0 \\
b) \quad \partial W / \partial f_2^\Delta &= \frac{2}{\sqrt{3}} M_\Delta v_2^\Delta + \frac{1}{\sqrt{3}} \lambda_{\Delta\varphi} v_2^\Delta (v_2^\varphi + \sqrt{3}v_1^\varphi) + 2\lambda_\Delta v_1^\Delta v_3^\Delta = 0 \\
c) \quad \partial W / \partial f_3^\Delta &= \frac{2}{\sqrt{3}} M_\Delta v_3^\Delta + \frac{1}{\sqrt{3}} \lambda_{\Delta\varphi} v_3^\Delta (v_2^\varphi - \sqrt{3}v_1^\varphi) + 2\lambda_\Delta v_1^\Delta v_2^\Delta = 0 \\
d) \quad \partial W / \partial f_1^\varphi &= \sqrt{2} M_\varphi v_1^\varphi + \frac{\lambda_{\xi\eta}}{2} (u_2^\xi z_1^\eta + u_1^\xi z_2^\eta) + \frac{\lambda_\Delta}{2} [(v_2^\Delta)^2 - (v_3^\Delta)^2] = 0 \\
e) \quad \partial W / \partial f_2^\varphi &= \sqrt{2} M_\varphi v_2^\varphi + \frac{\lambda_{\xi\eta}}{2} (u_1^\xi z_1^\eta - u_2^\xi z_2^\eta) + \frac{\lambda_\Delta}{2\sqrt{3}} [-2(v_1^\Delta)^2 + (v_2^\Delta)^2 + (v_3^\Delta)^2] = 0 \\
f) \quad \partial W / \partial f_1^\eta &= \frac{M_{\xi\eta}}{\sqrt{2}} u_1^\xi + \frac{\lambda_{\xi\eta}}{2} (v_1^\varphi u_2^\xi + v_2^\varphi u_1^\xi) + \sqrt{2} \lambda_{\sigma\eta} v_\sigma z_1^\eta = 0 \\
g) \quad \partial W / \partial f_2^\eta &= \frac{M_{\xi\eta}}{\sqrt{2}} u_2^\xi + \frac{\lambda_{\xi\eta}}{2} (v_1^\varphi u_1^\xi - v_2^\varphi u_2^\xi) + \sqrt{2} \lambda_{\sigma\eta} v_\sigma z_2^\eta = 0 \\
h) \quad \partial W / \partial f^\sigma &= \frac{\lambda_{\sigma\eta}}{\sqrt{2}} [(z_1^\eta)^2 + (z_2^\eta)^2] = 0 \\
i) \quad \partial W / \partial f_1^\xi &= \frac{1}{\sqrt{2}} M_{\xi\eta} z_1^\eta + \frac{1}{2} \lambda_{\xi\eta} (z_1^\eta v_2^\varphi + z_2^\eta v_1^\varphi) + \frac{1}{2} \lambda_{\xi\phi 1} [(v_2^\phi)^2 - (v_3^\phi)^2] + \frac{1}{2} \lambda_{\xi\phi 2} [(u_2^\phi)^2 - (u_3^\phi)^2] \\
&\quad + \frac{1}{2\sqrt{3}} \lambda_{\xi\phi 12} (2v_1^\phi u_1^\phi - v_2^\phi u_2^\phi - v_3^\phi u_3^\phi) = 0 \\
j) \quad \partial W / \partial f_2^\xi &= \frac{1}{\sqrt{2}} M_{\xi\eta} z_2^\eta + \frac{1}{2} \lambda_{\xi\eta} (z_1^\eta v_1^\varphi - z_2^\eta v_2^\varphi) + \frac{1}{2\sqrt{3}} \lambda_{\xi\phi 1} [-2(v_1^\phi)^2 + (v_2^\phi)^2 + (v_3^\phi)^2] \\
&\quad + \frac{1}{2\sqrt{3}} \lambda_{\xi\phi 2} [-2(u_1^\phi)^2 + (u_2^\phi)^2 + (u_3^\phi)^2] + \frac{1}{2} \lambda_{\xi\phi 12} (v_2^\phi u_2^\phi - v_3^\phi u_3^\phi) = 0 \\
k) \quad \partial W / \partial f_1^{\phi 1} &= \frac{1}{\sqrt{3}} (\lambda_{\xi\phi 12} u_1^\phi u_1^\xi - 2\lambda_{\xi\phi 1} u_2^\xi v_1^\phi) = 0 \\
l) \quad \partial W / \partial f_2^{\phi 1} &= u_1^\xi (\lambda_{\xi\phi 1} v_2^\phi - \frac{1}{2\sqrt{3}} \lambda_{\xi\phi 12} u_2^\phi) + u_2^\xi (\frac{\lambda_{\xi\phi 1}}{\sqrt{3}} v_2^\phi + \frac{1}{2} \lambda_{\xi\phi 12} u_2^\phi) = 0 \\
m) \quad \partial W / \partial f_3^{\phi 1} &= u_1^\xi (-\lambda_{\xi\phi 1} v_3^\phi - \frac{1}{2\sqrt{3}} \lambda_{\xi\phi 12} u_3^\phi) + u_2^\xi (\frac{\lambda_{\xi\phi 1}}{\sqrt{3}} v_2^\phi - \frac{1}{2} \lambda_{\xi\phi 12} u_2^\phi) = 0 \\
n) \quad \partial W / \partial f_1^{\phi 2} &= \frac{1}{\sqrt{3}} (\lambda_{\xi\phi 12} v_1^\phi u_1^\xi - 2\lambda_{\xi\phi 2} u_2^\xi u_1^\phi) = 0 \\
o) \quad \partial W / \partial f_2^{\phi 2} &= u_1^\xi (\lambda_{\xi\phi 2} u_2^\phi - \frac{1}{2\sqrt{3}} \lambda_{\xi\phi 12} v_2^\phi) + u_2^\xi (\frac{\lambda_{\xi\phi 1}}{\sqrt{3}} u_2^\phi + \frac{1}{2} \lambda_{\xi\phi 12} v_2^\phi) = 0 \\
p) \quad \partial W / \partial f_3^{\phi 2} &= u_1^\xi (-\lambda_{\xi\phi 2} u_3^\phi - \frac{1}{2\sqrt{3}} \lambda_{\xi\phi 12} v_3^\phi) + u_2^\xi (\frac{\lambda_{\xi\phi 2}}{\sqrt{3}} u_2^\phi - \frac{1}{2} \lambda_{\xi\phi 12} v_2^\phi) = 0
\end{aligned} \tag{34}$$

Eq. *h*) of eq. (34) implies $z_{1,2}^\eta = 0$. As first consequence we have that a possible solution of eqs. *f*) – *g*) and eqs. *k*) – *p*) is given by

$$(u_1^\xi, u_2^\xi) = (0, 0) \quad \text{and} \quad v_\sigma \neq 0. \tag{35}$$

By substituting $(z_1^\eta, z_2^\eta) = (0, 0)$, $(u_1^\xi, u_2^\xi) = (0, 0)$ and $v_\sigma \neq 0$ in the equations not yet solved it is easy to check that a possible solution for eqs. *a*) – *e*) is given by the vacuum configuration

$$\begin{aligned}
(v_1^\varphi, v_2^\varphi) &= (0, v^\varphi) \quad \text{with} \quad v^\varphi = \frac{M_\Delta}{\lambda_\Delta} \\
(v_1^\Delta, v_2^\Delta, v_3^\Delta) &= (v^\Delta, 0, 0) \quad \text{with} \quad v^\Delta = 6^{1/4} \frac{\sqrt{M_\varphi M_\Delta}}{\lambda_\Delta}.
\end{aligned} \tag{36}$$

Finally eqs. *i*) – *j*) are solved by the vacuum configuration

$$(v_1^\phi, v_2^\phi, v_3^\phi) = v^\phi (1, 1, 1) \quad \text{and} \quad (u_1^\phi, u_2^\phi, u_3^\phi) = u^\phi (1, 1, 1). \tag{37}$$

The solution found is not unique but can be stabilized once we add apposite SUSY soft breaking terms. In sec. III A we have assumed that the flavon vevs is of order $\lambda^2 \Lambda$. Therefore the next to leading order corrections to the Yukawa superpotential induced by the driving fields are sufficiently suppressed.

B. Model II : minimization of the potential

	$\hat{\Delta}$	$\hat{\sigma}$	$\hat{\phi}$	$\hat{\varphi}$	$\hat{\bar{\sigma}}$	$\hat{\xi}$	$\hat{\eta}$
$SU(2)$	1	1	1	1	1	1	1
S_4	3 ₁	1	2	2	1	1	1
Z_3	ω	1	1	1	1	ω	ω^2
Z_5	1	ω_5^2	ω_5^2	1	ω_5	1	1

TABLE IV: Scalar content of model I including both flavon and the driving field supermultiplets.

The flavon potential is obtained by the following part of the full superpotential

$$\begin{aligned} \mathcal{W} = & \lambda_{\Delta\xi} \hat{\xi} \hat{\Delta} \hat{\Delta} + \lambda_{\Delta} \hat{\Delta} \hat{\Delta} \hat{\Delta} + M_{\xi} \hat{\xi} \hat{\eta} + \lambda_{\xi} \hat{\xi} \hat{\xi} \hat{\xi} + \lambda_{\eta} \hat{\eta} \hat{\eta} \hat{\eta} \\ & + M_{\varphi} \hat{\varphi} \hat{\varphi} + \lambda_{\varphi} \hat{\varphi} \hat{\varphi} \hat{\varphi} + \lambda_{\phi} \hat{\sigma} \hat{\phi} \hat{\phi} + \lambda_{\sigma} \hat{\sigma} \hat{\sigma} \hat{\sigma}. \end{aligned} \quad (38)$$

By assuming the general vacuum configuration

$$\langle \Delta \rangle = (v_1^{\Delta}, v_2^{\Delta}, v_3^{\Delta}), \langle \varphi \rangle = (v_1^{\varphi}, v_2^{\varphi}), \langle \phi \rangle = (v_1^{\phi}, v_2^{\phi}), \langle \xi \rangle = v_{\xi}, \langle \eta \rangle = v_{\eta}, \langle \sigma \rangle = v_{\sigma}, \langle \bar{\sigma} \rangle = v_{\bar{\sigma}}, \quad (39)$$

the minimization of the scalar potential obtained in the SUSY limit gives the following set of equations

$$\begin{aligned} \partial \mathcal{W}_Y / \partial f_1^{\Delta} &= \sqrt{2} \lambda_{\Delta\xi} v_{\xi} v_1^{\Delta} + \sqrt{3} \lambda_{\Delta} v_2^{\Delta} v_3^{\Delta} = 0 \\ \partial \mathcal{W}_Y / \partial f_2^{\Delta} &= \sqrt{2} \lambda_{\Delta\xi} v_{\xi} v_2^{\Delta} + \sqrt{3} \lambda_{\Delta} v_1^{\Delta} v_3^{\Delta} = 0 \\ \partial \mathcal{W}_Y / \partial f_3^{\Delta} &= \sqrt{2} \lambda_{\Delta\xi} v_{\xi} v_3^{\Delta} + \sqrt{3} \lambda_{\Delta} v_1^{\Delta} v_2^{\Delta} = 0 \\ \partial \mathcal{W}_Y / \partial f^{\xi} &= \sqrt{3} \lambda_{\Delta\xi} [(v_1^{\Delta})^2 + (v_2^{\Delta})^2 + (v_3^{\Delta})^2] + M_{\xi} v_{\eta} + 3 \lambda_{\xi} v_{\xi}^2 = 0 \\ \partial \mathcal{W}_Y / \partial f^{\eta} &= M_{\xi} v_{\xi} + 3 \lambda_{\eta} v_{\eta}^2 = 0 \\ \partial \mathcal{W}_Y / \partial f_1^{\varphi} &= \sqrt{2} M_{\varphi} v_1^{\varphi} + 3 \lambda_{\varphi} v_1^{\varphi} v_2^{\varphi} = 0 \\ \partial \mathcal{W}_Y / \partial f_2^{\varphi} &= \sqrt{2} M_{\varphi} v_2^{\varphi} + \frac{3}{2} \lambda_{\varphi} [(v_1^{\varphi})^2 - (v_2^{\varphi})^2] = 0 \\ \partial \mathcal{W}_Y / \partial f_1^{\phi} &= \sqrt{2} \lambda_{\phi} v_1^{\phi} v_{\bar{\sigma}} = 0 \\ \partial \mathcal{W}_Y / \partial f_2^{\phi} &= \sqrt{2} \lambda_{\phi} v_2^{\phi} v_{\bar{\sigma}} = 0 \\ \partial \mathcal{W}_Y / \partial f^{\sigma} &= 2 \lambda_{\bar{\sigma}} v_{\sigma} v_{\bar{\sigma}} = 0 \\ \partial \mathcal{W}_Y / \partial f^{\bar{\sigma}} &= \frac{1}{\sqrt{2}} \lambda_{\phi} [(v_1^{\phi})^2 + (v_2^{\phi})^2] + \lambda_{\bar{\sigma}} v_{\sigma}^2 = 0. \end{aligned} \quad (40)$$

Discarding for the triplet and the doublets the trivial solutions that do not break S_4 , the solution of the system of eq. (40) is given by the following vacuum configuration

$$\begin{aligned} v_1^{\Delta} = v_2^{\Delta} = v_3^{\Delta} = v^{\Delta} \quad & \text{with} \quad v^{\Delta} = \sqrt{2} \frac{\lambda_{\Delta\xi} \lambda_{\eta}}{\lambda_{\Delta}} \frac{v_{\eta}^2}{M_{\xi}} \\ v_{\xi} = -3 \lambda_{\eta} \frac{v_{\eta}^2}{M_{\xi}} \quad & \text{with} \quad v_{\eta}^3 = -M_{\xi}^3 \frac{\lambda_{\Delta}^2}{\lambda_{\eta}^2 (2\sqrt{3} \lambda_{\Delta\xi}^3 + 27 \lambda_{\xi} \lambda_{\Delta}^2)} \\ (v_1^{\varphi}, v_2^{\varphi}) \neq (0, 0) \quad & \text{with} \quad \begin{cases} (0, \frac{2\sqrt{2}}{3} \frac{M_{\varphi}}{\lambda_{\varphi}}) \\ (\sqrt{\frac{2}{3}} \frac{M_{\varphi}}{\lambda_{\varphi}}, -\frac{\sqrt{2}}{3} \frac{M_{\varphi}}{\lambda_{\varphi}}) \\ (-\sqrt{\frac{2}{3}} \frac{M_{\varphi}}{\lambda_{\varphi}}, -\frac{\sqrt{2}}{3} \frac{M_{\varphi}}{\lambda_{\varphi}}) \end{cases} \\ v_{\sigma}^2 = -\frac{1}{\sqrt{2}} \frac{\lambda_{\phi}}{\lambda_{\sigma}} [(v_1^{\phi})^2 + (v_2^{\phi})^2] \neq 0 \text{ and } v_{\bar{\sigma}} = 0. \end{aligned} \quad (41)$$

The three solutions corresponding to $\langle\varphi\rangle$ are degenerate and corresponding to the breaking of S_3 to its 3 different Z_2 subgroups. Through appropriate choices of soft terms that break the discrete abelian symmetry Z_3 and Z_5 and not S_4 we can stabilize as absolute minimum the vacuum configuration $\langle\varphi\rangle \sim (0, 1)$.

V. CONCLUSION

In this paper we have discussed the idea that S_4 is the minimal discrete non abelian group naturally related to TBM in the lepton sector. We have shown that S_4 can yield exact TBM according to a general group theory analysis and we have presented two explicit model realizations of how TBM can be obtained in S_4 once the basis of its generators are fixed. In addition we have provided a detailed study of the corresponding scalar potentials. The two models require two triplets with different vev alignments. For each model we have built a potential that in the SUSY limit contains the minimum required. The problem of the triplet and doublet alignments is solved in a more economical way than in models based on A_4 [14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29]. To separate the charged lepton sector from the neutrino one we have introduced extra abelian symmetries. The construction of the potentials have not required additional symmetries than such extra abelian symmetries, but just the addition of “driving” fields that do not enter in the Yukawa part. We have studied neither the quark sector nor the possibility to embed such a model in a GUT theory. We leave these subjects for a future publication. It is worth to mention that in S_4 there is more freedom to generate the mixing in the quark sector than in A_4 . Indeed the doublet irreducible representation could play an important role as happens in T' [30].

Acknowledgments

We thank L. Merlo for very useful comments and discussions. Work supported by MEC grant FPA2008-00319/FPA, by EC RTN network MRTN-CT-2004-503369 and by Generalitat Valenciana ACOMP06/154.

-
- [1] P. F. Harrison, D. H. Perkins, and W. G. Scott, Phys. Lett. **B530**, 167 (2002), hep-ph/0202074.
 - [2] M. Maltoni, T. Schwetz, M. A. Tortola, and J. W. F. Valle, New J. Phys. **6**, 122 (2004), hep-ph/0405172.
 - [3] F. Feruglio and Y. Lin, Nucl. Phys. **B800**, 77 (2008), 0712.1528.
 - [4] W. Grimus and L. Lavoura, JHEP **08**, 013 (2005), hep-ph/0504153.
 - [5] M. Mitra and S. Choubey (2008), 0806.3254.
 - [6] A. Mondragon, M. Mondragon, and E. Peinado, AIP Conf. Proc. **1026**, 164 (2008), 0712.2488.
 - [7] C.-Y. Chen and L. Wolfenstein, Phys. Rev. **D77**, 093009 (2008), 0709.3767.
 - [8] Y. Koide, Eur. Phys. J. **C50**, 809 (2007), hep-ph/0612058.
 - [9] R. N. Mohapatra, S. Nasri, and H.-B. Yu, Phys. Lett. **B639**, 318 (2006), hep-ph/0605020.
 - [10] S. Morisi and M. Picariello, Int. J. Theor. Phys. **45**, 1267 (2006), hep-ph/0505113.
 - [11] F. Caravaglios and S. Morisi (2005), hep-ph/0503234.
 - [12] P. F. Harrison and W. G. Scott, Phys. Lett. **B557**, 76 (2003), hep-ph/0302025.
 - [13] S.-L. Chen, M. Frigerio, and E. Ma, Phys. Rev. **D70**, 073008 (2004), hep-ph/0404084.
 - [14] E. Ma and G. Rajasekaran, Phys. Rev. **D64**, 113012 (2001), hep-ph/0106291.
 - [15] K. S. Babu, E. Ma, and J. W. F. Valle, Phys. Lett. **B552**, 207 (2003), hep-ph/0206292.
 - [16] M. Hirsch et al., Phys. Rev. **D69**, 093006 (2004), hep-ph/0312265.
 - [17] E. Ma, Phys. Rev. **D70**, 031901 (2004), hep-ph/0404199.
 - [18] G. Altarelli and F. Feruglio, Nucl. Phys. **B720**, 64 (2005), hep-ph/0504165.
 - [19] S.-L. Chen, M. Frigerio, and E. Ma, Nucl. Phys. **B724**, 423 (2005), hep-ph/0504181.
 - [20] A. Zee, Phys. Lett. **B630**, 58 (2005), hep-ph/0508278.
 - [21] G. Altarelli and F. Feruglio, Nucl. Phys. **B741**, 215 (2006), hep-ph/0512103.
 - [22] B. Adhikary, B. Brahmachari, A. Ghosal, E. Ma, and M. K. Parida, Phys. Lett. **B638**, 345 (2006), hep-ph/0603059.
 - [23] J. W. F. Valle, J. Phys. Conf. Ser. **53**, 473 (2006), hep-ph/0608101.
 - [24] B. Adhikary and A. Ghosal, Phys. Rev. **D75**, 073020 (2007), hep-ph/0609193.
 - [25] E. Ma, Mod. Phys. Lett. **A22**, 101 (2007), hep-ph/0610342.
 - [26] G. Altarelli, F. Feruglio, and Y. Lin, Nucl. Phys. **B775**, 31 (2007), hep-ph/0610165.
 - [27] M. Hirsch, A. S. Joshipura, S. Kaneko, and J. W. F. Valle (0300), hep-ph/0703046.
 - [28] G. Altarelli (0500), arXiv:0705.0860 [hep-ph].
 - [29] F. Bazzocchi, S. Kaneko, and S. Morisi, JHEP **03**, 063 (2008), 0707.3032.

- [30] F. Feruglio, C. Hagedorn, Y. Lin, and L. Merlo, Nucl. Phys. **B775**, 120 (2007), hep-ph/0702194.
- [31] A. Aranda, Phys. Rev. **D76**, 111301 (2007), 0707.3661.
- [32] D. A. Eby, P. H. Frampton, and S. Matsuzaki (2008), 0810.4899.
- [33] P. H. Frampton and T. W. Kephart, Int. J. Mod. Phys. **A10**, 4689 (1995), hep-ph/9409330.
- [34] G. J. Ding, Phys. Rev. **D 78** (2008) 036011.
- [35] E. Ma, Phys. Lett. **B632**, 352 (2006), hep-ph/0508231.
- [36] C. Hagedorn, M. Lindner, and R. N. Mohapatra, JHEP **06**, 042 (2006), hep-ph/0602244.
- [37] Y. Cai and H.-B. Yu, Phys. Rev. **D74**, 115005 (2006), hep-ph/0608022.
- [38] H. Zhang, Phys. Lett. **B655**, 132 (2007), hep-ph/0612214.
- [39] Y. Koide, JHEP **08**, 086 (2007), 0705.2275.
- [40] C. Luhn, S. Nasri, and P. Ramond, J. Math. Phys. **48**, 073501 (2007), hep-th/0701188.
- [41] I. de Medeiros Varzielas, S. F. King, and G. G. Ross, Phys. Lett. **B648**, 201 (2007), hep-ph/0607045.
- [42] E. Ma, Phys. Lett. **B660**, 505 (2008), 0709.0507.
- [43] E. Ma, Mod. Phys. Lett. **A21**, 1917 (2006), hep-ph/0607056.
- [44] P. H. Frampton and T. W. Kephart, Phys. Rev. **D64**, 086007 (2001), hep-th/0011186.
- [45] F. Bazzocchi, M. Frigerio, and S. Morisi (2008), 0809.3573.
- [46] C. S. Lam, Phys. Rev. **D 78**, 073015 (2008), 0809.1185.