

Generalized Seesaw Mechanism of Neutrino and Bose-Einstein Condensation in the Modified O'Raifeartaigh Model

Tadafumi Ohsaku

Institut für Theoretische Physik, Universität zu Köln, 50937 Köln, Germany

(Dated: February 21, 2024)

The modified O'Raifeartaigh model from the context of the generalized seesaw mechanism of neutrino mass is investigated. In our evaluation of effective potentials of the theory, both the component field and the superspace formalisms to approach the problem are presented. In the component field formalism, we take into account the Bose-Einstein condensates in the scalar sector by the method of many-boson theory, i.e. we consider both the condensates and the Hartree-Fock-Bogoliubov-type self-energies of quantum fluctuations. The diagonalization of the mass matrix of the fermion sector gives the same functional forms of the mass eigenvalues in the generalized seesaw mechanism. The stability condition in the vicinity of the classical vacuum which shows the generalized seesaw situation is obtained by the examination of the mass eigenvalues of the scalar sector of the model. The superspace formalism will be devoted to a comparison between its result with that of the component field formalism. (Keywords: Supersymmetric Effective Theories, Neutrino Physics, Supersymmetry Breaking, Nonperturbative Effects)

PACS numbers: 11.30.Pb, 11.30.Qc, 14.60.Pq, 14.80.Mz

I. INTRODUCTION

The standard model of elementary particles is the most important achievement in modern physics, and still it gives us the horizon of particle phenomenology [1]. Supersymmetry (SUSY) [2-22] can be understood as one of the ways toward "beyond the standard model", from a viewpoint of particle phenomenology. In such a supersymmetric approach, a theory has both fermionic and bosonic degrees of freedom, and they interact with each other under a supersymmetric manner. Dynamics of interacting boson gases quite often show Bose-Einstein condensation (BEC) as a universal phenomenon. BEC was first found by Bose [23] and Einstein [24], and theory of BEC of an interacting nonrelativistic boson gas was constructed firstly by Bogoliubov [25]. The Bogoliubov theory has a very universal character, and it is the case that the theory can be applied to various interacting boson systems. Moreover, methods and concepts of BEC and superfluidity of boson gases are useful to examine/understand an interacting fermion system, for example, BCS (Bardeen-Cooper-Schrieffer) superconductivity [26] or chiral condensations in the Namu-Jona-Lasinio model (NJL) [27,28] and quantum chromodynamics (QCD) [29], from the context of spontaneous symmetry breakings. On the other hand, a SUSY multiplet must be broken in a model for phenomenology because we have not yet found any superpartner. Due to the (perturbative) nonrenormalization theorem, a SUSY breaking cannot take place in a perturbation theory, and it should be realized in a nonperturbative manner, i.e. a spontaneous SUSY breaking. A lot of modern particle theoreticians consider that a dynamical symmetry breaking is phenomenologically preferred for a SUSY breakdown [3,4,5,7,9,10,12].

The most important problem in modern particle physics which has been found by experimental results is the origin of masses, their hierarchy, and flavor violations of particles. Recent experimental observations confirmed that neutrinos should have very tiny masses, and the seesaw mechanism is one of candidates for providing an explanation to neutrino masses [30-34]. Hence, it is an interesting issue to make a SUSY model which will show a seesaw mechanism. In the ordinary seesaw mechanism, neutrino has both a Dirac and a right-handed Majorana mass terms. The references (35) and (36) discussed a generalization of the ordinary seesaw mechanism, added a (very tiny) left-handed Majorana mass, and some interesting results were obtained. It is well-known fact that the O'Raifeartaigh model breaks SUSY at its tree level [14]. Recently, the modified O'Raifeartaigh model has been examined in the context of meta-stable SUSY breaking [10-13]. Reference (11) gives mass eigenvalues of scalars and spinors: In fact the eigenvalues take quite similar structure with that of the generalized seesaw mechanism [35,36]. The purpose of this paper is to examine the modified O'Raifeartaigh model under the context of the generalized seesaw mechanism of neutrinos.

This paper is organized as follows. In Sec. II, we investigate the generalized seesaw mechanism of the so-called modified O'Raifeartaigh model in the component field formalism. It is suitable to employ several many-body-theoretical techniques in the component field formalism though it becomes more lengthy than the superspace formalism. While, we have to use the notion of component fields at some discussions also in the superspace formalism, especially if we wish to take into account BEC in the scalar sector of the theory. After introducing the modified O'Raifeartaigh model, we shortly discuss its symmetry property and the classical minimum. We consider it might be possible that

the solution of the classical minimum shows the generalized seesaw mass relation. Then, we employ the many-body-theoretical technique to take into account BEC in the scalar sector. By these preparation, the one-loop effective potential is calculated, and stability around the classical minimum will be investigated. Possibility of SUSY breakdown around the classical minimum also be examined. In a one-loop effective potential calculation, the loop expansion must converge rapidly enough, and thus the vacuum of a theory should have a semiclassical nature, will not obtain a radical modification by possible quantum corrections. This must be the case in our calculation, and thus we consider the situation where quantum corrections around a classical minimum are small. For comparison/supplement to the result of the component field formalism, a calculation of the one-loop effective potential in the superspace formalism is given in Sec. III. The summary and conclusion of this work is presented in Sec. IV.

We will follow the textbook of Wess and Bagger for the spinor algebra, gamma matrices and metric conventions throughout this paper [2]. (For example, the metric is $\eta = \text{diag}(-1; 1; 1; 1)$.)

II. COMPONENT FIELD FORMALISM

A. The Classical Solution

Our starting point is the following Lagrangian of the modified O'Raifeartaigh model [10-13] of three chiral matter fields:

$$\begin{aligned} \mathcal{L} = & X^\dagger X + \frac{Y}{+} + + Y \\ & + fX + \frac{g}{2}X + + + m_D + + + m_L \\ & + f^\dagger X^\dagger Y + \frac{g^\dagger}{2}X^\dagger Y + + + m_D^\dagger + + + m_L^\dagger : \end{aligned} \quad (1)$$

Here, X , Y are chiral (+; right, -; left) superfields. We regard X as neutrino superfields, m_D and m_L denote a Dirac and a left-handed Majorana mass parameters, respectively. If X takes a (very large) VEV compared with m_D and m_L , then the theory may show a seesaw-type situation in the mass matrix eigenvalues of its fermion sector. The usual (ordinary) seesaw situation will be achieved by $m_L \neq 0$. The mass dimensions of f and g become as follows: $f; [m_{\text{ass}}]^2, g; [m_{\text{ass}}]^0$. We consider the following global $U(1)_V$ (gauge) and $U(1)_A$ (chiral) transformations:

$$U(1)_V : + \rightarrow e^{i\nu} +; - \rightarrow e^{-i\nu} -; U(1)_A : + \rightarrow e^{i\alpha} +; - \rightarrow e^{-i\alpha} -; \nu; \alpha \in \mathbb{R} : \quad (2)$$

The Majorana mass term of the mass parameter m_L explicitly breaks both of these global symmetries. We can choose the charge of X to keep the coupling term $\frac{g}{2}X +$ invariant under these transformations as

$$U(1)_V : X \rightarrow e^{2i\nu} X; U(1)_A : X \rightarrow e^{2i\alpha} X : \quad (3)$$

The terms fX and $f^\dagger X^\dagger Y$ also explicitly break these global $U(1)$ symmetries. The Majorana mass term will break the $U(1)_R$ symmetry under the following charge assignment of the superfields X and Y [10-13]:

$$U(1)_R : - \rightarrow e^{i_R} -; + \rightarrow e^{i_R} +; X \rightarrow e^{2i_R} X; Y \rightarrow e^{i_R} Y; \nu; \alpha \in \mathbb{R} : \quad (4)$$

Another R -charge assignment is also possible:

$$U(1)_R : X \rightarrow X; + \rightarrow e^{i_R} +; - \rightarrow e^{i_R} - : \quad (5)$$

In this case, the R -symmetry will be restored at the limit $f \rightarrow 0$. Therefore, the term fX and the Majorana mass term are incompatible with respect to the $U(1)_R$ symmetry. As a result, there is no global $U(1)$ symmetry in our theory. The absence of R -axion in the modified O'Raifeartaigh model is discussed in Refs. [11,12] from the context of meta-stable SUSY breaking, and it is phenomenologically favorable. The ordinary O'Raifeartaigh model corresponds to the case $m_L = m_L^\dagger = 0$, it has an R -symmetry, and SUSY is broken at the tree level [12,14]. In the ordinary O'Raifeartaigh model, the classical solution becomes $+ = - = 0$ with $X = \text{arbitrary}$, and SUSY is spontaneously broken in the vacuum. The one-loop effective potential of the O'Raifeartaigh model was calculated in Ref. [15]. In that calculation, the degeneracy of vacua is lifted by the one-loop correction, and the origin of the potential becomes the only ground state. There is a Z_2 symmetry under $X \rightarrow -X$ in (1). The tree level part of the Lagrangian

will be obtained from the scalar potential by employing the Euler-Lagrange equations of the auxiliary fields of chiral multiplets:

$$V^{\text{tree}}[\phi; \chi] = F_X^2 + F_+^2 + F_-^2 = f + \frac{g}{2} \phi_+^2 + g \chi_+ + m_D \phi_+^2 + m_D \phi_-^2 + 2m_L \phi_+^2 : \quad (6)$$

In literature, the classical solution of V^{tree} is given as follows [??]

$$\phi_+^{\text{classical}} = \frac{m_D^2}{2gm_L}; \quad \phi_-^{\text{classical}} = \frac{2f}{g}; \quad \chi^{\text{classical}} = \frac{m_D}{2m_L} \frac{2f}{g} : \quad (7)$$

Here we have assumed that m_D, m_L, g and f are real valued to obtain the classical minimum. When these parameters are real, the classical solution shows the spontaneous Z_2 symmetry breakdown. All of the VEVs of the solution will go to infinity at the limit $g \rightarrow 0$, and $g = 0$ is a singular point for the solution. (This expression of the classical minimum of our model has a similarity with the classical solution of a Ginzburg-Landau-type ϕ^4 model which describes the low-energy property of the Ising ferromagnet [37]. Hence V^{tree} seems to have a relation with the Ising ferromagnet.) V^{tree} vanishes at the classical minimum and the $N = 1$ SUSY of this model is unbroken at the classical level [11,12]. We will see quantum corrections to the classical solution through the following loop expansion calculation, though we mainly investigate a possibility of the generalized seesaw situation in the vicinity of the classical solution in this work. As mentioned above, the ordinary O'Raifeartaigh model gives $h_X = i$ arbitrary at its classical solution with broken SUSY, and the one-loop correction gives the unique vacuum as the origin of the potential (the origin of (6) gives a finite energy and not supersymmetric). By introducing the left-handed Majorana mass term in (1), $\phi_+^{\text{classical}}$ has obtained the explicit expression given in (7) and then we can discuss the strength of the VEV $h_X = i$ which would give a right-handed Majorana mass parameter with some dynamics of our theory: If we set $m_L = 0$ from the beginning of our model, we cannot obtain an explicit expression for $h_X = i$ (at least at the classical level). This is crucial in the context of this work. The ordinary seesaw mechanism (the case $m_L = 0$) cannot be considered by the ordinary O'Raifeartaigh model. In this paper, we examine (i) when the classical solution can give the generalized seesaw situation, (ii) how the vicinity of the classical solution in the one-loop potential is stable and robust, (iii) whether the vicinity of the classical solution in the one-loop potential breaks SUSY or not. Usually, SUSY is broken if there is an R-symmetry in a theory, while SUSY will be kept if an R-symmetry is broken.

After eliminating the auxiliary fields of X and ϕ and perform integrations of Grassmann coordinates, one finds the expression of L in terms of component fields as follows:

$$L = \int d^4x \left[\int d^2\theta \left(\frac{1}{2} \phi_+^2 + g \chi_+ + m_D \phi_+^2 + m_D \phi_-^2 + 2m_L \phi_+^2 \right) + \int d^2\bar{\theta} \left(\frac{1}{2} \phi_-^2 + f + \frac{g}{2} \phi_+^2 + g \chi_+ + m_D \phi_+^2 + m_D \phi_-^2 + 2m_L \phi_+^2 \right) \right] : \quad (8)$$

The phases of mass parameters and χ are defined as

$$\chi = j_X e^{i\alpha_X}; \quad m_D = j_D e^{i\alpha_D}; \quad m_L = j_L e^{i\alpha_L}; \quad \alpha_X, \alpha_D, \alpha_L \in \mathbb{R} : \quad (9)$$

We can absorb only two of these phases α_X, α_D and α_L by a redefinition of fields. Hereafter, we set $m_D = j_D$ and $m_L = j_L$ by a field redefinition while keeping the phase degree of freedom of χ without loss of generality. Later, we will observe that mass eigenvalues of scalars and spinors are functions of α_X . In principle, if we take into account phase degrees of freedom of ϕ and χ , the classical solution of V^{tree} becomes ($f^y = f, g^y = g, m_D^y = m_D, m_L^y = m_L$ are imposed)

$$\begin{aligned} j_+ j_+ &= \frac{2f^y}{g} \frac{1}{\cos(2\alpha_X)} \frac{1}{\cos^2(2\alpha_X)} \frac{1}{1}; \\ j_+ j_- &= \frac{m_D j_+ j_-^h}{2m_L} \cos(\alpha_X + \alpha_D) \frac{1}{\cos^2(\alpha_X + \alpha_D)} \frac{1}{1}; \\ j_X j_+ &= \frac{m_D j_+ j_+^h}{g j_+ j_+} \cos(\alpha_X + \alpha_D) \frac{1}{\cos^2(\alpha_X + \alpha_D)} \frac{1}{1}; \quad = j_+ e^{i\alpha_X} : \end{aligned} \quad (10)$$

By the HFB approximation, the cubic interactions of fluctuations will also be dropped from our Lagrangian. Hence we get

$$\frac{\eta^2}{4} j_+^4 = \frac{\eta^2}{4} [4J_1 + 4j_+^c j_+^{\sim y} \tilde{\nu}_+ + J_2^y + (\frac{c^y}{+})^2 \tilde{\nu}_+ \tilde{\nu}_+ + J_2 + (\frac{c}{+})^2 \tilde{\nu}_+^y \tilde{\nu}_+^y + j_+^c j_+^4]; \quad (17)$$

and

$$\begin{aligned} \eta^2 j_x^2 j_+^2 j_+^2 = & \eta^2 [j_x^c j_+^c j_+^2 + (j_x^c j_+^2 + K_1) \tilde{\nu}_+^2 + (j_+^c j_+^2 + J_1) \tilde{\nu}_x^2 \\ & + (\frac{c^y}{x} \frac{c}{+} + K_3) \tilde{\nu}_+^y \tilde{\nu}_x + (\frac{c^y}{+} \frac{c}{x} + K_3^y) \tilde{\nu}_x^y \tilde{\nu}_+ + (\frac{c^y}{x} \frac{c^y}{+} + K_2^y) \tilde{\nu}_+ \tilde{\nu}_x + (\frac{c}{+} \frac{c}{x} + K_2) \tilde{\nu}_x^y \tilde{\nu}_+^y] \end{aligned} \quad (18)$$

From the classical solution, we guess $\eta^2 j_+^c j_+^2$ and $\eta^2 j_x^c j_+^2$ take values of $O(\eta^2)$, and thus a matrix elements given by a polynomial of them are not small enough to neglect from our Lagrangian. While, we hope $0 < \eta^2 < 1$ to be satisfied for convergence of a perturbative series/diagrams in terms of η^2 . This condition could conflict with the seesaw condition $g^2 m_D^2 \ll m_L^2$: A realization of the generalized seesaw situation is a non-trivial problem in our theory. In the next subsection, we will evaluate the one-loop effective potential of our theory. We hope the potential captures the essential feature of quantum dynamics of the system (1) even at the one-loop level. On the other hand, we simply drop $\frac{g}{2} (\tilde{\nu}_x^y + \tilde{\nu}_+^y + \tilde{\nu}_x^y) + h.c.$ which will give a coupling between the scalar and spinor sectors in L.

Now, we examine the variational condition, namely the vanishing condition of the terms linear in fluctuating scalars. From (8), we obtain the linear terms as follows:

$$\begin{aligned} & \tilde{\nu}_+^h [j_D^c j_+^{\sim y} + f^y g^c + g m_D^y \frac{c^y}{x} + 2m_L^y m_D^{\sim y}] \\ & + \eta^2 j_+^c j_+^2 + J_1 + 2j_x^c j_+^2 + 2K_1 \frac{c^y}{+} + \frac{J_2^y + 2K_3^y}{2} \frac{c}{+} + K_2^y \frac{c}{x} + K_3^y \frac{c^y}{x} \\ & \sim \tilde{\nu}_+^h [j_D^c j_+^2 + 4j_L^c j_+^2] + (g^y m_D^c + 2m_L^y m_D^{\sim y}) \tilde{\nu}_+^c \\ & \tilde{\nu}_x^h [g m_D^y \frac{c}{+} + \eta^2 J_1 \frac{c^y}{x} + K_2^y \frac{c}{+}] \\ & + h.c. \end{aligned} \quad (19)$$

They must vanish in our treatment of BEC. All terms given above have mass dimension $[Mass]^4$. We will employ a kind of Popov approximation to our HFB theory [38], i.e. $J_2 = K_2 = K_3 = 0$ (all of the anomalous self-energies will be dropped). At the classical solution (7), the vanishing condition of the coefficient function of $\tilde{\nu}_x$ gives

$$J_1 = J_1^y = \frac{2f}{g} \quad (20)$$

This expression would be modified under a renormalization of bare parameters as $J_1 = 2f^{(ren)} = g^{(ren)}$. We confirm that the coefficient of $\tilde{\nu}_+$ vanishes identically at the classical solution (7). From the vanishing condition of the coefficient of $\tilde{\nu}_+$ at (7), one finds

$$K_1 = K_1^y = g^2 \frac{m_D^4 + 2m_L m_D^3}{4m_L^2} \quad gf \quad (21)$$

This expression of K_1 takes a large value at the seesaw condition $m_D \ll m_L$.

Next, we will introduce several fields of the following definitions for the convenience of our discussion:

$$\begin{aligned} & \begin{pmatrix} \tilde{\nu}_x^c \\ \tilde{\nu}_x^y \end{pmatrix}^T; \quad \tilde{\nu}_x^c \quad \begin{pmatrix} \tilde{\nu}_x^c \\ \tilde{\nu}_x^y \end{pmatrix}^T; \quad \tilde{\nu}_M^c \quad \begin{pmatrix} \tilde{\nu}_M^c \\ \tilde{\nu}_M^y \end{pmatrix}^T; \quad \tilde{\nu}_M^c \quad \begin{pmatrix} \tilde{\nu}_+^c \\ \tilde{\nu}_+^y \end{pmatrix}^T; \quad \tilde{\nu}_M^c \quad \begin{pmatrix} \tilde{\nu}_+^c \\ \tilde{\nu}_+^y \end{pmatrix}^T; \\ & \begin{pmatrix} \tilde{\nu}_x^c \\ \tilde{\nu}_x^y \end{pmatrix}^T; \quad \tilde{\nu}_x^c \quad \begin{pmatrix} \tilde{\nu}_x^c \\ \tilde{\nu}_x^y \end{pmatrix}^T; \quad \tilde{\nu}_M^c \quad \begin{pmatrix} \tilde{\nu}_M^c \\ \tilde{\nu}_M^y \end{pmatrix}^T; \quad \tilde{\nu}_M^c \quad \begin{pmatrix} \tilde{\nu}_+^c \\ \tilde{\nu}_+^y \end{pmatrix}^T; \quad \tilde{\nu}_M^c \quad \begin{pmatrix} \tilde{\nu}_+^c \\ \tilde{\nu}_+^y \end{pmatrix}^T; \end{aligned} \quad (22)$$

Here, χ is a Majorana, ψ_R and ψ_L are right- and left-handed Majorana fields, respectively. T denotes transposition operation of a matrix. The Lagrangian density will be rewritten in the following form by these fields:

$$\mathcal{L} = V^{\text{tree}}[\psi; \chi] + \frac{1}{2} \psi^B + \frac{1}{2} \psi^F; \quad (23)$$

The matrices B and F are defined as follows:

$$\begin{aligned} B & \begin{pmatrix} B_{XX} & B_{XM} & B_{MM} \\ B_{MX} & B_{MX} & B_{MX} \end{pmatrix}; \quad \begin{pmatrix} B_{++} & B_{+} \\ B_{+} & B_{+} \end{pmatrix}; \quad \begin{pmatrix} B_{XM} & B_{X+} \\ B_{X+} & B_{X+} \end{pmatrix}; \quad \begin{pmatrix} B_{MX} & B_{+X} \\ B_{+X} & B_{+X} \end{pmatrix}; \\ F & \begin{pmatrix} F_{XX} & F_{XM} & F_{MM} \\ F_{MX} & F_{MX} & F_{MX} \end{pmatrix}; \quad \begin{pmatrix} F_{XX} & i\mathcal{I} \\ F_{XM} & g_{+P}^c + g_{+P}^{cy} \end{pmatrix}; \quad 0; \\ F & \begin{pmatrix} F_{MX} & g_{+P}^c + g_{+P}^{cy} \\ 0 & g_{+P}^{cy} \end{pmatrix}; \quad \begin{pmatrix} F_{MM} & i\mathcal{I} \\ F_{MX} & g_{+P}^c + g_{+P}^{cy} \end{pmatrix}; \quad \begin{pmatrix} m_D & \\ & 2m_L \end{pmatrix}; \end{aligned} \quad (24)$$

The definitions $\mathcal{I} = \frac{1}{2} \gamma^5 \gamma^0 \gamma^1 \gamma^2 \gamma^3$ and $P = \frac{1-i\gamma^5}{2}$ have been used. The Hermiticity $(\psi^B)^y = \psi^B$ is satisfied. Entries of B become such that,

$$\begin{aligned} B_{XX} & \begin{pmatrix} \mathcal{J}_1^2 (J_1 + j_+^c j_+^f) & 0 \\ 0 & \mathcal{J}_1^2 (J_1 + j_+^c j_+^f) \end{pmatrix}; \quad \begin{pmatrix} m_D^2 & 4m_L^2 \\ 0 & m_D^2 \end{pmatrix}; \quad \begin{pmatrix} 0 & 4m_L^2 \\ m_D^2 & 4m_L^2 \end{pmatrix}; \\ B_{++} & \begin{pmatrix} m_D^2 & \mathcal{J}_1^2 (J_1 + K_1 + j_+^c j_+^f + j_+^c j_+^f) \\ \frac{\mathcal{J}_1^2}{4} (J_2^y + (j_+^{cy})^2) & \frac{f g^y}{2} \end{pmatrix}; \quad \begin{pmatrix} m_D^2 & \frac{\mathcal{J}_1^2}{4} (J_2 + (j_+^c)^2) \\ m_D^2 & \mathcal{J}_1^2 (J_1 + K_1 + j_+^c j_+^f + j_+^c j_+^f) \end{pmatrix}; \\ B_{X+} & \begin{pmatrix} \mathcal{J}_1^2 (K_3 + \frac{y}{3} + \frac{cy}{+} \frac{c}{X}) & g_{m_D}^c \mathcal{J}_1^2 (K_2 + \frac{c}{+} \frac{c}{X}) \\ g_{m_D}^{cy} \mathcal{J}_1^2 (K_2 + \frac{y}{2} + \frac{cy}{+} \frac{cy}{X}) & \mathcal{J}_1^2 (K_3 + \frac{cy}{X} \frac{c}{+}) \end{pmatrix}; \\ B_X & \begin{pmatrix} g_{m_D}^{cy} & 0 \\ 0 & g_{m_D}^c \end{pmatrix}; \quad \begin{pmatrix} m_D (g_{+P}^{cy} + 2m_L) & 0 \\ 0 & m_D (g_{+P}^c + 2m_L) \end{pmatrix}; \end{aligned} \quad (25)$$

All of the off-diagonal elements of B are coming from particle-number-non-conserving interactions and/or mean fields of L under the HFB approximation. The diagonalizations of B and F will give "quasiparticle" excitation energy spectra of scalar and spinor fields in terms of the bare parameters/fields. Especially we have an interest on whether the spinor χ becomes massive or not under the one-loop quantum correction. If there is no massless fermion, then there is no Nambu-Goldstone (NG) fermion, and the Nambu-Goldstone theorem implies the absence of spontaneous SUSY breaking in our theory [9,12].

C. The One-loop Effective Potential

In this subsection, we will evaluate and examine the one-loop effective potential of our theory. We obtain the generating functional of our theory as follows:

$$\begin{aligned} Z & \int \mathcal{D} \psi \mathcal{D} \chi \exp \left\{ i \int d^4x \left[V^{\text{tree}}[\psi; \chi] + \frac{1}{2} \psi^B + \frac{1}{2} \psi^F \right] + (\text{source}) \right\}; \end{aligned} \quad (26)$$

The one-loop contribution to the effective potential is evaluated to be

$$\begin{aligned} V^{(1)}[\psi; \chi] & = V^{B(1)} + V^{F(1)}; \\ V^{B(1)} & = \frac{i}{2} \ln \det B = \frac{i}{2} \ln \det B_{MM} + \frac{i}{2} \ln \det \begin{pmatrix} B_{XX} & B_{XM} \\ B_{XM} & B_{MX} \end{pmatrix}; \\ V^{F(1)} & = \frac{i}{2} \ln \det F = \frac{i}{2} \ln \det F_{MM} + \frac{i}{2} \ln \det \begin{pmatrix} F_{XX} & F_{XM} \\ F_{XM} & F_{MX} \end{pmatrix}; \end{aligned} \quad (27)$$

The effective action is found to be

$$\Gamma(\text{com po}) = \int d^4x \left[V^{\text{tree}}[\psi; \chi] + V^{(1)}[\psi; \chi] \right]; \quad (28)$$

Note that M_X^F can become imaginary when inside the square root of the denominator of (36) takes a negative value. By putting the classical solution (7), one obtains

$$M_X^F = \frac{4f \mu \eta_L}{m_D^2 \sqrt{2(1 - \cos \chi)}}: \quad (37)$$

M_X^F becomes very small and will behave as a pseudo-NG fermion when $f \mu \eta_L \ll m_D$, and it vanishes at $m_L = 0$, while $(M_X^F)^2$ is always a positive quantity. It is an interesting fact that M_X^F will diverge under $\chi \rightarrow 0$, namely, not well-defined in the limit. (In the case of a non-SUSY dynamical model of the generalized seesaw mechanism, $\chi = 0$ is chosen as the vacuum state [36].) Therefore, a careful examination on χ is important (crucial) in our theory. As examined in the subsection A, χ of the classical solutions will take 0 or π . $M_X^F \rightarrow 0$ ($f \mu \eta_L = m_D^2$) if we choose $\chi = \pi$. We assume there is no spontaneous Lorentz symmetry breaking in the fermion sector. As a result, the spectrum of χ -particle must take the following Lorentz symmetric form:

$$E_X^F = \sqrt{k^2 + M_X^F{}^2}: \quad (38)$$

Therefore, we get the one-loop contribution of the fermion sector as

$$V^{F(1)} = \frac{i}{2} \text{Tr} \ln (k_0 - E_X^F)^2 (k_0 + E_X^F)^2 (k_0 - E_+^F)^2 (k_0 + E_+^F)^2 (k_0 - E_-^F)^2 (k_0 + E_-^F)^2: \quad (39)$$

We summarize the result of our analysis of the fermion sector: (i) Our model at $m_L = 0$ corresponds to the ordinary O'Raifeartaigh model, has R-symmetry and SUSY is broken at the ground state, and we have confirmed that an NG fermion appears [9,12]. (ii) Our model at $m_L \neq 0$, namely the modified O'Raifeartaigh model [11,12], will give the generalized seesaw mechanism at the point (7), while χ has a finite mass and SUSY seems not broken. We will proceed our examination to the boson sector of our theory.

In our treatment for the scalar sector, first we solve the secular equation $\det M_{MM}^B = 0$. Though the secular equation is quartic in the d'Alembertian, fortunately, we can diagonalize M_{MM}^B analytically because its secular equation will be factorized into a product of two quadratic equations of k . The results is

$$\det M_{MM}^B = (k_0 - E_{M1+}^B)(k_0 + E_{M1+}^B)(k_0 - E_{M1-}^B)(k_0 + E_{M1-}^B) \\ (k_0 - E_{M2+}^B)(k_0 + E_{M2+}^B)(k_0 - E_{M2-}^B)(k_0 + E_{M2-}^B): \quad (40)$$

There is no degeneracy in the spectra obtained from $\det M_{MM}^B = 0$. Here, the energy eigenvalues become

$$E_{M1-}^B(k) = \sqrt{k^2 + (M_{M1-}^B)^2}; \quad E_{M2-}^B(k) = \sqrt{k^2 + (M_{M2-}^B)^2}; \\ M_{M1-}^B = \frac{c_2 + c_3 - \mu \eta}{2} \frac{1}{2} \sqrt{c_2^2 - g + \mu \eta^2 + 4j_1^2 f} \\ M_{M2-}^B = \frac{c_2 + c_3 + \mu \eta}{2} \frac{1}{2} \sqrt{c_2^2 - g - \mu \eta^2 + 4j_1^2 f} \\ c_1 = m_D (g_X^{cy} + 2m_L); \quad c_2 = m_D^2 + 4m_L^2; \\ c_3 = m_D^2 + \mu \eta^2 (J_1 + K_1 + j_+^c f + j_X^c f); \quad c_4 = \frac{\mu \eta^2}{4} (J_2 + (\frac{c}{+})^2) - \frac{fg^y}{2}: \quad (41)$$

$j_1^2 f$ includes the phase χ . If we put the expressions at the classical solution for μ, J_1 and K_1 to c_3 and c_4 with employing the Popov approximation $J_2 = 0$, we get

$$M_{M1-}^B = M_{M2-}^B \\ = m_D^2 + \frac{\mu \eta^2 j_X^c f}{2} + 2m_L^2 + \frac{m_D^4}{8m_L^2} + \frac{m_D^3}{4m_L} \frac{5}{2} \mu \eta f \\ \frac{1}{2} \sqrt{\frac{\mu \eta^2 j_X^c f}{4} + m_L^2 + \frac{m_D^4}{16m_L^2} + \frac{m_D^3}{8m_L} \frac{5}{4} \mu \eta f^2 + m_D^2 \frac{\mu \eta^2 j_X^c f}{4} + m_L^2 + \mu \eta j_X^c \eta_L \cos \chi} \quad \#_{1=2} \quad (42)$$

Therefore, we conclude

$$M^F < M_{M1-}^B; M_{M2-}^B; \quad (43)$$

at $f < 0, g > 0$. The mass eigenvalues of $M_{M_{1+}}^B$ and $M_{M_{2+}}^B$ become tachyonic at

$$M_{M_{1+}}^B; c_2(c_3 - j_4) < j_1 j^2; \quad M_{M_{2+}}^B; c_2(c_3 + j_4) < j_1 j^2; \quad (44)$$

and an appearance of tachyon indicates the instability of vacuum state [12,45]. In our condition (44) of tachyonic masses, c_3 and c_4 include the HFB self-energies. At the classical solution (7) with $J_2 = 0$, the tachyon condition (44) becomes such that

$$2f - g J_1 + K_1 = 5f \frac{1}{g} \frac{m_D^4}{4m_L^2} + \frac{m_D^3}{2m_L} > 0: \quad (45)$$

Here we have assumed f/g as real. (Again, we wish to rewrite that f, J_1, K_1 , and J_2 have mass dimension $[\text{Mass}]^2$, while g is dimensionless.) Hence, the vicinity of the classical solution is stable if $f < 0$ and $g > 0$. Due to the HFB self-energies and c_+ , there are several differences between M_{MM}^B and the mass eigenvalues obtained from M_{MM}^B : The mass spectra of bosons and fermions are not symmetric (namely, not supersymmetric) in our theory. We regard the vacuum energy as the order parameter of SUSY-breaking, we must examine the local/global structure of the one-loop effective potential to clarify whether the vacuum energy vanishes or not before concluding a breakdown of SUSY.

In the determinant $\det(M_{XX}^B - M_{XM}^B (M_{MM}^B)^{-1} M_{MX}^B g)$, we wish to concentrate upon the vicinity of the classical solution (7). At the point (7) with the neglect of J_2 , M_{++}^B in the expression of (25) becomes diagonal. This helps us to evaluate the mass eigenvalue of M_X^B in an analytic manner. Then we get $(M_X^B)^2$ in terms of bare parameters as follows:

$$\begin{aligned} (M_X^B)^2 &= \lim_{p^2 \rightarrow 0} (\mathcal{A} j - \mathcal{B} j); \\ \mathcal{A} &= a + \frac{j(j^2 + c^2)b + c j j^2 (d e^y + y d^y e)j}{bc j^3}; \quad \mathcal{B} = \frac{j(j + y)(y b - d e^y)j}{bc j^3}; \\ a &= j^2(J_1 + j_+^c j^2); \quad b = m_b^2 - 4m_L^2; \\ c &= m_b^2 - j^2(J_1 + K_1 + j_+^c j^2 + j_X^c j^2); \quad d = y m_D - c y; \quad e = m_b (g^y - c y + 2m_L); \\ &= j^2(K_3^y + \frac{c y}{+ X}); \quad y m_D - c = j^2(K_2 + \frac{c}{+ X}); \end{aligned} \quad (46)$$

These $a; b; c; d; e$; are matrix elements of M^B (see (25)). We also set four-momentum as $p = 0$ in the bosonic matrices. We examine the stability conditions of M_X^B . Especially, we have interest on its behavior in the vicinity of the classical solution with the seesaw condition $j j^2 j_X^c j^2 = m_D^2 - m_L^2$. A direct evaluation from (46) with (7), (20), (21), and by employing a Popov approximation $J_2 = K_2 = K_3 = 0$ gives

$$\begin{aligned} (M_{X+}^B)^2 &= (M_X^B)^2 \\ &= \frac{1}{(m_D^2 + 4m_L^2)(3gf + \frac{m_D^4}{4m_L^2} + \frac{m_D^3}{2m_L})} \\ &\quad \frac{h}{48g^2 f^2 m_L^2 + 2g^2 f^2 m_D^2 + 8g f m_D^3 m_L + 6g f m_D^4 + 3g f \frac{m_D^5}{m_L} + \frac{g f m_D^6}{2 m_L^2}}; \end{aligned} \quad (47)$$

Hence, if we take into account the seesaw condition $m_D = m_L$, then we obtain $g > 0$ and $f < 0$ is the stability condition of $(M_X^B)^2$. A rough estimation gives

$$(M_X^B)^2 \approx 2gf - O(gf); \quad (48)$$

We conclude that, with taking into account (45) and (48), the generalized seesaw mechanism can take place under $f < 0$ and $g > 0$ (we have assumed that $j f j (m_L = m_D) = 1$). It is worth noticing that $(M_X^B)^2 = m^4 = 8m_L^2$ (negative) if we set $K_1 = 0$, and thus the HFB self-energy K_1 is important for stability of the potential at the classical solution. Finally one finds

$$E_X^B = \frac{q}{k^2 + (M_X^B)^2}; \quad (49)$$

and we obtain the one-loop contribution of the scalar sector as follows:

$$\begin{aligned} V^{B(1)} &= \frac{i}{2} \text{Tr} (k_0 - E_{X+}^B) (k_0 + E_{X+}^B) (k_0 - E_X^B) (k_0 + E_X^B) \\ &\quad (k_0 - E_{M_{1+}}^B) (k_0 + E_{M_{1+}}^B) (k_0 - E_{M_1}^B) (k_0 + E_{M_1}^B) \\ &\quad (k_0 - E_{M_{2+}}^B) (k_0 + E_{M_{2+}}^B) (k_0 - E_{M_2}^B) (k_0 + E_{M_2}^B); \end{aligned} \quad (50)$$

It is a well-known fact that the naive dimensional regularization, suitable to keep a gauge invariance in a non-SUSY gauge theory, will break SUSY through the regularization. To circumvent of this problem is relatively easier in non-gauge models, while the problem is severe in SUSY gauge theories, and the method of "dimensional reduction" regularization seems more suitable [18]. Since the Lagrangian we consider here is not a gauge model, here we employ a simple cutoff scheme for regularizations of integrals. After performing the four-dimensional momentum integration, the one-loop contribution to the effective potential will be obtained as follows:

$$\begin{aligned}
V^{(1)} = & \frac{1}{16^2} \left(M_{X^+}^B + M_X^B + M_{M_{1+}}^B + M_{M_1}^B + M_{M_{2+}}^B + M_{M_2}^B \right)^2 (M_X^F + M_+^F + M^F)^2 \\
& + 4 \ln \frac{(1 + M_{X^+}^B)^2 (1 + M_X^B)^2 (1 + M_{M_{1+}}^B)^2 (1 + M_{M_1}^B)^2 (1 + M_{M_{2+}}^B)^2 (1 + M_{M_2}^B)^2}{(1 + M_X^F)^2 (1 + M_+^F)^2 (1 + M^F)^2} \\
& M_{X^+}^B \ln \left(1 + \frac{2}{M_{X^+}^B} \right) + M_X^B \ln \left(1 + \frac{2}{M_X^B} \right) + 2M_X^F \ln \left(1 + \frac{2}{M_X^F} \right) + 2M_+^F \ln \left(1 + \frac{2}{M_+^F} \right) + 2M^F \ln \left(1 + \frac{2}{M^F} \right) \\
& M_{M_{1+}}^B \ln \left(1 + \frac{2}{M_{M_{1+}}^B} \right) + M_{M_1}^B \ln \left(1 + \frac{2}{M_{M_1}^B} \right) + M_{M_{2+}}^B \ln \left(1 + \frac{2}{M_{M_{2+}}^B} \right) + M_{M_2}^B \ln \left(1 + \frac{2}{M_{M_2}^B} \right); \quad (51)
\end{aligned}$$

where, Λ denotes the four-momentum cutoff. By the standard method for handling the effective potential, namely, remove contributions they will vanish at $\Lambda \rightarrow \infty$ (there is no divergent constant due to $N = 1$ SUSY), one obtains

$$\begin{aligned}
V^{(1)} = & \frac{1}{16^2} \left(M_{X^+}^B \ln \frac{M_{X^+}^B}{\Lambda^2} + M_X^B \ln \frac{M_X^B}{\Lambda^2} + 2M_X^F \ln \frac{M_X^F}{\Lambda^2} + 2M_+^F \ln \frac{M_+^F}{\Lambda^2} + 2M^F \ln \frac{M^F}{\Lambda^2} \right. \\
& \left. + M_{M_{1+}}^B \ln \frac{M_{M_{1+}}^B}{\Lambda^2} + M_{M_1}^B \ln \frac{M_{M_1}^B}{\Lambda^2} + M_{M_{2+}}^B \ln \frac{M_{M_{2+}}^B}{\Lambda^2} + M_{M_2}^B \ln \frac{M_{M_2}^B}{\Lambda^2} \right); \\
& (M_X^B; M_X^F; M_{M_1}^B; M_{M_2}^B; M^F); \quad (52)
\end{aligned}$$

We have arrived at a generalization of the so-called SUSY Coleman-Weinberg potential discussed in Refs. [11,12] (see also, Ref. [46]). In our $V^{(1)}$, the one-loop contribution of X -field is also included. For obtaining a one-loop potential which will not diverge into the negative-energy direction at the limit $j_X^c \rightarrow \infty$ [47], we should impose both

$$(M^F) < (M_{M_1}^B)^2; (M_{M_2}^B)^2 \quad (53)$$

and

$$(M_X^F)^2 < (M_X^B)^2; \quad (54)$$

We have known from (43) that (53) is satisfied, while if

$$x > 1; \quad 1 > g > 0; \quad 0 > f(m_L = m_D^2) > -1; \quad (55)$$

then (54) is satisfied. We should set model parameters with respect to these relations. The mass eigenvalues will degenerate under using (7) for μ with the Popov approximation. Therefore, we will denote them as follows:

$$M_X^B = M_{X^+}^B = M_X^B; \quad M_+^B = M_{M_{1+}}^B = M_{M_{2+}}^B; \quad M^B = M_{M_1}^B = M_{M_2}^B; \quad (56)$$

Since $m_D, f < m_L$, especially we have interest on a situation

$$M_X^F = M_X^B < M_+^F < M_+^B = M^F < M^B; \quad (57)$$

We have arrived at the order of mass eigenvalues and it is the crucial result for our discussion hereafter.

Since our effective potential and mass eigenvalues have similarities with those of the Minimal Supersymmetric Standard Model (MSSM) [19,20], let us utilize some methods/results from it. In theory of MSSM, SUSY is explicitly broken by a vacuum energy and several soft mass parameters, while it is not broken in the Lagrangian level as the starting point of our model. Thus, discussion on renormalization would become simpler than that of MSSM. A renormalization-group invariant calculation for renormalization of our $V^{\text{tree}} + V^{(1)}$ is subtle, because it includes many different mass parameters/scales [19,20,48-53]. Since our interest is to examine a possibility of a realization of

the generalized seesaw mechanism in the vicinity of the classical solution (7) with taking into account the one-loop contribution (52), we concentrate on a VEV of χ under the situation (55). Unfortunately, it is difficult to find a global minimum of the potential $V^{\text{tree}} + V^{(1)}$ because it has many parameters, μ^2 , μ_X^2 , J_1 and K_1 which should be determined variationally. For example, if we put the expressions of classical solution (7) for condensate $\langle \chi \rangle$ to reduce variational parameters and try to find a minimum with respect to variation of μ_X^2 , the potential might give a non-vanishing vacuum energy (hence SUSY is broken) because this procedure corresponds to a restriction of trial functions in the variation: To achieve a true vacuum might be difficult. In the usual prescription of renormalization, a running coupling will be used to remove a renormalization point from a theory to get a physical (renormalization-group invariant) potential, though this procedure is difficult in our case. To make our problem tractable for our purpose, we will use the following definition of $V^{(1)}$ [50,53] by taking into account the Appelquist-Carazzone decoupling theorem [54]:

$$V^{(1)} = \frac{1}{8} \sum_l \chi^h \left(\mu^2 (M_1^B)^2 (M_1^B)^4 \ln \frac{(M_1^B)^2}{2} + \mu_X^2 (M_1^F)^2 (M_1^F)^4 \ln \frac{(M_1^F)^2}{2} \right); \quad (l = X; +; -); \quad (58)$$

(A quite clear example of the decoupling theorem can be found in Ref. [20].) Here, $\theta(x)$ is the Heaviside step function, it has been introduced to define mass thresholds inside the potential. We have changed the regularization method to \overline{MS} (modified minimal subtraction scheme). $\mu^2 = \mu^2(\mu^2)$, where μ^2 denotes the \overline{MS} renormalization scale. Of course, M_1^B and M_1^F are functions of μ_X^2 . The logarithmic functions appeared in the above equation must satisfy $j \ln(M^2 = \mu^2) < 1$ (μ^2 : a renormalization point) for the justification for our loop expansion. We should find the situation where $V^{(1)}(\mu^2) = 0$ and $\frac{d}{d\mu^2}(V^{\text{tree}} + V^{(1)}) = 0$ are simultaneously satisfied [50]. It is a hard task to arrive from the complete theory to an effective theory of lowest region in (57) with running parameters with μ^2 . In such a top-down approach, as we know from (57), we have totally six decoupling scales until we arrive at the region $\mu^2 < (M_X^F)^2$ where all particles are decoupled, and then V^{tree} alone gives the renormalization-group invariant potential [50,53]. Hence, first we wish to consider the case $(M_X^F)^2 < \mu^2 < \text{others}$. In this case, the potential of the one-loop contribution can be written down as follows:

$$V^{(1)} = \frac{1}{8} (M_X^F)^4 \ln \frac{(M_X^F)^2}{2}; \quad (59)$$

Here, we simply have assumed that the effect of decoupled particles is already included by a renormalization of parameter. This $V^{(1)}$ gives a positive contribution to our one-loop potential. After put the classical solution (7) to χ of this $V^{(1)}$, choose $\mu_X^2 = \mu_X^2$, and take the derivative of $V^{\text{tree}} + V^{(1)}$ with respect to μ_X^2 we get $h_{\chi^i} = m_D^2 = (2g m_L)$ from the stationary condition: We find that the effective field theory of this renormalization point/scale gives the same expression for VEV of χ with its classical solution, shows the generalized seesaw mechanism. Needless to say, we will also obtain $h_{\chi^i} = m_D^2 = (2g m_L)$ at the complete decoupled region $\mu^2 < (M_X^F)^2$ because $V^{(1)} = 0$. Since the improved V^{tree} is the "exact" potential (with satisfying the matching condition) [50,53], an observation will find that the vacuum is supersymmetric in the energy scale $\mu^2 < (M_X^F)^2$. We conclude that, under a reasonable choice of model parameters with respect to the conditions for stability of the vicinity of the classical solution (7) and a justification on convergence of the loop expansion, certainly (7) is robust against a quantum correction, will not obtain a radical modification, and thus the generalized seesaw mechanism takes place.

III. SUPERSPACE FORMALISM

In this section we will calculate the one-loop effective potential in superfield formalism [2,17,55], though we have to use components of superfields at several points of our discussions, especially for our consideration on BEC. For example, it seems difficult to consider the HFB approximation of quantum fluctuations of scalars in the superfield formalism, and thus the selfenergies as $J_1; K_1$ do not appear in our superspace formalism. It is a problem inherent in the superspace formalism, and as a result, the one-loop contribution of the superspace formalism is different from that of the component field formalism. Moreover, it seems difficult to examine the generalized seesaw mechanism by our superfield formalism of one-loop potential because mass eigenvalues of fermion and boson sectors will not be derived under a direct manner. Therefore the component field formalism is better to describe the dynamics and physical property of the scalar sector with having the BEC. The purpose of this section is to make a comparison between the two formalisms. The examination on the generalized seesaw mechanism is beyond scope of this section. We employ the background field method, the standard method of superspace formalism [17], to take into account the

BEC in our model:

$$\begin{aligned}
 X &= X^c + X^{\sim}; \quad X^c = \frac{c}{X} + F_X^c; \quad X^{\sim} = \tilde{X} + X + F_X^{\sim}; \\
 &= c + \tilde{X}; \quad c = c + F^c; \quad \tilde{X} = \tilde{X} + X + F_X^{\sim};
 \end{aligned}
 \tag{60}$$

Similar to the case of component field formalism given in the previous section, again we assume the condensates $\frac{c}{X}$ and c are independent on spacetime coordinates. The Lagrangian will be converted into the following form :

$$\begin{aligned}
 L &= L^c + \mathbb{F}; \\
 L^c &= X^{cy}X^c + \frac{cy}{4} \frac{c}{X} + \frac{cy}{4} c + fX^c + \frac{g}{2} X^c \frac{c}{X} + m_D \frac{c}{X} + m_L c^c + (\text{h.c.}); \\
 \mathbb{F} &= X^y X^{\sim} + \frac{1}{2} y M \frac{D^2}{4} g \frac{c}{X} X^{\sim} + \frac{D^{-2}}{4} g^y \frac{cy}{X} X^y \tilde{y} :
 \end{aligned}
 \tag{61}$$

Here, we have used the equivalent relations $\int d^4x = D^2=4$ and $\int d^4x = \frac{D^{-2}}{4}$ under the integration of d^4x inside the action functional of the theory, and have dropped terms linear in the fluctuating super fields X^{\sim}, \tilde{X} . We have introduced several matrix notations defined as follows:

$$M = \begin{pmatrix} \tilde{X}^+; \tilde{X}^-; \tilde{X}^y; \tilde{X}^{\sim} \end{pmatrix}^T; \quad \begin{pmatrix} \frac{D^2 D^{-2}}{16} & 0 \\ \frac{D^{-2} D^2}{4} C & 0 \end{pmatrix} \begin{pmatrix} \frac{D^2}{4} C^y \\ 0 \end{pmatrix} \delta^8(z-z^0); \quad C = m_D \frac{1}{2} + m_L \frac{1}{2} + \frac{g}{2} X^c \frac{1}{2}; \tag{62}$$

The sigma matrices (the definition: $\sigma^0 = \frac{1}{2} \sigma_2$, while $\sigma^1; \sigma^2; \sigma^3$ are the ordinary Pauli matrices) act on the two-dimensional chirality space $(+; -)$. The chiral and antichiral delta functions are defined as

$$\frac{\delta^8(z^0)}{\delta^8(z)} = \frac{D^{-2}}{4} \delta^8(z-z^0); \quad \frac{\delta^y(z^0)}{\delta^y(z)} = \frac{D^2}{4} \delta^8(z-z^0); \quad z = (x; i); \tag{63}$$

The generating functional will be written down in the following form :

$$\begin{aligned}
 Z &= \int D X^c D X^y D \tilde{X}^+ D \tilde{X}^{\sim} D \tilde{X}^y \exp i \int d^4x L + (\text{sources}) \\
 &= \int D X^c D X^y \exp i \int d^4x L^c + X^y X^{\sim} \frac{1}{2} + \frac{i}{2} \text{Tr} \ln M + G; \\
 G &= \frac{i}{2} \int d^8z \int d^8z^0 \frac{1}{2} J^y(z) M^{-1} \delta^8(z-z^0) J^y(z^0);
 \end{aligned}
 \tag{64}$$

Here, $d^8z = d^4x d^2 \theta^2$. To obtain the final expression of Z in (64), we have neglected contributions of (anti)chiral sources. The definition of J is

$$J = \begin{pmatrix} g \frac{c}{X} \\ 0 \end{pmatrix} \frac{1+\sigma_3}{2} \begin{pmatrix} 0 \\ g^y \frac{cy}{X} X^y \frac{1+\sigma_3}{2} \end{pmatrix}; \tag{65}$$

By putting the components of X^c and c , we can confirm the fact that L^c becomes

$$L^c = V^{\text{tree}} [c; \frac{c}{X}]; \tag{66}$$

Hence the tree level potential is the same in both of the formalisms. Next, we divide M as follows:

$$\begin{aligned}
 M &= M_0 \quad M^0; \quad M_0 = \begin{pmatrix} \frac{D^2 D^{-2}}{16} & 0 \\ 0 & \frac{D^{-2} D^2}{16} \end{pmatrix} \begin{pmatrix} 0 \\ 0 \end{pmatrix}; \\
 M_0^{-1} &= \frac{1}{2} \begin{pmatrix} \frac{D^2 D^{-2}}{16} & 0 \\ 0 & \frac{D^{-2} D^2}{16} \end{pmatrix} \begin{pmatrix} 0 \\ 0 \end{pmatrix}; \quad M^0 = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + \frac{D^2}{4} C^y \begin{pmatrix} 0 \\ 0 \end{pmatrix};
 \end{aligned}
 \tag{67}$$

The one-loop effective action $\frac{i}{2}\text{Tr}\ln M$ is evaluated to be

$$\begin{aligned} \stackrel{(1)}{\text{(super)}} \frac{i}{2}\text{Tr}\ln M &= \frac{i}{2}\ln \text{Det} M_0 + \frac{i}{2}\text{Tr}\ln(1 - M_0^{-1}M^0) = \frac{i}{2}\text{Tr}\ln(1 - M^0 M_0^{-1}) \\ &= \lim_{z^0 \rightarrow z^0} \frac{i}{2}\text{tr} \int d^8z \ln(1 - \frac{1}{16}C^Y C^X \frac{D^2 \bar{D}^2}{16} \delta(z-z^0)) \end{aligned} \quad (68)$$

We have dropped $\frac{i}{2}\ln \text{Det} M_0^{-1}$ because it does not contribute to $\stackrel{(1)}{\text{(super)}}$. The relations $M_0^{-1}M^0 = 1 - M^0 M_0^{-1}$ and the commutator $[M_0^{-1}, M^0] = 0$ have been used. From the following identity in superspace,

$$\frac{D^2 \bar{D}^2}{16} (\delta^0)^2 (\delta^0)^2 = \delta^0; \quad (69)$$

the effective potential is found to be

$$\begin{aligned} V_{\text{(super)}}^{(1)} &= \frac{\stackrel{(1)}{\text{(super)}}}{\int d^4x} = \frac{i}{2}\text{tr} \int d^2\theta \int d^2\bar{\theta} \frac{d^4p}{(2\pi)^4} \frac{1}{p^2} \ln p^2 + C^Y C^X \\ &= \frac{1}{2}\text{tr} \int d^2\theta \ln(1 + \frac{C^Y C^X}{2}) + C^Y C^X \ln(1 + \frac{C^2}{C^Y C^X}) \\ &= \frac{\int j^c j^c}{4} \int F_X^c j^c \ln(1 + \frac{C^2}{M_C j^c}) + \frac{\int j^c j^c}{4} \int F_X^c j^c \int X^c j^c \frac{C^2}{(C^2 + M_C j^c) M_C j^c} \\ &\quad \int F_X^c j^c \frac{\int j^c j^c}{4} \ln \frac{C^2}{M_C j^c}; \quad (C^2 \neq 1); \\ M_C j^c &= m_D^2 + m_L^2 + \frac{\int j^c j^c}{4} \int X^c j^c; \end{aligned} \quad (70)$$

Of course, the mass dimension of $V_{\text{(super)}}^{(1)}$ is $[\text{mass}]^4$. Both V^{tree} and $V_{\text{(super)}}^{(1)}$ vanish simultaneously at the classical vacuum (7) and SUSY is not broken. To make our calculation on G in (64) tractable, we approximate M^{-1} by replacing $C \rightarrow m_D$. Then we get

$$\begin{aligned} G &= \int d^8z \int j^c j^c \int X^c \frac{1}{m_D^2} X^Y + X^Y \frac{1}{m_D^2} X^c \\ &\quad + \int d^6z (g^c)^2 (\frac{C}{+})^2 X^c \frac{m_D}{m_D^2} X^c + \int d^6z (g^Y)^2 (\frac{C^Y}{+})^2 X^Y \frac{m_D}{m_D^2} X^Y; \end{aligned} \quad (71)$$

Obviously, G includes a Kahler potential and (anti)chiral superpotentials of the fluctuating X -eld. From a consideration by the Wick theorem, one finds $(\frac{C}{+})^2$ in the Kahler potential corresponds to the propagator $\langle \tilde{\psi}_+^Y \tilde{\psi}_+^Y \rangle$, while $m_D = (\frac{C}{+})^2$ in the chiral and antichiral superpotential parts of G came from $\langle \tilde{\psi}_+^Y \tilde{\psi}_+^Y \rangle$ and $\langle \tilde{\psi}_+^Y \tilde{\psi}_+^Y \rangle$, respectively. (An examination of mass dimensions of these propagators is also helpful.) Because $(\frac{C}{+})^2 X^c$ or $(1 = (\frac{C}{+})^2) X^c$ are chiral super elds, $\frac{D^2}{4}$ can be inserted between them inside the integration $\int d^8z$. Therefore we get

$$\begin{aligned} \int d^8z X^Y X^c \quad iG &= \frac{1}{2} \int d^8z (X^Y; X^c) M_X \quad X^Y; \\ M_X &= \begin{pmatrix} 1 & \frac{g^2}{2} j^c j^c \frac{1}{m_D^2} \\ \frac{g^2}{2} (\frac{C}{+})^2 (\frac{D}{4}) \frac{m_D}{m_D^2} & 1 \end{pmatrix} \frac{1}{\frac{g^2}{2} j^c j^c \frac{1}{m_D^2}} A; \end{aligned} \quad (72)$$

Integration of $D X^Y D X^c$ will give $\text{Det}^{-1} M_X$, and this determinant gives a polynomial of F_+^c and F_+^{cY} . Because $F_+^c = F_+^{cY} = 0$ at (7), the one-loop contribution of $\text{Det}^{-1} M_X$ is also vanish and we conclude that SUSY is not broken at the classical solution (7).

IV . CONCLUSION

In summary, we have examined the mass spectra of scalars and spinors of the modified O'Raifeartaigh model by our evaluation of the one-loop effective potential in the component field formalism, especially in the vicinity of the classical solution (7) of the model, from the context of the generalized seesaw mechanism. The BEC in the scalar sector has been considered, while the spinor sector has a mathematical similarity with relativistic theory of superconductivity [56,57]. Therefore, some parts of our formulation has some similarities with that of theory of supersymmetric (color-)superconductivity [58,59], though the intrinsic dynamics of them are quite different. We have emphasized that it becomes possible for us to examine the mass spectra for the generalized seesaw mechanism of neutrino by introducing the left-handed Majorana mass term (namely, a modification [11,12]) to the ordinary O'Raifeartaigh model. Our calculation at the one-loop level of the effective potential of the component field formalism indicates that SUSY is not broken in the theory due to the absence of an NG fermion, and have concluded that SUSY is not broken at the classical vacuum of the one-loop potential of the super field formalism.

In this paper, we have discussed several VEVs of scalars. It is interesting for us to consider some possible relations between inflation of cosmology, scalar fields of (1), a (generalized) seesaw mechanism of neutrino, and a spontaneous SUSY breaking. The scalar field χ seems to have a special role in a determination of local/global minima of an effective potential of (1), while its VEV determines a right-handed Majorana mass parameter in our theory. Thus it is interesting for us to investigate a possible scenario of a relation between χ and an inflation for our further investigation. It might be possible to investigate the lightest fermion χ -field could become a candidate of dark matter. In the strong CP problem of QCD, a theta angle gives us an important issue on axion. An investigation of a relation between χ and QCD theta angle (and also the Peccei-Quinn mechanism [60]) is far beyond scope of this paper, though it is also an interesting problem.

Present address: Department of Physics, University of Texas at Austin, tadafum@physics.utexas.edu

- ¹ S. Weinberg, A Model of Leptons, Phys. Rev. Lett. 19, 1264 (1967).
- ² J. Wess and J. Bagger, Supersymmetry and Supergravity, (Princeton University Press, Princeton, USA, 1991).
- ³ E. Witten, Dynamical Breaking of Supersymmetry, Nucl. Phys. B 185, 513 (1981).
- ⁴ E. Witten, Constraints of Supersymmetry Breaking, Nucl. Phys. B 202, 253 (1982).
- ⁵ I. A. Heck, M. Dine and N. Seiberg, Dynamical Supersymmetry Breaking in Four-Dimensions and its Phenomenological Implications, Nucl. Phys. B 256, 557 (1985).
- ⁶ A. E. Nelson and N. Seiberg, R-Symmetry Breaking versus Supersymmetry Breaking, Nucl. Phys. B 416, 46 (1994).
- ⁷ J. Bagger, R. Poppitz and L. Randall, The R-axion from Dynamical Supersymmetry Breaking, Nucl. Phys. B 426, 3 (1994).
- ⁸ G. F. Giudice and R. Rattazzi, Theories with Gauge-Mediated Supersymmetry Breaking, Phys. Rep. 322, 419 (1999).
- ⁹ Y. Shadmi and Y. Shim'an, Dynamical Supersymmetry Breaking, Rev. Mod. Phys. 72 25 (2000).
- ¹⁰ K. Intriligator, N. Seiberg and D. Shih, Dynamical SUSY Breaking in Meta-Stable Vacua, J. High Energy Phys. 04 021 (2006).
- ¹¹ K. Intriligator, N. Seiberg and D. Shih, Supersymmetry Breaking, R-Symmetry Breaking and Metastable Vacua, J. High Energy Phys. 07 017 (2007).
- ¹² K. Intriligator and N. Seiberg, Lectures on Supersymmetry Breaking, Class. Quantum Grav. 24, S741 (2007).
- ¹³ S. Ray, Some Properties of Meta-Stable Supersymmetry-Breaking Vacua in Wess-Zumino Models, Phys. Lett. B 642, 137 (2006).
- ¹⁴ L. O'Raifeartaigh, Spontaneous Symmetry Breaking for Chiral Scalar Superfields, Nucl. Phys. B 96, 331 (1975).
- ¹⁵ M. H. Uq, On Spontaneous Breakdown of Fermion Number Conservation and Supersymmetry, Phys. Rev. D 14, 3548 (1976).
- ¹⁶ M. H. Uq, Evaluation of Effective Potential in Superspace, Phys. Rev. D 16, 1733 (1977).
- ¹⁷ A. Yu. Petrov, Quantum Superfield Supersymmetry, arXiv:hep-th/0106095.
- ¹⁸ I. Jack and D. R. T. Jones, Regularisation of Supersymmetric Theories, in 'Perspectives on Supersymmetry', ed. G. Kane (World Scientific, Singapore, 1998), [hep-ph/9707278].
- ¹⁹ M. S. Carena, J. R. Espinosa, M. Quiros and C. E. M. Wagner, Analytic Expressions for Radiatively Corrected Higgs Masses and Couplings in the MSSM, Phys. Lett. B 355, 209 (1995).
- ²⁰ M. S. Carena, M. Quiros, C. E. M. Wagner, Effective Potential Methods and the Higgs Mass Spectrum in the MSSM, Nucl. Phys. B 461, 407 (1996).
- ²¹ M. Dine, N. Seiberg and S. Thomas, Higgs Physics as a Window beyond the MSSM, Phys. Rev. D 76, 095004 (2007).
- ²² E. Dudas, S. Lavignac and J. P. A. M. Entier, A Light Neutralino in Hybrid Models of Supersymmetry Breaking, arXiv:0808.0562.
- ²³ S. N. Bose, Plancks Gesetz und Lichtquantenhypothese, Z. Phys. 26, 178 (1924).
- ²⁴ A. Einstein, Quantentheorie des einatomigen idealen Gases, Sitzungsberichte der Preussischen Akademie der Wissenschaften, Phys. Math. Kl. L 261, 3 (1924).
- ²⁵ N. N. Bogoliubov, J. Phys. (Moscow) 11, 23 (1947).

- ²⁶ J. Bardeen, L. N. Cooper and J. R. Schrieffer, Theory of Superconductivity, *Phys. Rev.* 108, 1175 (1957).
- ²⁷ Y. Nambu and G. Jona-Lasinio, Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. I, *Phys. Rev.* 122, 345 (1961).
- ²⁸ Y. Nambu and G. Jona-Lasinio, Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. II, *Phys. Rev.* 124, 246 (1961).
- ²⁹ K. Hagiwara, Theory of Dynamical Symmetry Breaking, *Prog. Theor. Phys. Suppl.* 104, 1 (1991).
- ³⁰ M. Gell-Mann, P. Ramond and R. Slansky, Complex Spinors and Unified Theories, in *Supergravity* (P. Nieuwenhuizen and D. Z. Freedman, eds.), (North-Holland, Amsterdam (1975)), p.315.
- ³¹ T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe* (O. Sawada and A. Sugamoto, eds.) (KEK, Tsukuba, Japan (1979)), p.95.
- ³² R. N. Mohapatra and G. Senjanovic, Neutrino Mass and Spontaneous Parity Violation, *Phys. Rev. Lett.* 44, 912 (1980).
- ³³ C. T. Hill, M. A. Luty and E. A. Paschos, Electroweak Symmetry Breaking by Fourth-Generation Condensates and the Neutrino Spectrum, *Phys. Rev. D* 43, 3011 (1991).
- ³⁴ R. S. Chivukula, B. A. Dobrescu, H. Georgi and C. T. Hill, Top Quark Seesaw Theory of Electroweak Symmetry Breaking, *Phys. Rev. D* 59, 075003 (1999).
- ³⁵ T. Ohsaku, Dynamical Dirac Mass Generation in the Supersymmetric Nambu–Jona-Lasinio Model with the Seesaw Mechanism of Neutrinos, arXiv: 0801.1256, submitted for publication.
- ³⁶ T. Ohsaku, Collective Excitations in the Nambu–Jona-Lasinio Model with Left-Right-Asymmetric Majorana Mass Terms, arXiv: 0802.1286, submitted for publication.
- ³⁷ G. Parisi, *Statistical Field Theory*, (Addison-Wesley, Reading, MA, USA, 1988).
- ³⁸ A review on studies of non-relativistic interacting boson gas: J. O. Andersen, Theory of the Weakly Interacting Bose Gas, *Rev. Mod. Phys.* 76, 599 (2004).
- ³⁹ A. A. Abrikosov, L. P. Gor'kov and I. E. Dzyaloshinskii, *Methods of Quantum Field Theory in Statistical Physics* (Dover, New York, 1963).
- ⁴⁰ A. L. Fetter and D. J. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, New York, 1971).
- ⁴¹ J. W. Negele and H. O. Rland, *Quantum Many-Particle Systems* (Addison-Wesley, New York, 1987).
- ⁴² J. I. Kapusta, *Finite-Temperature Field Theory* (Cambridge University Press, Cambridge, 1989).
- ⁴³ L. P. Gor'kov, *Sov. Phys. JETP* 7, 505 (1958).
- ⁴⁴ Y. Nambu, Quasi-particles and Gauge Invariance in the Theory of Superconductivity, *Phys. Rev.* 117, 648 (1960).
- ⁴⁵ E. Weinberg and A. Wu, Understanding Complex Perturbative Effective Potential, *Phys. Rev. D* 36, 2474 (1987).
- ⁴⁶ S. Coleman and E. Weinberg, Radiative Corrections as the Origin of Spontaneous Symmetry Breaking, *Phys. Rev. D* 7, 1888 (1973).
- ⁴⁷ I. V. Kriev and A. Linde, On the Vacuum Stability Problem in Gauge Theories, *Nucl. Phys. B* 117, 265 (1976).
- ⁴⁸ M. Bando, T. Kugo, N. Maekawa and H. Nakano, Improving the Effective Potential, *Prog. Theor. Phys.* 90, 405 (1993).
- ⁴⁹ C. Ford, Multiscale Renormalization Group Improvement of the Effective Potential, *Phys. Rev. D* 50, 7531 (1994).
- ⁵¹ J. A. Casas, V. D. I. Clemente and M. Quiros, The Effective Potential in the Presence of Several Mass Scales, *Nucl. Phys. B* 553, 511 (1999).
- ⁵¹ M. Quiros, Status of Effective Potential Calculations, hep-ph/9509385.
- ⁵² M. Quiros, Constraints on the Higgs Boson Properties from the Effective Potential, hep-ph/9703412.
- ⁵³ J. A. Casas, V. D. I. Clemente, A. Ibarra and M. Quiros, Massive Neutrinos and the Higgs Boson Mass Window, *Phys. Rev. D* 62, 053005 (2000).
- ⁵⁴ T. Appelquist and J. Carazzone, Infrared Singularities and Massive Fields, *Phys. Rev. D* 11, 2856 (1975).
- ⁵⁵ T. Ohsaku, Supersymmetric Nambu–Jona-Lasinio Model on $N = 1=2$ Four-Dimensional Non(anti)commutative Superspace, *J. High Energy Phys.* 02, 021 (2008).
- ⁵⁶ T. Ohsaku, BCS and Generalized BCS Superconductivity in Relativistic Quantum Field Theory: Formulation, *Phys. Rev. B* 65, 024512 (2002).
- ⁵⁷ T. Ohsaku, BCS and Generalized BCS Superconductivity in Relativistic Quantum Field Theory. II. Numerical Calculations, *Phys. Rev. B* 66, 054518 (2002).
- ⁵⁸ T. Ohsaku, Dynamical Chiral Symmetry Breaking and Superconductivity in the Supersymmetric Nambu–Jona-Lasinio Model at finite Temperature and Density, *Phys. Lett. B* 634, 285 (2006).
- ⁵⁹ T. Ohsaku, Dynamical Chiral Symmetry Breaking, Color Superconductivity, and Bose-Einstein Condensation in an $SU(N_c) \times U(N_f)_L \times U(N_f)_R$ -invariant Supersymmetric Nambu–Jona-Lasinio Model at finite Temperature and Density, *Nucl. Phys. B* 803, 299 (2008).
- ⁶⁰ R. D. Peccei and H. R. Quinn, Constraints Imposed by CP Conservation in the Presence of Pseudoparticles, *Phys. Rev. D* 16, 1791 (1977).