

# The Number of Supertraces on the Superalgebra of Observables of Rational Calogero Model based on the Root System

S.E. Konstein<sup>\*†</sup>

I.E.Tamm Department of Theoretical Physics,  
P. N. Lebedev Physical Institute,  
119991, Leninsky Prospect 53, Moscow, Russia.

and

R. Stekolshchik<sup>‡</sup>

ECI Telecomm Ltd.,  
Osivim 30, Petah-Tikva, Israel.

## Abstract

In the Coxeter group  $W(R)$  generated by the root system  $R$ , let  $Q(R)$  be the number of conjugacy classes having no eigenvalue  $-1$ . The superalgebra  $H_{W(R)}$  of observables of the rational Calogero model based on the root system  $R$  possesses  $Q(R)$  supertraces. The numbers  $Q(R)$  are determined for all irreducible root systems (hence for all root systems).

---

<sup>\*</sup>E-mail: konstein@lpi.ru

<sup>†</sup>This work was supported by the RFBR (grant No. 08-02-01118 (S.K.)), and by the grant LSS-1615.2008.2 (S.K.).

<sup>‡</sup>E-mail: rs2@biu.013.net.il

# 1 The superalgebra of observables

The superalgebra  $H_{W(R)(\nu)}$  of observables of the rational Calogero model based on the root system  $R$  is defined in the following way.

For any nonzero  $\vec{v} \in V = \mathbb{R}^N$  define the reflections  $R_{\vec{v}}$  as follows:

$$R_{\vec{v}}(\vec{x}) = \vec{x} - 2 \frac{(\vec{x}, \vec{v})}{(\vec{v}, \vec{v})} \vec{v} \quad \text{for any } \vec{x} \in V. \quad (1)$$

Here  $(\cdot, \cdot)$  stands for the inner product in  $V$ , i.e.,  $(\vec{x}, \vec{y}) = \sum_{i=1}^N x_i y_i$ , where the  $x_i$  are the coordinates of vector  $\vec{x}$ :  $x_i \stackrel{\text{def}}{=} (\vec{x}, \vec{e}_i)$ , and the vectors  $\vec{e}_i$  constitute an orthonormal basis in  $V$ :  $(\vec{e}_i, \vec{e}_j) = \delta_{ij}$ . The reflections (1) have the following properties

$$R_{\vec{v}}(\vec{v}) = -\vec{v}, \quad R_{\vec{v}}^2 = 1, \quad (R_{\vec{v}}(\vec{x}), \vec{u}) = (\vec{x}, R_{\vec{v}}(\vec{u})), \quad \text{for any } \vec{v}, \vec{x}, \vec{u} \in V.$$

**Definition 1.** *The finite set of vectors  $R \subset V$  is a root system if  $R$  is  $R_{\vec{v}}$ -invariant for any  $\vec{v} \in R$  and the group  $W(R)$  generated by all reflections  $R_{\vec{v}}$  with  $\vec{v} \in R$  (called a Coxeter group) is finite.*

In this paper, we consider noncrystallographic irreducible root systems ( $H_3$ ,  $H_4$  and  $I_2(n)$ ) together with crystallographic ones ( $A_n$ ,  $B_n$ ,  $C_n$ ,  $D_n$ ,  $E_n$ ,  $F_4$ ,  $G_2$ ) [1, 2].

Let  $\mathcal{H}^\alpha$  ( $\alpha = 0, 1$ ) be two copies of  $V$  with orthonormal bases  $a_i^\alpha$  ( $i = 1, \dots, N$ ), respectively. For every vector  $\vec{v} = \sum_{i=1}^N v_i \vec{e}_i \in V$  let  $v^\alpha \in \mathcal{H}^\alpha$  be the vectors  $v^\alpha = \sum_{i=1}^N v_i a_i^\alpha$ , so the bilinear forms on  $\mathcal{H}^0 \oplus \mathcal{H}^1$  can be defined as  $(x^\alpha, y^\beta) = (\vec{x}, \vec{y})$ , where  $\vec{x}, \vec{y} \in V$  and  $x^\alpha, y^\alpha \in \mathcal{H}^\alpha$  are their copies. The reflections  $R_{\vec{v}}$  act on  $\mathcal{H}^\alpha$  as follows

$$R_{\vec{v}}(h^\alpha) = h^\alpha - 2 \frac{(h^\alpha, v^\alpha)}{(\vec{v}, \vec{v})} v^\alpha, \quad \text{for any } h^\alpha \in \mathcal{H}^\alpha.$$

Thus, the  $W(R)$ -action on the spaces  $\mathcal{H}^\alpha$  is defined.

Let  $\nu$  be a set of constants  $\nu_{\vec{v}}$  with  $\vec{v} \in R$  such that  $\nu_{\vec{v}} = \nu_{\vec{w}}$  if  $R_{\vec{v}}$  and  $R_{\vec{w}}$  belong to one conjugacy class of  $W(R)$ .

**Definition 2.**  *$H_{W(R)(\nu)}$  is the associative algebra of polynomials in the  $a_i^\alpha$  with coefficients in the group algebra  $\mathbb{C}[W(R)]$  subject to the relations*

$$\begin{aligned} gh^\alpha &= g(h^\alpha)g \quad \text{for any } g \in W(R), \quad \text{and } h^\alpha \in \mathcal{H}^\alpha \\ [h_1^\alpha, h_2^\beta] &= \varepsilon^{\alpha\beta} \left( (\vec{h}_1, \vec{h}_2) + \sum_{\vec{v} \in R} \nu_{\vec{v}} \frac{(\vec{h}_1, \vec{v})(\vec{h}_2, \vec{v})}{(\vec{v}, \vec{v})} R_{\vec{v}} \right) \quad \text{for any } h_1^\alpha, h_2^\beta \in \mathcal{H}^\alpha, \end{aligned} \quad (2)$$

where  $\varepsilon^{\alpha\beta}$  is the antisymmetric tensor,  $\varepsilon^{01} = 1$ .

This algebra has faithful representation via Dunkl differential-difference operators [3] acting on the space of smooth functions on  $V$ . Namely, let

$$D_i = \frac{\partial}{\partial x_i} + \frac{1}{2} \sum_{\vec{v} \in R} \nu_{\vec{v}} \frac{v_i}{(\vec{x}, \vec{v})} (1 - R_{\vec{v}})$$

and [4, 5]

$$a_i^\alpha = \frac{1}{\sqrt{2}} (x_i + (-1)^\alpha D_i), \quad \alpha = 0, 1. \quad (3)$$

The reflections  $R_{\vec{v}}$  transform the deformed creation and annihilation operators (3) as vectors:

$$R_{\vec{v}}a_i^\alpha = \sum_{j=1}^N \left( \delta_{ij} - 2 \frac{v_i v_j}{(\vec{v}, \vec{v})} \right) a_j^\alpha R_{\vec{v}}.$$

Since  $[D_i, D_j] = 0$  [3], it follows that

$$[a_i^\alpha, a_j^\beta] = \varepsilon^{\alpha\beta} \left( \delta_{ij} + \sum_{\vec{v} \in R} \nu_{\vec{v}} \frac{v_i v_j}{(\vec{v}, \vec{v})} R_{\vec{v}} \right),$$

which manifestly coincides with (2).

The commutation relations (2) suggest to define the *parity*  $\pi$  by setting:

$$\pi(a_i^\alpha) = 1 \text{ for any } \alpha, i, \quad \pi(g) = 0 \text{ for any } g \in W(R)$$

and consider  $H_{W(R)}(\nu)$  as a superalgebra.

We say that  $H_{W(R)}(\nu)$  is the *superalgebra of observables of the Calogero model based on the root system  $R$* .

Obviously,  $\mathbb{C}[W(R)]$  is a subalgebra of  $H_{W(R)}(\nu)$ .

Observe an important property of superalgebra  $H_{W(R)}(\nu)$ : the Lie superalgebra of its inner derivations<sup>1</sup> contains  $\mathfrak{sl}_2$  generated by

$$T^{\alpha\beta} = \frac{1}{2} \sum_{i=1}^N \{a_i^\alpha, a_i^\beta\}$$

which commute with  $\mathbb{C}[W(R)]$ , i.e.,  $[T^{\alpha\beta}, R_{\vec{v}}] = 0$ , and act on  $a_i^\alpha$  as on  $\mathfrak{sl}_2$ -vectors:

$$[T^{\alpha\beta}, a_i^\gamma] = \varepsilon^{\alpha\gamma} a_i^\beta + \varepsilon^{\beta\gamma} a_i^\alpha. \quad (4)$$

The restriction of operator  $T^{01}$  in the representation (3) on the subspace of  $W(R)$ -invariant functions on  $V$  is a second-order differential operator which is the well-known Hamiltonian of the rational Calogero model [6] based on the root system  $R$  [7]. One of the relations (4), namely,  $[T^{01}, a_i^\alpha] = -(-1)^\alpha a_i^\alpha$ , allows one to find the wave functions of the equation  $T^{01}\psi = \epsilon\psi$  via the usual Fock procedure with the vacuum  $|0\rangle$  such that  $a_i^0|0\rangle=0$  for any  $i$  [5]. After  $W(R)$ -symmetrization these wave functions become the wave functions of Calogero Hamiltonian.

## 2 Supertraces on $H_{W(R)}(\nu)$

**Definition 3.** Any linear complex-valued function  $str(\cdot)$  on the superalgebra  $\mathcal{A}$  such that

$$str(fg) = (-1)^{\pi(f)\pi(g)} str(gf)$$

for any  $f, g \in \mathcal{A}$  with definite parity  $\pi(f)$  and  $\pi(g)$  is called a *supertrace*.

---

<sup>1</sup>Let  $\mathcal{A}$  be arbitrary associative superalgebra. Then, the operators  $\mathcal{D}_x$  which act on  $\mathcal{A}$  via  $\mathcal{D}_x(y) = [x, y]$  (supercommutator) constitute the Lie superalgebra of *inner* derivations.

The restriction of every supertrace on  $\mathbb{C}[W(R)]$  is completely determined by its values on  $W(R) \subset \mathbb{C}[W(R)]$  and the function  $str$  is a central function on  $W(R)$ , i.e., it is constant on the conjugacy classes.

Let us introduce the grading  $E$  on the vector space of  $\mathbb{C}[W(R)]$ . Consider the subspace  $\mathcal{E}(g) \subset V$ :

$$\mathcal{E}(g) = \{x \in V \mid gx = -xg\} \text{ for } g \in W(R).$$

Set

$$E(g) = \dim \mathcal{E}(g).$$

The following theorem was proved in [8]

**Theorem 1.** *The number of linearly independent supertraces  $Q(R)$  on  $H_{W(R)}(\nu)$  does not depend on  $\nu$  and is equal to the number of conjugacy classes in  $W(R)$  with  $E(g) = 0$ .*

This Theorem helps to find the number  $Q(R)$  for an arbitrary root system  $R$ .

Theorem 1 implies, evidently, the following statement

**Theorem 2.**

$$Q(R_1 \oplus R_2) = Q(R_1)Q(R_2).$$

Therefore, the problem of finding  $Q(R)$  is reduced to the problem of finding  $Q(R)$  for irreducible root systems  $R$ .

### 3 The numbers $Q(R)$ for irreducible root systems

$R$	$Q(R)$	for the proof, see
$A_{n-1}$	the number of partitions of $n$ into a sum of odd positive integers	[9]
$B_n, C_n, BC_n$	the number of partitions of $n$ into a sum of positive integers	[10]
$D_n$	the number of partitions of $n$ into a sum of positive integers with even number of even integers	[10]
$E_6$	9	Appendix 1
$E_7$	12	Appendix 2
$E_8$	30	Appendix 3
$F_4$	9	Appendix 4
$G_2$	3	[11]
$H_3$	4	Appendix 5
$H_4$	20	Appendix 6
$I_2(n)$	$\left\lfloor \frac{n+1}{2} \right\rfloor$	[11]

# Appendices

## Appendix 1. $E_6$

The conjugacy classes of the Weyl group  $E_6$  are described in Table 9 of [12].  
The following 9 classes have not the root  $-1$ :

$$\phi, A_2, A_2^2, A_4, D_4(a_1), A_2^3, E_6, E_6(a_1), E_6(a_2).$$

## Appendix 2. $E_7$

The conjugacy classes of the Weyl group  $E_7$  are described in Table 10 of [12].  
The following 12 classes have not the root  $-1$ :

$$\phi, A_2, A_2^2, A_4, D_4(a_1), A_2^3, A_4 \times A_2, A_6, D_6(a_1), E_6, E_6(a_1), E_6(a_2).$$

## Appendix 3. $E_8$

The conjugacy classes of the Weyl group  $E_8$  are described in Table 11 of [12].  
The following 30 classes have not the root  $-1$ :

$$\begin{aligned} &\phi, A_2, A_2^2, A_4, D_4(a_1), A_2^3, A_4 \times A_2, A_6, D_4(a_1) \times A_2, D_6(a_1), \\ &E_6, E_6(a_1), E_6(a_2), A_2^4, A_4^2, A_8, D_4(a_1)^2, D_8(a_1), D_8(a_3), \\ &E_6 \times A_2, E_6(a_2) \times A_2, E_8, E_8(a_i) \ (i = 1, \dots, 8). \end{aligned}$$

## Appendix 4. $F_4$

The conjugacy classes of the Weyl group  $F_4$  are described in Table 8 of [12].  
The following 9 classes have not the root  $-1$ :

$$\phi, A_2, \tilde{A}_2, B_2, A_2 \times \tilde{A}_2, D_4(a_1), B_4, F_4, F_4(a_1).$$

## Appendix 5. $H_3$

Let  $k = (\sqrt{5} + 1)/2$ . Then the reflections

$$\begin{aligned} a &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ b &= \frac{1}{2} \begin{pmatrix} 1 & k & k-1 \\ k & 1-k & -1 \\ k-1 & -1 & k \end{pmatrix} \\ c &= \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \end{aligned}$$

corresponding to the roots  $\vec{e}_2$ ,  $\frac{1}{2}(-\vec{e}_1 + k\vec{e}_2 + k^{-1}\vec{e}_3)$  and  $\vec{e}_1$  satisfy the relations

$$a^2 = b^2 = c^2 = 1, \quad (ab)^5 = (bc)^3 = (ac)^2 = 1$$

and define the Coxeter group  $H_3$ .

As  $H_3 = S_5^e \times C_2$  (See [1], Ch.VI, Sect. 4, Ex.11 d), p. 284; [13], p.160 and references there), where  $S_5^e$  is the group of even permutations of 5 elements,  $C_2 = \{1, -1\}$ , ( $|S_5^e| = 60$ ,  $|H_3| = 120$ ),  $H_3$  has 10 conjugacy classes, 5 with positive determinant and 5 with negative one.

The conjugacy classes with positive determinant are described by their representatives

The representative	The characteristic polynomial
1	$(1 - t)^3$
$ac$	$(1 - t)(1 + t)^2$
$bc$	$(1 - t)(t^2 + t + 1)$
$ab$	$(1 - t)[t^2 + (1 - k)t + 1]$
$abab$	$(1 - t)(t^2 + kt + 1)$

Each of this characteristic polynomials has the root  $+1$ , and so each characteristic polynomials of any conjugacy class with negative determinant has the root  $-1$ . The conjugacy class with representative  $ac$  also has the root  $-1$ .

So, the number of conjugacy classes without root  $-1$  is equal to 4.

## Appendix 6. $H_4$

According to [13], all 34 conjugacy classes of  $H_4$  are described by their representatives acting on the space of quaternions

$$g_{lr} : x \mapsto lxr^* \tag{A6.1}$$

and

$$g_p^* : x \mapsto px^*. \tag{A6.2}$$

and all 25 pairs of unit quaternions  $l$  and  $r$  and 9 unit quaternions  $p$  are listed in Table 3 of [13].

Each operator (A6.2) has the root  $-1$ . Indeed, the equation  $px^* = -x$  has nonzero solution  $x = -1 + p$  if  $p \neq 1$ , and  $x$  is an arbitrary imaginary quaternion if  $p = 1$ .

Each operator (A6.1) has not the root  $-1$  if and only if

$$l_0 + r_0 \neq 0. \tag{A6.3}$$

Indeed, the determinant of the map  $x \mapsto lx + xr$  is equal to  $4(l_0 + r_0)^2$ .

There are 20 pairs of  $l, r$  in Table 3 of [13] satisfying the condition (A6.3), namely,  $K_i$  with  $i = 1, 4, 6, 8, 10 - 25$ .

## References

- [1] N.Bourbaki, "Groupes et algèbres de Lie", ch.IV-ch.VI, Hermann, Paris, 1968.
- [2] James E.Humphreys, "Reflection groups and Coxeter groups", Cambridge University Press, 1990.
- [3] C.F.Dunkl, Trans. Am. Math. Soc. **311** (1989) 167.
- [4] A. Polychronakos, Phys. Rev. Lett. **69** (1992) 703.
- [5] L. Brink, H. Hansson and M.A. Vasiliev, Phys. Lett. **B286** (1992) 109.
- [6] F. Calogero, J. Math. Phys., **10** (1969) 2191, 2197; *ibid* **12** (1971) 419.
- [7] M. A.Olshanetsky and A. M. Perelomov, Phys. Rep., **94** (1983) 313.
- [8] S.E. Konstein, Supertraces on the Superalgebra of Observables of Rational Calogero Model based on the Root System, arXiv:hep-th/9803213.
- [9] S.E. Konstein and M.A. Vasiliev, J. Math. Phys. **37** (1996) 2872.
- [10] S.E. Konstein, Teor. Mat. Fiz., **111** (1997) 252-262.
- [11] S.E. Konstein, Rational Calogero models based on rank-2 root systems: supertraces on the superalgebras of observables, arXiv:math/9801001.
- [12] R.W.Carter, Conjugacy classes in the weyl group, Compositio Mathematica, **25** n°1, (1972) 1-59.
- [13] Larry C.Grove, "The characters of the hecatonicosahedroidal group", Journal für die reine und angewandte Mathematik, **265** (1974) 160-169.