Levy ights, dynamical duality and fractional quantum mechanics

Piotr G arbaczewski
Institute of Physics, University of Opole, 45-052 Opole, Poland
February 21, 2024

A bstract

We discuss dual time evolution scenarios which, albeit running according to the same real time clock, in each considered case may be mapped among each other by means of a suitable analytic continuation in time procedure. This dynamical duality is a generic feature of di usion-type processes. Technically that involves a familiar transformation from a non-Hermitian Fokker-Planck operator to the Hermitian operator (e.g. Schrodinger Hamiltonian), whose negative is known to generate a dynamical semigroup. Under suitable restrictions upon the generator, the semigroup admits an analytic continuation in time and ultimately yields dual motions. We analyze an extension of the duality concept to Levy ights, free and with an external forcing, while presuming that the corresponding evolution rule (fractional dynamical semigroup) is a dual counterpart of the quantum motion (fractional unitary dynamics).

PACS numbers: 02.50 Ey, 05.20.-y, 05.40 Jc

1 Brownian motion inspirations

1.1 Diusion-type processes and dynamical semigroups

The Langevin equation for a one-dimensional stochastic division process in an external conservative force eld $F = (r \ V): \underline{x} = F(x) + \frac{p}{2D}b(t)$, where b(t) stands for the normalized white noise hb(t)i = 0, $hb(t^0)b(t)i = (t t^0)$, gives rise to the corresponding Fokker-Planck equation for the probability density -(x;t):

$$Q_{+}^{-} = D - r (F -) :$$
 (1)

By m eans of a standard substitution $(x;t) = (x;t) \exp[V(x)=2D]$, [1], we pass to a generalized di usion equation for an auxiliary function (x;t):

$$\theta_t = D \qquad V(x)$$
 (2)

P resented at the 21 M arian Sm oluchow ski Sym posium on Statistical Physics

where a compatibility condition $V(x) = (1-2)[(F^2-2D) + rF]$ needs to be respected. This transform ation assigns the role of the dynamics generator to the Hermitian (eventually self-ad-pint) operator H = D V.

Under suitable restrictions upon V(x), \hat{H} becomes a legitim ate generator of a contractive dynamical semigroup $\exp(\hat{H}t)$, t=0. If additionally the dynamical semigroup is amenable to an analytic continuation in time, the contractive semigroup operator $\exp(\hat{H}t)$ can be related with the unitary operator $\exp(\hat{H}t)$ via so-called \hat{W} ick rotation t! it. This duality observation underlies our forthcoming discussion and generalizations to Levy ights fram ework.

1.2 Free propagation and its analytic continuation in time

The standard theory of G aussian di usion-type processes takes the W iener process as the "free noise" model, with the Laplacian as the "noise" generator. It is an element of folk lore that the related dissipative sem igroup dynam ics exp (tD) = exp (t $\hat{H_0}$) (and thus the heat equation) can be m apped into the unitary dynam ics exp (itD) = exp (it $\hat{H_0}$) (and thus the free Schrodinger equation), by m eans of an analytic continuation in time procedure, [2]. A parameter D m ay be interpreted dimensionally as D = ~=2m , or D = k_B T=m (E instein's uctuation-dissipation statement).

Quite often, this mapping is represented by a formalit! time transformation of the free Schrodinger picture dynamics (one should be aware that to execute a mapping for concrete solutions, a proper adjustment of the time interval boundaries is necessary):

$$iQ_{+} = D 4 ! Q_{+} = D 4 ; (3)$$

where the notation for solutions of the heat equation has been adopted, to stay in conform ity with the forthcom ing more general discussion, where (x;t) needs not to be a probability density, [2]-[4].

The mapping is usually exemplied in terms of integral kernels g and k as follows, c.f. also [5]:

$$(x;t) = dx^0 g(x x^0;t) (x^0;0)$$
 (4)

g(x
$$x^{0}$$
;t) = k(x x^{0} ; it) = (4 iD t) $^{1=2} \exp\left[-\frac{(x - x^{0})^{2}}{4iD t}\right]$

and

$$Z (x;t) = dx^0k (x x^0;t) (x^0;0) (5)$$

$$k(x x^{0};t) \stackrel{!}{=} g(x x^{0}; it) = (4 D t)^{1=2} \exp\left[-\frac{(x x^{0})^{2}}{4D t}\right];$$

where the initial t = 0 data need to be properly adjusted. Here, $g(x x^0;t)$ is an integral kernel of the unitary evolution operator: [exp(iDt)](x;0) = (x;t). The

heat kernel k (x x^0 ; y) plays the same role with respect to the contractive sem igroup operator: [exp(Dt)](x;0).

The special choice of

$$(x;0) = (^2)^{1=4} \exp \frac{x^2}{2^2}$$
 (6)

im plies

$$(x;t) = \frac{2^{-1-4}}{-} (^{2} + 2iDt)^{-1-2} \exp \frac{x^{2}}{2(^{2} + 2iDt)}$$
 (7)

and

$$(x;t) = (x; it) = -\frac{2^{-1-4}}{2(2+2Dt)^{-1-2}} \exp \frac{x^2}{2(2+2Dt)^{-0}}$$
 (8)

 $w \pm h \quad (x;0) = (x;0).$

We note that $= j \hat{j} = is$ a quantum mechanical probability density on R for all times

$$(x;t) = \frac{2}{(4+4D^2t^2)} \exp \frac{2x^2}{4+4D^2t^2} :$$
 (9)

The real solution (x;t) of the heat equation is not a probability density $\overline{}(x;t) = (x;t)$, unless multiplied by an appropriate real function (x;t) which solves the time adjoint heat equation (that becomes an ill-posed dynamical problem if considered carelessly).

Case 1: Since $(x;t) = [2 (^2 + 2D t)]^{1=2} \exp[x^2=2(^2 + 2D t)]$ actually is an example of the free B rownian motion probability density for all t = 0, we infer

$$-(x;t) = (4^{2})^{1-4} \quad (x;t) = (10)$$

where $(x;t) = (4^{-2})^{1-4}$ is interpreted as a trivial (constant) solution of the time adjoint heat equation $\theta_t = D$. We stress that $= (4^{-2})^{-1-4}$. This, booking redundant observation, will prove quite useful in a more general fram ework to be introduced in below.

Case 2: A complex conjugate (x;t) = (x;t) of (x;t), Eq. (7), solves the time-adjoint Schrodinger equation $i e_t = D$. Hence a time-symmetric approach to the analytic continuation in time might look more compelling. Indeed

$$(x;t) = (x;it) = -\frac{2^{-1-4}}{2} (2Dt)^{-1-2} \exp \frac{x^2}{2(2Dt)}$$
 (11)

is a legitim ate solution of the time-adjoint heat equation $@_t = D$ as long as t2 [T=2; + T=2] where $T={}^2=D$.

In the present case, both time adjoint equations set well de ned Cauchy problems (at least in the just de ned time interval). The subtle point is that the would be

"initial" data for the backward in time evolution, in fact need to be the term inal data, given at the end-point T=2 of the considered time-interval.

The only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands, is the heat kernel (3): $k(x x^0;t!)$ the only propagation tool, we have in hands (3): $k(x x^0;t!)$ the only propagation tool, we have in hands (3): $k(x x^0;t!)$ the only propagation tool, we have in hands (3): $k(x x^0;t!)$ the hands (3): $k(x x^0;t!)$ the hands (3): $k(x x^0;t!)$ the h

The original quantum mechanical probability density = $j \hat{j}$ = , Eq. (7), is mapped into a Brownian bridge (pinned Brownian motion) probability density:

(x; it) =
$$\frac{1}{2}$$
 (x;t) = () (x;t) = $\frac{2}{(4 + 4D^2t^2)}$ exp $\frac{2}{4}$ x² (12)

The price paid for the time-symmetric appearance of this formula is a limitation of the admissible time span to a nite time-interval of length $T = {}^{2}=D$.

Case 3: To make a direct comparison of Case 2 with the previous Case 1, let us con ne the time interval of Case 2 to [0;+T=2]. Now, a conditional Brownian motion connects $\overline{}(x;0)=(x;0)=(^2)^{1=2}\exp(x^2=^2)$ with $\overline{}(x;t!+T=2)$ of Eq. (10). Because of $T=^2=D$, as t! T=2, instead of a regular function we arrive at the linear functional (generalized function), here represented by the D irac delta (x). Note that $(x + x^0)$ is a standard initial t=0 value of the heat kernel t=0.

This behavior is faithfully reproduced by the time evolution of (x;t) and (x;t) that compose $\overline{}(x;t) = ()(x;t)$ for t 2 [0;T=2]. The initial value of (x;0) = (x;0), Eq. (6), is propagated forward in accordance with Eq. (8) to $(x;T=2) = (4 \quad 2)^{1-4} \exp(x^2-4 \quad 2)$.

In parallel, (x;t) of (11) interpolates backwards between (x;T=2) $(4^2)^{1=4}$ (x) and (x;0) = (x;0). We have here employed an identity (ax) = (1=jaj) (x). Because of f(x) (x) f(0) (x), we arrive at (x;T=2) = (x) (x;T=2) (x).

1.3 Schrodinger's boundary data problem

The above discussion provides particular solutions to so-called Schrodinger boundary data problem, under an assum ption that a M arkov stochastic process which interpolates between two a priorigiven probability densities (x;0) and (x;T=2) can be modeled by means of the G auss probability law (e.g. in terms of the heat kernel). That incorporates the free Brownian motion (W interprocess) and all its conditional variants, Brownian bridges being included, [3,4] and [6]-[8], c.f. also [2].

For our purposes the relevant information is that, if the interpolating process is to display the Markov property, then it has to be specified by the joint probability measure (A and B are Borel sets in R):

$$m (A;B) = dx dym (x;y)$$
(13)

where $_{R}^{R}$ m (x;y)dy = (x;0); and $_{R}^{R}$ m (x;y)dx = (y;T=2). From the start, we assign densities to all m easures to be dealt with, and we assume the functional form of the density m (x;y)

$$m(x;y) = f(x)k(x;0;y;T=2)g(y)$$
 (14)

to involve two unknown functions f(x) and g(y) which are of the same sign and nonzero, while k(x;s;y;t) is any bounded strictly positive (dynamical semigroup) kernel dened for all times 0 s < t T=2. For each concrete choice of the kernel, the above integral equations are known to determine functions f(x); g(y) uniquely (up to constant factors).

By denoting (x;t) = R f (z)k(z;0;x;t)dz and (x;t) = R k (x;t;z;T=2)g(z)dz it follows that

$$-(x;t) = (x;t) (x;t) = p(y;s;x;t) - (y;s)dy;$$
 (15)

$$p(y;s;x;t) = k(y;s;x;t) - \frac{(x;t)}{(y;s)};$$

for all 0 s < t T=2. The above p(y;s;x;t) is the transition probability density of the pertinent M arkov process that interpolates between -(x;0) and -(x;T=2). Cases 1 through 3 are particular examples of the above reasoning, once k(x;s;y;t) is specified to be the heat kernel (3) and the corresponding boundary density data are chosen. Clearly, (x;0) = f(x) while (x;T=2) = g(x).

We recall that in case of the free evolution, by setting (x;t) = const, as in Case 1, we exclively transform an integral kernel k of the L¹ (R) norm-preserving sem igroup into a transition probability density p of the M arkov stochastic process. Then -.

2 Free noise models: Levy ights and fractional (Levy) sem igroups

The Schrodinger boundary data problem is amenable to an immediate generalization to in nitely divisible probability laws which induce contractive semigroups (and their kernels) for general Gaussian and non-Gaussian noise models. They allow for various jump and jump-type stochastic processes instead of a diusion process.

A subclass of stable probability laws contains a subset that is associated in the literature with the concept of Levy ights. At this point let us invoke a functional analytic lore, where contractive sem igroup operators, their generators and the pertinent integral kernels can be directly deduced from the Levy-K hitchine form ula, compare e.g. [8].

Let us consider sem igroup generators (H am iltonians, up to dim ensional constants) of the form $\hat{H} = F$ (\hat{p}), where $\hat{p} = F$ ir stands for the momentum operator (up to the

disregarded \sim or 2m D factor) and for 1 < k < +1, the function F = F(k) is real valued, bounded from below and locally integrable. Then,

$$\exp(t\hat{H}) = \exp[tF(k)]dE(k)$$
 (16)

where t 0 and dE (k) is the spectral measure of p.

Because of

$$(E (k)f)(x) = \frac{1}{2} \sum_{k=1}^{Z_k} \exp(ipx)f(p)dp$$
 (17)

where f is the Fourier transform of f, we learn that

$$Z_{+1}$$
 [exp(th)]f(x) = [exp(tF(k))dE(k)f](x) = (18)

$$\frac{1}{p + 1} = \exp[tF(k)] \frac{d}{dk} \left[\exp(ipx)f(p)dp\right] dk = (22)$$

$$\frac{1}{p + 1} = \exp(tF(k)) \exp(ikx)f(k) dk = \exp(tF(p))f(p) + (x)$$

where the superscript _ denotes the inverse Fourier transform.

Let us set

$$k_t = \frac{1}{2} [exp(t F(p))]$$
: (19)

Then the action of exp ($t\hat{H}$) can be given in terms of a convolution (i.e. by means of an integral kernel k_t k (x y;t) = k (y;0;x;t)):

$$\exp(t\hat{H})f = [\exp(tF(p))f(p)] = fk$$
 (20)

w here

(f g) (x)
$$=$$
 g (x z) f (z) dz: (21)

We shall restrict considerations only to those F (p) which give rise to positivity preserving sem igroups: if F (p) satisfies the celebrated Levy-K hintchine formula, then k_t is a positive measure for all t 0. The most general case refers to a combined contribution from three types of processes: deterministic, Gaussian, and the jump-type process.

We recall that a characteristic function of a random variable X completely determines a probability distribution of that variable. If this distribution admits a density we can write E [exp(ipX)] = $_{\rm R}^{-}$ (x) exp(ipx)dx which, for in nitely divisible probability laws, gives rise to the Levy-K hintchine formula

E [exp(ipX)] = expfi p (
2
=2)p 2 + 2 [exp(ipy) 1 $\frac{ipy}{1+y^{2}}$] (dy)g (22)

where $\mbox{(dy)}$ stands for the so-called Levy measure. In terms of M arkov stochastic processes all that amounts to a decomposition of X $_{\rm t}$ into

$$X_t = t + B_t + J_t + M_t$$
 (23)

where B $_{\rm t}$ stands for the free B rownian motion (W iener process), $J_{\rm t}$ is a Poisson process while M $_{\rm t}$ is a general jump-type process (m ore technically, m artingale with jumps).

By disregarding the determ in istic and jump-type contributions in the above, we are left with the Wiener process $X_t = B_t$. For a Gaussian $(x) = (2^2)^{-12} \exp(x^2 + 2^2)$ we directly evaluate E [exp(ipx)] = exp($x^2 + 2^2 + 2^2$).

Let us set 2 = 2D t. We get E [exp (ipX $_t$)] = exp (tD $_t$) and subsequently, by employing p! p = ir, we arrive at the contractive sem igroup operator exp (tD) where the one-dimensional Laplacian = d^2 =dx² has been introduced. That amounts to choosing a special version of the previously introduced H am iltonian H = F (p) = D p². Note that we can get read of the constant D by rescaling the time parameter in the above.

P resently, we shall concentrate on the integral part of the Levy-K hintchine formula, which is responsible for arbitrary stochastic jump features. By disregarding the determ inistic and B rownian motion entries we arrive at:

$$F(p) = \exp(ipy) \quad 1 \quad \frac{ipy}{1 + y^2}] (dy)$$
 (24)

where (dy) stands for the appropriate Levy measure. The corresponding non-Gaussian Markov process is characterized by

$$E \left[\exp \left(ipX_{t} \right) \right] = \exp \left[tF \left(p \right) \right] \tag{25}$$

with F (p), (22). A coordingly, the contractive sem igroup generator may be defined as follows: F (\hat{p}) = \hat{H} .

For concreteness we can mention some explicit examples of non-Gaussian Markov sem igroup generators. F (p) = p where < 2 and > 0 stands for the intensity parameter of the Levy process, upon p! p = p ir gives rise to a pseudo-di erential operator \hat{H} = p often named the fractional Hamiltonian. Note that, by construction, it is a positive operator (quite alike D).

The corresponding jump-type dynamics is interpreted in terms of Levy ights. In particular

$$F(p) = \dot{p}j! \quad \hat{H} = F(p) = \dot{j}r \quad \dot{j} = \dot{j}r \quad \dot{j}r \quad \dot{j} = \dot{j}r \quad \dot{j$$

refers to the Cauchy process.

Since we know that the probability density of the free Brownian motion is a solution of the Fokker-Planck (here, simply - heat) equation

$$Q_{+}^{-} = D \qquad (27)$$

it is instructive to set in comparison the pseudo-di erential Fokker-Planck equation which corresponds to the fractional Hamiltonian and the fractional semigroup exp(th^) = $\exp(jj^{-2})$

$$Q_{t}^{-} = \dot{j} \dot{j}^{=2-} : \tag{28}$$

Asmentioned in the discussion of Case 1, instead of in the above we can insert —, while remembering that const.

3 Free fractional Schrodinger equation

Fractional H am iltonians $\hat{H} = jj^{=2}$ with < 2 and > 0 are self-adjoint operators in suitable $L^2(R)$ domains. They are also positive operators, so that the respective fractional sem igroups are holomorphic, i. e. we can replace the time parameter t by a complex one = t + is; t > 0 so that

$$\exp \left(\begin{array}{cc} Z \\ \exp \left(\begin{array}{cc} \hat{H} \end{array} \right) = \exp \left(\begin{array}{cc} F(k) \end{array} \right) dE(k) : \tag{29}$$

Its action is de ned by

$$[\exp(\hat{H})]f = [(f \exp(\hat{F})] = f k : (30)$$

Here, the integral kernel reads $k = \frac{1}{2} [\exp(F)]$. Since \hat{H} is selfad joint, the lim it t # 0 leaves us with the unitary group $\exp(is\hat{H})$, acting in the same way: $[\exp(is\hat{H})]f = [f\exp(isF)]$, except that now $k_{is} := \frac{1}{2} [\exp(isF)]$ in general is not a probability measure.

In view of unitarity, the unit ball in L^2 is an invariant of the dynamics. Hence probability densities, in a standard form = can be associated with solutions of the free fractional (pseudodiferential) Schrodinger equations:

$$i\theta_{t} (x;t) = j j^{=2} (x;t)$$
 (31)

with initial data (x;0). Attempts towards formulating the fractional quantum mechanics can be found in Refs. [8, 11, 12, 13].

All that amounts to an analytic continuation in time, in close a nity with the Gaussian pattern (1):

$$iQ_t = jj^{=2}$$
 ! $Q_t = jj^{=2}$ (32)

We assume that and thence the corresponding const.

Stable stochastic processes and their quantum counterparts are plagued by a com-mon disease: it is extremely hard, if possible at all, to produce insightful analytic solutions. To get a avor of intricacies to be faced and the level of technical diculties,

we shall reproduce some observations in regard to the Cauchy dynamical sem igroup and its unitary (quantum) partner. For convenience we scale out a parameter .

For the Cauchy process, whose generator is jr j, we dealwith a probabilistic classics:

$$-(x;t) = \frac{1}{t^2 + x^2} = k (y;s;x;t) = \frac{1}{(t + s)^2 + (x + y)^2}$$
(33)

where 0 < s < t. We have hexp[ipX (t)]i = $\binom{R}{R} \exp(ipx)^{-}(x;t) dx = \exp[tF(p)] = \exp(tpt)$ and

$$-(x;t) = k(y;s;x;t) - (y;s) dy$$
 (34)

for all t> s 0. We recall that $\lim_{t \neq 0} \frac{t}{(x^2 + t^2)}$ (x).

The characteristic function of k(y;s;x;t) for y;s xed, reads $exp[ipy \quad jp](t \quad s)]$, and the Levy measure needed to evaluate the Levy-K hintchine integral reads:

$$_{0} (dy) := \lim_{t \neq 0} \left[\frac{1}{t} k (0;0;y;t) \right] dy = \frac{dy}{y^{2}} :$$
 (35)

To pass to a dual Cauchy-Schrodinger dynam ics, we need to perform an analytic continuation in time. We deal with a holomorphic fractional sem igroup $\exp(-tjr)$, = t+ is, (27). It is clear that $\exp(-tjr)$ and $\exp(-tjr)$ have a common, identity operator $\lim_{n \to \infty} t = 0$.

An analytic continuation of the Cauchy kernel by m eans of (28) gives rise to:

$$k_t(x) = \frac{1}{x^2 + t^2}$$
 ! $g_s(x) = k_{is}(x) = \frac{1}{2}[(x + s) + (x + s)] + \frac{1}{2}P\frac{is}{x^2 + s^2};$ (36)

where P indicates that a convolution of the integral kernel with any function should be considered as a principal value of an improper integral, [8]. This should be compared with an almost trivial outcome of the previous mapping (2)! (3). Here, we employ the usual notation for the Dirac delta functionals, and the new time labels is a remnant of the limiting procedure t # 0 in t = t + t is.

The function denoted by is= $(x^2 s^2)$ com es from the inverse Fourier transform of $\frac{1}{2}$ sgn (p) sin (sp). Because of

$$[sgn (p)] = i \frac{2}{-P} (\frac{1}{x})$$
(37)

where P $(\frac{1}{x})$ stands for the functional de ned in term sofa principal value of the integral. Using the notation s for the D irac delta functional (x s):

$$\begin{array}{ccc}
\text{r} & \underline{} \\
\text{[sin (sp)]} & = i & \frac{1}{2} (s_s & s_s)
\end{array}$$
(38)

we realize that

$$\frac{1}{x^2} \frac{\text{is}}{\text{s}^2} = \frac{\text{i}}{2} \left(s \right) \quad P \left(\frac{1}{x} \right) \tag{39}$$

is given in term s of the implicit convolution of two generalized functions. Obviously, a propagation of an initial function $_{0}$ (x) to time t > 0:

$$(x;t) = \int_{\mathbb{R}}^{\mathbb{R}} g(x \quad x^{0};t) \quad _{0}(x^{0})dx^{0}$$
 (40)

gives a solution of the fractional (C auchy) Schrodinger equation $i\theta_t = jr j$.

In comparison with the Gaussian case of Section 1, one important dierence must be emphasized. The improper integrals, which appear while evaluating various convolutions, need to be handled by means of their principal value. Therefore, a simple it! transform ation recipe no longer works on the level of integral kernels and respective and functions.

One explicit example is provided by the incongruence of (31) and (34) with respect to the formalt! it mapping. Another is provided by considering specic solutions of pseudo-di erential equations (30).

To that end, let us consider $_0(x) = (2=)^{1-2}\frac{1}{1+x^2}$, together with $= (2)^{1-2}$. Then, $(x;0) = \frac{1}{(1+x^2)}$ is an L (R) normalized C auchy density, while $_0(x)$ itself is the L² (R) normalized function. C learly:

$$(x;t) = [exp(t;t)] (x;t) = \begin{cases} z \\ k(y;0;x;t) \end{cases} (y;0)dy = \begin{cases} 2 \\ \frac{1+t}{x^2+(1+t)^2} \end{cases} (41)$$

while the corresponding (x;t) with $_0(x) = _0(x)$ reads (for details see e.g. [8]):

$$(x;s) = [\exp(isjr)]_{0}(x) = \frac{1}{2}[_{0}(x+s) + _{0}(x-s)] +$$

$$\frac{i}{2}[(x-s)_{0}(x-s) - (x+s)_{0}(x+s)] :$$

$$(42)$$

4 Dynam ical duality in external potentials: fractional Schrodinger sem igroups and Levy ights

4.1 Schrodinger sem igroups for Smoluchow skiprocesses

Considerations of Section 1, where the free quantum dynamics and free Brownian motion were considered as dual dynamical scenarios, can be generalized to an externally perturbed dynamics, [2]. Namely, one knows that the Schrodinger equation for a quantum particle in an external potential V(x), and the generalized heat equation are connected by analytic continuation in time, known to take the Feynman-Kac (holomorphic semigroup) kernel into the Green function of the corresponding quantum mechanical problem.

$$i\theta_t = D + V ! \theta_t = D V :$$
 (43)

Here V = V (x) = 2m D.

For V = V(x); $x \ge R$, bounded from below, the generator $\hat{H} = 2m D^2 4 + V$ is essentially selfad joint on a natural dense subset of L^2 , and the kernel $k(x;s;y;t) = [exp[(t s)\hat{H})](x;y)$ of the related dynam ical sem igroup $exp(t\hat{H})$ is strictly positive. The quantum unitary dynam ics $exp(t\hat{H})$ is the an obvious result of the analytic continuation in time of a dynam ical sem igroup.

A ssum ptions concerning the admissible potential may be relaxed. The necessary demands are that \hat{H} is self-adjoint and bounded from below. Then the respective dynamical semigroup is holomorphic.

The key role of an integral kernel of the dynam ical sem igroup operator has been elucidated in formulas (11)-(13), where an explicit form of a transition probability density of the M arkov di usion process was given. We have determined as well the time development of (x;t) and (x;t), so that (x;t) = (x;t) is a probability density of the pertinent process.

If we a priori consider (x;t) in the functional form $(x;t) = \exp(x;t)$, so that $(x;t) = (x;t) \exp(x;t)$, and properly de ne the forward drift b(x;t) = 2D r (x;t) in the pertinent Fokker-P lanck equation:

$$Q_{t}^{-} = D - r (b^{-})$$
 (44)

we can recast a di usion problem in terms of a pair of time adjoint generalized heat equations

$$Q_{+} = D \qquad V \tag{45}$$

and

$$Q_{+} = D + V ; \qquad (46)$$

i. e. as the Schrödinger boundary data problem, where an interpolating stochastic process is uniquely determined by a continuous and positive Feynman-Kac kernel of the Schrödinger sem igroup exp (th $\hat{}$), where H $\hat{}$ = D + V.

If our departure point is the Fokker-Planck (or Langevin) equation with the a priori prescribed potential function (x;t) for the forward drift b(x;t), then the backward equation (44) becomes an identity from which V directly follows, in terms of and its derivatives, [6, 7]:

$$V(x;t) = \theta_t + \frac{1}{2}(\frac{b^2}{2D} + rb)$$
 (47)

For the time-independent drift potential, which is the case for standard Smoluchowski di usion processes, we get (c.f. also [1], where the a transformation of the Fokker-Planck equations (42) into an associated Herm itian problem (43) is described in detail):

$$V(x) = \frac{1}{2} \left(\frac{b^2}{2D} + rb \right)$$
 (48)

Notice that (x) is de ned up to an additive constant.

To give an example of a pedestrian reasoning based on the above procedure in case of a concrete Smoluchowski di usion processes, let us begin from the Langevin equation for the one-dimensional stochastic process in the external conservative force $\operatorname{eld} F(x) = (r \ V)(x)$ (to keep in touch with the previous notations, note that V):

$$\frac{dx}{dt} = F(x) + \frac{P}{2DB}(t)$$
 (49)

where B (t) stands for the normalized white noise: hB (t) i = 0, hB (t) i

$$Q_{t}^{-} = D - r (F -)$$
 (50)

and by means of a substitution $(x;t) = (x;t) \exp[V(x)=2D]$, [1], can be transformed into the generalized di usion equation for an auxiliary function (x;t):

$$Q_{t} = D \qquad V \tag{51}$$

where the consistency condition (reconciling the functional form of V with this for F)

$$V = \frac{1}{2} \frac{F^2}{2D} + rF : (52)$$

directly comes from the time-adjoint equation

$$Q_{+} \qquad 0 = D \qquad + V \tag{53}$$

with $(x) = \exp[V(x)=2D]$.

For the 0 mstein-U hlenbeck process b(x) = F(x) = x and accordingly

$$V(x) = \frac{{}^{2}x^{2}}{4D} = \frac{1}{2} : (54)$$

is an explicit functional form of the potential V, present in previous form ulas (41)-(44).

4.2 Fractional sem igroups and perturbed Levy ights

External perturbations in the additive form:

$$i\theta_t (x;t) = jj^{-2} (x;t) + V(x) (x;t)$$
 (55)

were considered in the fram ework of fractional quantum mechanics, [11]-[13], c.f. also [8, 9]. With the dual dynamics concept in mind, Eq. (30), we expect that an analytic continuation in time (if admitted) takes us from the fractional Schrodinger equation to the fractional analog of the generalized di usion equation:

$$Q_t = jj^{-2} V :$$
 (56)

The time-adjoint equation has the form

$$Q_t = \dot{j} \dot{j}^{=2} + V ; \qquad (57)$$

We shall be particularly interested in the time-independent (x;t) (x), an assum ption a neto that involved in the passage from (44)-(46).

Herm itian fractional problems of the form (48) and/or (49) have also been studied in Refs. [14, 15, 16]. However, the major (albeit implicit, never openly stated) assumption of Refs. [14, 15, 16] was to consider the so-called step Levy process instead of the jumptype Levy process proper.

This amounts to introducing a lower bound on the length of admissible jumps: arbitrarily small jumps are then excluded. That allows to by-pass a serious technical obstacle. Indeed, for a pseudo-di erential operator =2, the action on a function from its domain can be greatly simplied by disregarding jumps of length not exceeding a xed > 0, see e.g. Refs. §, 9]:

Compare e.g. Eq. (2) in [15] and Eq. (6) in [16]. Note that these Authors have skipped the minus sign that must appear on the right-hand-side of both formulas (50).

As a side comment, let us point out that the principal integral value issues of Section 3 would not arise in our previous discussion of Cauchy ights and their generators, if arbitrarily small jumps were eliminated from the start. Nonetheless, if the #0 limit is under control, the step process can be considered as a meaningful approximation of the fully-edged (perturbed) jump-type Levy process. This approximation problem has been investigated in detail, in the construction of the perturbed Cauchy process, governed by the Hermitian dynamical problem (53), with the input (55), under suitable restrictions on the behavior of V, [9].

Let us come back to time-adjoint fractional equations (54) and (55). We have -(x;t) = ()(x;t) and employ the trial ansatz of Section 4.2:

$$(x;t)$$
 $(x) = \exp[(x)]$ (59)
 $(x;t) = -(x;t) \exp[(x)]$:

A coordingly (55) in plies, com pare e.g. [14] for an independent argum ent:

$$V = \exp() j j^{-2} \exp()$$
 (60)

to be compared with Eq. (8) in Ref. [15]. In view of (54) we have

$$Q_t^- = Q_t = \exp()[jj^{-2}\exp()] + V^{-2} = rj;$$
 (61)

Langevin-style description of perturbed Levy ights (determ in istic component plus the Levy noise contribution) are known, [17, 18, 19], to generate fractional Fokker-Planck equations of the form

$$Q_{t}^{-} = r (F^{-}) \qquad j j^{=2-} \stackrel{:}{=} r j \qquad (62)$$

where F = r V r, we face problem swhich are left unsettled at the present stage of our investigation:

- (i) M ay the stochastic processes driving (58) and/or (59) coincide under any circum stances, or basically not at all?
- (ii) G ive an insightful/useful de nition of the probability current j(x;t) in both considered cases, while remembering that for fractional derivatives the composition rule for consecutive (R iesz) derivatives typically breaks down.

Both problems (i) and (ii) have have an immediate resolution in case of di usion-type processes, where by departing from the Langevin equation one infers Fokker-Planck and continuity equations. In turn, these equations can be alternatively derived by means of the Schrodinger boundary data problem, provided its integral kernel stems from thee Schrodinger sem igroup, both in the free and perturbed cases. The stochastic di usion process (corresponding to that associated with the Langevin equation) is then reconstructed as well. Thence, the Schrodinger loop gets closed.

W hile passing to Levy processes, we have demonstrated that, with suitable reservations, this Schrodinger "loop" can be completed in case of free Levy ights. However, the "loop" remains incomplete (neither de nitely proved or disproved) for perturbed Levy ights.

At this point we should mention clear indications [14] that, once discussing Levy ights, we actually encounter two di erent classes of processes with incompatible dynamical properties. One class is related to the Langevin equation, another – term ed topological – relies on the "potential landscape" provided by the elective potential V(x). An extended discussion of the latter problem has been postponed to the forthcoming paper, c.f. [23].

A cknow ledgem ent: Partial support from the Laboratory for Physical Foundations of Inform ation Processing is gratefully acknow ledged.

R eferences

- [1] H.Risken, The Fokker-Planck equation, Sringer-Verlag, Berlin, 1989
- [2] P.G arbaczewski, Phys. Rev. E 78, 031101, (2008)
- [3] J-C. Zam brini, J. Math. Phys. 27, 2307. (1986)
- [4] J.C. Zambrini, Phys. Rev. A 35, 3631, (1987)
- [5] M .S.W ang, Phys.Rev.A 37, 1036, (1988)
- [6] Ph.Blanchard and P.Garbaczewski, Phys. Rev. E 49, 3815, (1994)
- [7] P.G arbaczewski and R.O lkiewicz, J.M ath. Phys. 37, 732, (1996)
- [8] P.G arbaczew ski, J.R.K lauder and R.O Ikiew icz, Phys. Rev. E 51, 4114, (1995)
- [9] P.G arbaczewski and R.O lkiewicz, J.M ath. Phys. 40, 1057, (1999)
- [10] P.G arbaczewski and R.O Ikiewicz, J.M ath. Phys. 41, 6843, (2000)
- [11] N. Laskin, Phys. Rev. E 62,3135, (2000)
- [12] N. Laskin, Phys. Rev. E 66, 056108, (2002)
- [13] N. Cufaro Petroni and M. Pusterla, Physica A 388, 824, (2009)
- [14] D.Brockmann and I.Sokolov, Chem. Phys. 284, 409, (2002)
- [15] D. Brockmann and T. Geisel, Phys. Rev. Lett. 90, 170601, (2003)
- [16] D. Brockmann and T. Geisel, Phys. Rev. Lett. 91, 048303, (2003)
- [17] S. Jespersen, R. Metzler and H. C. Fogedby, Phys. Rev. E 59, 2736, (1999)
- [18] P.D.Ditlevsen, Phys. Rev. E 60, 172, (1999)
- [19] V.V. Janovsky et al., Physica A 282, 13, (2000)
- [20] A. Chechkin et al., Chem. Phys. 284, 233, (2002)
- [21] A.A.Dubkov and B. Spagnolo, Acta Phys. Pol. A 38, 1745, (2007)
- [22] A.A.Dubkov, B. Spagnolo and V.V. Uchaikin, Int. J. Bifurcations and Chaos, 18, 2549, (2008)
- [23] P.G arbaczewski, Levy ights and Levy-Schrodinger sem igroups, arX iv:0902.3536