

GENERALIZED LANDAU-LIFSHITZ SYSTEMS AND LIE ALGEBRAS ASSOCIATED WITH HIGHER GENUS CURVES

S. IGONIN, J. VAN DE LEUR, G. MANNO, AND V. TRUSHKOV

Abstract. The Wahlquist-Estabrook prolongation method allows to obtain for some PDEs a Lie algebra that is responsible for Lax pairs and Backlund transformations of certain type. We study the Wahlquist-Estabrook algebra of the n -dimensional generalization of the Landau-Lifshitz equation and construct an epimorphism from this algebra onto an infinite-dimensional quasigraded Lie algebra $L(n)$ of certain matrix-valued functions on an algebraic curve of genus $1 + (n-3)2^{n-2}$. For $n = 3; 4; 5$ we prove that the Wahlquist-Estabrook algebra is isomorphic to the direct sum of $L(n)$ and a 2-dimensional abelian Lie algebra. Using these results, for any n a new family of Miura type transformations (differential substitutions) parametrized by points of the above mentioned curve is constructed. As a by-product, we obtain a representation of $L(n)$ in terms of a finite number of generators and relations, which may be of independent interest.

1. Introduction

In the last 25 years it has been well understood how to construct integrable PDEs from infinite-dimensional Lie algebras (see, e.g., [6, 7, 9, 22, 25] and references therein). The present paper addresses the inverse problem: given a system of PDEs, how to determine whether it is related to an infinite-dimensional Lie algebra and how to recover this Lie algebra?

A partial answer to this question is provided by the so-called Wahlquist-Estabrook prolongation method [4, 26], which is an algorithmic procedure that for a given $(1+1)$ -dimensional system of PDEs constructs a Lie algebra called the Wahlquist-Estabrook algebra (or the WE algebra in short). The WE algebra is responsible for Lax pairs and Backlund transformations of certain type. The method gives this algebra in terms of generators and relations. For some PDEs the explicit structure of the WE algebra was revealed, and as a result one obtained interesting infinite-dimensional Lie algebras (see, e.g., [3, 5, 14, 19] and references therein).

In the original method of Wahlquist and Estabrook the obtained algebras lack any invariant coordinate-free meaning. Recently the WE algebra has been included in a sequence of Lie algebras that have a remarkable geometric interpretation: they are the analogue of the topological fundamental group for the category of PDEs [11, 12]. All infinite-dimensional quotients of these Lie algebras have a coordinate-free meaning as symmetry algebras of certain coverings of PDEs [11] (the notion of coverings of PDEs by Krasilshchik and Vinogradov [2, 15] is a far-reaching geometric generalization of Wahlquist-Estabrook pseudopotentials). Moreover, an effective necessary condition for existence of a Backlund transformation connecting two given PDEs has been obtained in terms of these Lie algebras [11], which has allowed to prove for the first time ever that some PDEs are not connected by any Backlund transformation [11]. In our opinion, these results strongly suggest to compute and study the WE algebras for more PDEs.

1991 Mathematics Subject Classification. 37K30, 37K35.

Key words and phrases. Wahlquist-Estabrook prolongation structures, generalized Landau-Lifshitz systems, quasigraded infinite-dimensional Lie algebras, Miura type transformations.

In this paper we apply the Wahlquist-Estabrook method to the following system

$$(1) \quad S_t = S_{xx} + \frac{3}{2} h S_x; S_x i S_x + \frac{3}{2} h S; R S i S_x; \quad h S; S i = 1;$$

where $S = (s^1(x;t); \dots; s^n(x;t))$ is a column n -vector of dimension $n > 1$, $h; i$ is the standard scalar product, and $R = \text{diag}(r_1; \dots; r_n)$ is a constant diagonal matrix with $r_i \notin r_j$ for $i \notin j$. This system was introduced in [8] and possesses a Lax pair (a zero-curvature representation) parametrized by points of the following algebraic curve

$$(2) \quad \lambda_i^2 - \lambda_j^2 = r_j - r_i; \quad i, j = 1; \dots; n;$$

in the space C^n with coordinates $\lambda_1; \dots; \lambda_n$. According to [8], this curve is of genus $1 + (n-3)2^{n-2}$.

System (1) has also an infinite number of symmetries, conservation laws [8], and a Backlund auto-transformation with a parameter [1]. Soliton-like solutions of (1) can be found in [1]. According to [8], for $n = 3$ this system coincides with the higher symmetry (the commuting flow) of third order for the well-known Landau-Lifshitz equation (see, e.g., [10]). Thus system (1) is an n -dimensional generalization of the Landau-Lifshitz equation.

Note that, to our knowledge, before the present paper the Wahlquist-Estabrook method was never applied to a system of PDEs connected with an algebraic curve of genus greater than 1, and the explicit structure of the WE algebra was not computed for any system with more than two dependent variables (unknown functions $s^i(x;t)$).

For arbitrary n we construct an epimorphism from the WE algebra of (1) onto the finite-dimensional Lie algebra $L(n)$ of certain $so_{n,1}$ -valued functions on the curve (2). Note that $L(n)$ is not graded, but is quasigraded [16]

$$L(n) = \bigoplus_{i=1}^M L_i; \quad [L_i; L_j] = L_{i+j} + L_{i+j-2};$$

$$\dim L_{2k-1} = n; \quad \dim L_{2k} = \frac{n(n-1)}{2} \quad 8 \leq k \leq N;$$

For $n = 3; 4; 5$ we prove that the WE algebra is isomorphic to the direct sum of $L(n)$ and a two-dimensional abelian Lie algebra. In particular, for $n = 3$ the WE algebra of (1) is isomorphic to the WE algebra of the anisotropic Landau-Lifshitz equation [19]. For $n = 2$ the curve (2) is rational and system (1) belongs to the well-studied class of scalar evolutionary equations [5, 14], so we skip the case $n = 2$.

To achieve this, we prove that the algebra $L(n)$ is isomorphic for any $n \geq 3$ to the Lie algebra given by generators $p_1; \dots; p_n$ and relations

$$(3) \quad [p_i; [p_i; p_k]] - [p_j; [p_j; p_k]] = (r_j - r_i)p_k \quad \text{for } i \notin k; j \notin k;$$

$$[p_i; [p_j; p_k]] = 0 \quad \text{for } i \notin k; i \notin j; j \notin k;$$

In our opinion, this algebraic result may be of independent interest. For $n = 3$ it was proved in [19].

It is known that for a given evolutionary system

$$(4) \quad u_t^i = F^i(u^j; u_x^k; u_{xx}^l; \dots); \quad i = 1; \dots; m;$$

the WE algebra helps also to find a modified system

$$(5) \quad v_t^i = H^i(v^j; v_x^k; v_{xx}^l; \dots); \quad i = 1; \dots; m;$$

connected with (4) by a Backlund transformation of Miura type (sometimes also called a differential substitution)

$$(6) \quad u^i = g^i(v^j; v_x^k; v_{xx}^l; \dots); \quad i = 1; \dots; m;$$

That is, for any solution $v^1; \dots; v^m$ of (5) functions (6) form a solution of (4), and for any solution $u^1; \dots; u^m$ of (4) locally there exist functions $v^1; \dots; v^m$ satisfying (5), (6).

For system (1) a modified system (5) and a transformation (6) were not known. Using a suitable vector field representation of the WE algebra, we find a family of systems (5) and transformations (6) for (1). This family is parametrized by points of the curve (2). Apparently, this is the first example of a family of Miura type transformations parametrized by points of a curve of genus greater than 1.

We would like to make also the following observation. Clearly, relations (3) look somewhat similar to equations (2). And indeed, formula (19) and Theorem 3 below explain how the generator p_i is related to ψ_i . Note that, as we show in Sections 3.1, 3.2, 3.3, at least for $n = 3; 4; 5$ relations (3) arise from the internal jet space geometry of system (1) using the Wahlquist-Estabrook procedure. Therefore, the Wahlquist-Estabrook method and its generalization in [11, 12] may be useful also for the following problem: given a system of PDEs, which is suspected to be integrable, how to recover an algebraic curve that most naturally parametrizes a possible Lax pair for this system? For example, applying the Wahlquist-Estabrook procedure to system (1) without any knowledge of Lax pairs or algebraic curves behind (1), one obtains relations (3) for the WE algebra. Since p_i should correspond to some matrix-valued functions on an algebraic curve, looking at (3) it is not so hard to guess that the curve should be of the form (2). It would be very interesting to make this observation into a rigorous construction for more PDEs.

Several more integrable systems associated with the curve (2) were introduced in [8, 22, 23, 24], where the opposite approach is taken: they start with an infinite-dimensional Lie algebra very similar to our algebra $L(n)$ and construct integrable systems from this Lie algebra. Note that a representation of the Lie algebra in terms of a finite number of generators and relations was not obtained in [8, 22, 23, 24].

The functions $S = (s^1(x;t); \dots; s^n(x;t))$ in (1) and the parameters $\psi_i; r_i$ in (2) may take values in $\mathbb{C}$ or $\mathbb{R}$. In this paper the $\mathbb{C}$ -valued case is studied, but all results and proofs are valid also in the $\mathbb{R}$ -valued case, if one replaces $\mathbb{C}$ by $\mathbb{R}$ in the definitions.

The paper is organized as follows.

In Section 2 we present a rigorous definition of WE algebras for arbitrary evolutionary systems and, therefore, for any systems of PDEs that can be written in evolutionary form. For this we use formal power series with coefficients in Lie algebras. For example, such series occur in computations of the WE algebra of the f-Gordon equation $u_{tt} = u_{xx} = f(u)$ [21], which, as is well known, can be rewritten in evolutionary form as follows

$$(7) \quad \begin{aligned} u_t &= q; \\ q_t &= u_{xx} + f(u); \end{aligned}$$

We discuss also possible generalizations of the Wahlquist-Estabrook ansatz.

In Sections 3, 4 the above mentioned results on the WE algebra of system (1) are obtained, and in Section 5 the family of Miura type transformations is constructed. The appendix contains the proof of technical Lemma 1.

The following abbreviations are used in the paper: WE = Wahlquist-Estabrook, ZCR = zero-curvature representation.

2. The general definition of WE algebras

Originally the Wahlquist-Estabrook prolongation method was formulated in terms of differential forms. We prefer the vector fields version of it, which goes as follows. For a given m -component evolutionary system of PDEs of order $d \geq 1$

$$(8) \quad u_t^i = F^i(u^1; \dots; u^m; u_1^1; \dots; u_1^m; \dots; u_d^1; \dots; u_d^m); \quad i = 1; \dots; m; \quad u_k^i = \frac{\partial^k u^i}{\partial x^k};$$

consider the infinite-dimensional jet space with the coordinates

$$(9) \quad x, t; u_k^i; \quad i = 1, \dots, m; \quad k = 0, 1, 2, \dots; \quad u_0^i = u^i.$$

The total derivative operators

$$D_x = \frac{\partial}{\partial x} + \sum_{i,k} u_{k+1}^i \frac{\partial}{\partial u_k^i}; \quad D_t = \frac{\partial}{\partial t} + \sum_{i,k} D_x^{k+1}(F^i) \frac{\partial}{\partial u_k^i}$$

are commuting vector fields on this space.

Let be an $(m \times d)$ -matrix with entries

$$z_{ik} \in \mathbb{Z}_+; \quad i = 1, \dots, m; \quad k = 0, \dots, d-1.$$

Denote the set of such matrices by M . Denote by U the following product of coordinates (9)

$$U = \prod_{\substack{i=1, \dots, m; \\ k=0, \dots, d-1}} u_k^i z_{ik}.$$

Consider two formal power series

$$(10) \quad X = \sum_{2M} A U; \quad T = \sum_{2M} B U;$$

in the variables

$$(11) \quad u_k^i; \quad 1 \leq i \leq m; \quad 0 \leq k \leq d-1;$$

where the coefficients $A; B$ are elements of some (not specified yet) Lie algebra. The equation

$$(12) \quad [D_x + X; D_t + T] = D_x(T) - D_t(X) + [X; T] = 0$$

in the space of formal power series is equivalent to some Lie algebraic relations for the elements $A; B$. Here

$$D_x(T) = \sum_{2M} B D_x(U); \quad D_t(X) = \sum_{2M} A D_t(U);$$

$$[X; T] = \sum_{2M} [A; B] U U;$$

Let F be the free Lie algebra generated by all the letters $A; B$ for $; 2M$. The quotient of F over the above mentioned relations arising from equation (12) is called the Wahlquist-Estabrook Lie algebra of system (8) (or the WE algebra in brief). From now on $A; B$ are elements of the WE algebra. Then (10) is the most general solution of (12) provided that $X; T$ are power series in variables (11).

Remark 1. From any system (8) equation (12) implies that $X; T$ are of the form

$$(13) \quad X = \sum_{i=1}^{K_1} C_i f_i; \quad T = \sum_{j=1}^{K_2} D_j g_j; \quad k_1, k_2 \in \mathbb{N};$$

where $f_i; g_j$ are analytic functions of variables (11), the functions $f_1, \dots, f_{k_1}$ are linearly independent, the functions $g_1, \dots, g_{k_2}$ are linearly independent, and $C_i; D_j$ are elements of the WE algebra. Expanding $f_i; g_j$ as power series in (11), we obtain that $A; B$ from (10) are linear combinations of $C_i; D_j$ and, therefore, in this case $C_i; D_j$ can be taken as another set of generators of the same WE algebra.

However, the cases when $X; T$ are formal power series and cannot be presented as finite sums of analytic functions do also occur. For example, this happens for the f-Gordon equation $u_{tt} - u_{xx} = f(u)$ [21], which can be rewritten in evolutionary form (7).

Remark 2. A natural question arises, what happens if in equation (12) one considers power series X, T in the variables

$$u_k^i; \quad 1 \leq i \leq m; \quad 0 \leq k \leq s;$$

for arbitrary s ? It turns out that if $s > d - 1$ then before solving (12) one should simplify X, T by so-called gauge transformations [11, 12], and then one also obtains certain Lie algebras [11, 12], which are generally bigger than the WE algebra for $s = d - 1$. For example, for the Krichever-Novikov equation the WE algebra for $s = d - 1$ is trivial, but the case $s = d$ does produce an interesting Lie algebra [14].

Coordinate-free meaning of the Lie algebras for arbitrary s is studied in [11, 12]. In this way one obtains new geometric invariants of PDEs, which allow to prove, for example, that some PDEs are not connected by any Backlund transformations [11, 12].

Let g be a Lie algebra. Recall [10] that g -valued functions $M; N$ of a finite number of variables (9) form a zero-curvature representation (ZCR in short) for system (8) if

$$(14) \quad [D_x + M; D_t + N] = D_x(N) - D_t(M) + [M; N] = 0;$$

If $\dim g < 1$ then $M; N$ are supposed to be analytic or smooth g -valued functions, while if $\dim g = 1$ then $M; N$ are formal power series with coefficients in g . For example, the power series $X; T$ form a ZCR with values in the WE algebra.

It is known that every Lax pair for a $(1+1)$ -dimensional system of PDEs determines a ZCR. In addition to the Wahlquist-Estabrook technique, some other methods to obtain ZCRs for a given system of PDEs also exist (see, e.g., [17, 18, 20] and references therein).

Since (10) is the most general solution of (12) and equation (14) is similar to (12), we obtain the following result.

Proposition 1. Suppose that $M; N$ are g -valued functions of variables (11) and form a ZCR. Expand $M; N$ as power series in (11)

$$M = \sum_{2M}^X M^U; \quad N = \sum_{2M}^X N^U; \quad M; N \in 2g;$$

Then the map $A \mapsto M; B \mapsto N$ determines a homomorphism from the WE algebra to g . If the coefficients $M; N; \dots \in 2M$, generate the whole Lie algebra g then this homomorphism is surjective.

3. Computations for the generalized Landau-Lifshitz system

In order to study the WE algebra of system (1), we need to resolve (locally) the constraint $hS; S = 1$ for the vector $S = (s^1(x;t); \dots; s^n(x;t))$. Following [8], we do it as

$$(15) \quad s^i = \frac{2u^i}{1 + h u; u^i}; \quad i = 1; \dots; n-1; \quad s^n = \frac{1 - h u; u^i}{1 + h u; u^i};$$

where u is an $(n-1)$ -dimensional vector with the components $u^1(x;t); \dots; u^{n-1}(x;t)$, and $h; \cdot$ is the standard scalar product. Then one can rewrite system (1) (as [8])

$$(16) \quad u_t = u_{xxx} - 6h u; u_x i^{-1} u_{xx} + 6h u; u_{xx} i^{-1} + 24h u; u_x i^2 - 2 \sum_{i=1}^{n-1} 6h u; u_i h u_x; u_x i^{-2} u_x + 6h u_x; u_{xx} i^{-1} - 12h u; u_x i h u_x; u_x i^{-2} u + \frac{3}{2} r_n + 4 \sum_{i=1}^{n-1} (r_i - r_n) (u^i)^2 u_x;$$

where $\epsilon = 1 + h u; u^i$ and $r_1; \dots; r_n$ are distinct complex numbers that are the entries of the matrix $R = \text{diag}(r_1; \dots; r_n)$ from system (1).

Let $E_{ij} \in \mathfrak{gl}_{n+1}(\mathbb{C})$ be the matrix with (i,j) -th entry equal to 1 and all other entries equal to zero. Recall that the Lie subalgebra $\mathfrak{so}_{n,1} \subset \mathfrak{gl}_{n+1}(\mathbb{C})$ has the following basis

$$E_{ij}, E_{ji}; \quad i < j \leq n; \quad E_{1,n+1} + E_{n+1,1}; \quad l = 1, \dots, n;$$

From the results of [8, 25] one can obtain the following $\mathfrak{so}_{n,1}$ -valued ZCR of system (1)

$$(17) \quad M = \sum_{i=1}^n s^i (E_{i,n+1} + E_{n+1,i});$$

$$(18) \quad N = D_x^2(M) + [D_x(M), M] + (r_1 + \frac{1}{2})M + \frac{1}{2}(S;RS) + \frac{3}{2}(S_x;S_x)M;$$

Here $s_1, \dots, s_n$ are complex parameters satisfying equations (2).

Remark 3. It was noticed in [22] that the formulas $x_i = \frac{1}{2} + r_i$; $y = \prod_{i=1}^n x_i$ provide a mapping from the curve (2) to the hyperelliptic curve $y^2 = \prod_{i=1}^n (x - r_i)$. However, according to [8], the curve (2) itself is not hyperelliptic.

If $S = (s^1, \dots, s^n)$ is given by formulas (15) then (17), (18) determines a ZCR for system (16).

In the algebra $\mathbb{C}[s_1, \dots, s_n]$ consider the ideal $I \subset \mathbb{C}[s_1, \dots, s_n]$ generated by the polynomials $\frac{1}{2}x_i^2 + r_i x_j$; $i, j = 1, \dots, n$. Denote by $\bar{\cdot}$ the image of $\cdot$ in the quotient algebra $\mathbb{Q} = \mathbb{C}[s_1, \dots, s_n]/I$, which is equal to the algebra of polynomial functions on the curve (2).

Consider the infinite-dimensional Lie algebra over $\mathbb{C}$

$$\mathfrak{gl}_{n+1}(\mathbb{C}) \otimes_{\mathbb{C}} \mathbb{Q} = \mathfrak{gl}_{n+1}(\mathbb{Q});$$

$$[g_1, g_2] = [g_1, g_2]; \quad g_i \in \mathfrak{gl}_{n+1}(\mathbb{C}); \quad g_i \in \mathbb{Q};$$

and the elements

$$(19) \quad Q_i = (E_{i,n+1} + E_{n+1,i}) \in \mathfrak{so}_{n,1} \otimes_{\mathbb{C}} \mathbb{Q} \subset \mathfrak{gl}_{n+1}(\mathbb{C}) \otimes_{\mathbb{C}} \mathbb{Q}; \quad i = 1, \dots, n;$$

Denote by $L(n) \subset \mathfrak{so}_{n,1} \otimes_{\mathbb{C}} \mathbb{Q}$ the Lie subalgebra generated by $Q_1, \dots, Q_n$.

Obviously, the element $\frac{1}{2}x_i^2 + r_i x_j \in \mathbb{Q}$ does not depend on i . For $i, j = 1, \dots, n$, $i \neq j$, and $k \in \mathbb{N}$ consider the following elements of $\mathfrak{so}_{n,1} \otimes_{\mathbb{C}} \mathbb{Q}$

$$Q_i^{2k-1} = (E_{i,n+1} + E_{n+1,i})^{k-1}; \quad Q_{ij}^{2k} = (E_{ij} - E_{ji})^{k-1};$$

For $i, j, a, b = 1, \dots, n$, $i \neq j$, $a \neq b$, and $k_1, k_2 \in \mathbb{N}$ one has

$$(20) \quad [Q_{ij}^{2k_1}; Q_{ab}^{2k_2}] = a_j Q_{ib}^{2(k_1+k_2)} - i_b Q_{aj}^{2(k_1+k_2)} + j_b Q_{ai}^{2(k_1+k_2)} - i_a Q_{jb}^{2(k_1+k_2)} + r_i i_b Q_{aj}^{2(k_1+k_2-1)} - r_j a_j Q_{ib}^{2(k_1+k_2-1)} + r_i i_a Q_{jb}^{2(k_1+k_2-1)} - r_j j_b Q_{ai}^{2(k_1+k_2-1)};$$

$$(21) \quad [Q_{ij}^{2k_1}; Q_a^{2k_2-1}] = a_j Q_i^{2k_1+2k_2-1} - i_a Q_j^{2k_1+2k_2-1} - r_j a_j Q_i^{2k_1+2k_2-3} + r_i i_a Q_j^{2k_1+2k_2-3};$$

$$(22) \quad [Q_i^{2k_1-1}; Q_j^{2k_2-1}] = Q_j^{2(k_1+k_2-1)}; \quad [Q_i^{2k_1-1}; Q_i^{2k_2-1}] = 0;$$

Since $Q_i^1 = Q_i$ and $Q_{ij}^{2k} = Q_{ji}^{2k}$, from (20), (21), (22) we obtain that

$$Q_i^{2k-1}; \quad Q_{ij}^{2k}; \quad i, j = 1, \dots, n; \quad i < j; \quad k = 1, 2, 3, \dots;$$

span the Lie algebra $L(n)$. It is easily seen that these elements are linearly independent over $\mathbb{C}$ and, therefore, form a basis of $L(n)$.

For $k \in \mathbb{N}$ set

$$L_{2k-1} = \mathbb{H} Q_i^{2k-1}; \quad i = 1, \dots, n; \quad L_{2k} = \mathbb{H} Q_{ij}^{2k}; \quad i, j = 1, \dots, n; \quad i < j;$$

Here and below for elements $v_1, \dots, v_s$ of a vector space the expression $\langle v_1, \dots, v_s \rangle$ denotes the linear span of $v_1, \dots, v_s$ over $\mathbb{C}$. Then from (20), (21), (22) one gets

$$L(n) = \sum_{i=1}^M L_i; \quad [L_i, L_j] = L_{i+j} + L_{i+j-2}.$$

Therefore, the Lie algebra $L(n)$ is quasigraded [16].

Clearly, formulas (17), (18) can be regarded as a ZCR with values in the algebra $L(n)$. In particular, $M = \sum_{i=1}^n s^i Q_i$. Combining this with Proposition 1, we obtain the following.

Theorem 1. For any $n \geq 3$ we have an epimorphism from the WE algebra of system (1) onto the infinite-dimensional Lie algebra $L(n)$.

Let us give a complete description of the WE algebra of (16) for small n . For $n = 2$ system (16) is a scalar equation of the form $u_t = u_{xxx} + f(u; u_x; u_{xx})$. Since for such equations the WE algebras have already been studied quite extensively (see, e.g., [5, 14] and references therein) and the curve (2) is rational for $n = 2$, we skip the case $n = 2$. For $n = 3, 4, 5$ the WE algebras are studied below.

3.1. The case $n = 3$. According to Section 2, we must solve equation (12) for

$$X = X(u^1; u^2; u_x^1; u_x^2; u_{xx}^1; u_{xx}^2); \quad T = T(u^1; u^2; u_x^1; u_x^2; u_{xx}^1; u_{xx}^2):$$

If we differentiate (12) with respect to the variables $u_{xxxx}^1, u_{xxxx}^2, u_{xxxx}^1, u_{xxxx}^2$, we get that X depends only on $(u^1; u^2)$. Next, differentiating (12) with respect to $u_{xxx}^1, u_{xxx}^2, u_{xxx}^1, u_{xxx}^2$ several times, one obtains

$$X = X(u^1; u^2); \quad T = \frac{\partial X}{\partial u^1} u_{xx}^1 + \frac{\partial X}{\partial u^2} u_{xx}^2 + F_1(u^1; u^2; u_x^1; u_x^2);$$

where

$$F_1 = \frac{1}{2} \frac{\partial^2 X}{\partial u^1 \partial u^1} + 6u^1 \frac{\partial X}{\partial u^1} - 6u^2 \frac{\partial X}{\partial u^2} - (u_x^1)^2 \\ - \frac{\partial^2 X}{\partial u^1 \partial u^2} + 6u^2 \frac{\partial X}{\partial u^1} + 6u^1 \frac{\partial X}{\partial u^2} - u_x^1 u_x^2 \\ - \frac{1}{2} \frac{\partial^2 X}{\partial u^2 \partial u^2} + 6u^2 \frac{\partial X}{\partial u^2} - 6u^1 \frac{\partial X}{\partial u^1} - (u_x^2)^2 + \left[\frac{\partial X}{\partial u^1}; X \right] u_x^1 + \left[\frac{\partial X}{\partial u^2}; X \right] u_x^2 + F_2(u^1; u^2);$$

and $\Delta = 1 + (u^1)^2 + (u^2)^2$.

Then the expression

$$(23) \quad [D_x + X; D_t + T] = D_x(T) - D_t(X) + [X; T]$$

becomes a third degree polynomial in $u_x^1; u_x^2$ with coefficients depending on $u^1; u^2$. By putting equal to zero the third degree coefficients, we get a system of linear PDEs of order 3 for the function $X(u^1; u^2)$. The general solution of this system is

$$(24) \quad X = D_0 + u^1 D_1 + u^2 D_2 + (u^1)^2 D_3 + (u^2)^2 D_4 + 1; \quad \Delta$$

where D_i do not depend on $u^1; u^2$ and, therefore, belong to a set of generators of the WE algebra.

Taking into account, that system (16) arises from (1) by means of (15), it is more convenient to rewrite (24) as

$$X = 2C_1 u^1 + 2C_2 u^2 + C_3 \Delta - (u^1)^2 - (u^2)^2 - 1 + C_0;$$

where C_i are also elements of the W E algebra. By putting to zero the coefficients of (23) at the monomials $(u_x^1)^2$ and $(u_x^2)^2$, we obtain

$$(25) \quad [C_0; C_1] = [C_0; C_2] = [C_0; C_3] = 0;$$

Now, by putting the coefficients of u_x^1 and u_x^2 equal to zero, one gets a system of two first order differential equations for $F_2(u^1; u^2)$. Its compatibility conditions are satisfied if and only if the following relations hold

$$(26) \quad [C_0; [C_1; C_2]] = [C_0; [C_1; C_3]] = [C_0; [C_2; C_3]] = 0;$$

$$(27) \quad [C_1; [C_2; C_3]] = [C_2; [C_3; C_1]] = [C_3; [C_1; C_2]] = 0;$$

$$(28) \quad [C_1; [C_1; C_3]] - [C_2; [C_2; C_3]] = (r_2 - r_1)C_3;$$

$$(29) \quad [C_1; [C_1; C_2]] - [C_3; [C_3; C_2]] = (r_3 - r_1)C_2;$$

$$(30) \quad [C_2; [C_2; C_1]] - [C_3; [C_3; C_1]] = (r_3 - r_2)C_1;$$

Note that (26) follows from (25) using the Jacobi identity.

Then we are able to compute $F_2(u^1; u^2)$.

$$\begin{aligned} F_2 = & (2[C_3; [C_1; C_3]] + 3C_1r_3)(u^1)^5 + (3C_2r_3 - 2[C_3; [C_2; C_3]])(u^1)^4u^2 \\ & + (6C_1r_3 - 4[C_3; [C_1; C_3]])(u^1)^3(u^2)^2 + (4[C_3; [C_2; C_3]] + 6C_2r_3)(u^1)^2(u^2)^3 \\ & + (2[C_3; [C_1; C_3]] + 3C_1r_3)u^1(u^2)^4 + (3C_2r_3 - 2[C_3; [C_2; C_3]])(u^2)^5 \\ & + (2C_3r_2 + 2[C_2; [C_2; C_3]] - 2C_3r_1 + 3C_3r_3)(u^1)^4 \\ & + (4[C_2; [C_2; C_3]] + 2C_3r_2 - 2C_3r_1 + 6C_3r_3)(u^1)^2(u^2)^2 \\ & + (2[C_2; [C_2; C_3]] + 3C_3r_3)(u^2)^4 + (4C_1r_1 - 4[C_3; [C_1; C_3]] + 2C_1r_3)(u^1)^3 \\ & + (2C_2r_3 - 4[C_3; [C_2; C_3]] + 4C_2r_1)(u^1)^2u^2 + (4[C_3; [C_1; C_3]] + 4C_1r_2 + 2C_1r_3)u^1(u^2)^2 \\ & + (4C_2r_2 + 2C_2r_3 - 4[C_3; [C_2; C_3]])(u^2)^3 + (4[C_2; [C_2; C_3]] + 4C_3r_2 + 2C_3r_1)(u^1)^2 \\ & + (4[C_2; [C_2; C_3]] + 6C_3r_2)(u^2)^2 + (2[C_3; [C_1; C_3]] + 3C_1r_3)u^1 \\ & + (3C_2r_3 - 2[C_3; [C_2; C_3]])u^2 + C_3r_3 + 2C_3r_2 + 2[C_2; [C_2; C_3]]^3 + C^0; \end{aligned}$$

where C^0 is another generator of the W E algebra.

Then (23) depends only on $u^1; u^2$ and vanishes if and only if

$$(31) \quad [C_3; C^0] = 0;$$

$$(32) \quad [C_0; C^0] = 0;$$

$$(33) \quad [C_3; [C_2; [C_2; C_3]]] = 0;$$

$$(34) \quad [C_3; [C_3; [C_3; C_1]]] = \frac{3}{2}r_3[C_1; C_3] + [C_1; C^0];$$

$$(35) \quad [C_3; [C_3; [C_3; C_2]]] = \frac{3}{2}r_3[C_2; C_3] + [C_2; C^0];$$

$$(36) \quad [[C_2; C_3]; [C_1; C_2]] = \frac{1}{2}r_3[C_1; C_3] + [C_1; C^0] + r_2[C_1; C_3];$$

$$(37) \quad [C_2; [C_2; [C_2; C_3]]] = \frac{1}{2}r_3[C_2; C_3] - [C_2; C^0] - r_2[C_2; C_3];$$

Therefore, the W E algebra of (16) for $n = 3$ is given by the generators C^0, C_0, C_1, C_2, C_3 and all relations obtained in this subsection. Let us simplify the structure of these relations. Taking into account (25) and (27), relations (31), (32), (33), (36), (37) imply

$$(38) \quad [C^0 + r_2 + \frac{1}{2}r_3 C_3 + [C_2; [C_2; C_3]]; C_i] = 0; \quad i = 0; 1; 2; 3;$$

Then it is easily seen that all relations follow from (25), (27), (28), (29), (30), (38) using the Jacobi identity. For example, let us prove this for (33) and (34), for all other relations the statement can be proved analogously. Applying $\text{ad} C_3$ to (28), $\text{ad} C_2$ to (29), and $\text{ad} C_1$ to (30), one obtains

$$[C_i; [C_j; [C_j; C_i]]] = [C_i; [C_k; [C_k; C_i]]] \quad \text{for } f_i; j; k; g = f_1; 2; 3; g:$$

Besides, by the Jacobi identity, we have $[C_i; [C_j; [C_j; C_i]]] = [C_j; [C_i; [C_i; C_j]]]$. Therefore,

$$\begin{aligned} [C_3; [C_2; [C_2; C_3]]] &= [C_2; [C_3; [C_3; C_2]]] = [C_2; [C_1; [C_1; C_2]]] = \\ [C_1; [C_2; [C_2; C_1]]] &= [C_1; [C_3; [C_3; C_1]]] = [C_3; [C_1; [C_1; C_3]]] = [C_3; [C_2; [C_2; C_3]]]; \end{aligned}$$

which implies (33).

Using (30), (27), and the Jacobi identity, one obtains

$$\begin{aligned} [C_3; [C_3; [C_3; C_1]]] &= [C_3; [C_2; [C_2; C_1]]] + (r_2 - r_3) [C_3; C_1] = \\ [C_3; [C_2; [C_2; C_1]]] &+ (r_2 - r_3) [C_3; C_1] = [C_1; [C_2; [C_2; C_3]]] + (r_2 - r_3) [C_3; C_1]; \end{aligned}$$

Combining this with (38) for $i = 1$, we get (34).

Thus the WE algebra is isomorphic to the direct sum $\mathfrak{g}(3) \oplus A$, where $\mathfrak{g}(3)$ is the Lie algebra given by the generators $C_1; C_2; C_3$ and relations (27), (28), (29), (30), and A is the two-dimensional abelian Lie algebra spanned by $C_0, C^0 + r_2 + \frac{1}{2}r_3 C_3 + [C_2; [C_2; C_3]]$.

3.2. The case $n = 4$. According to Section 2, we must solve equation (12) for

$$X = X(u^1; u^2; u^3; u_x^1; u_x^2; u_x^3; u_{xx}^1; u_{xx}^2; u_{xx}^3); \quad T = T(u^1; u^2; u^3; u_x^1; u_x^2; u_x^3; u_{xx}^1; u_{xx}^2; u_{xx}^3):$$

Similarly to the previous subsection we obtain the following.

$$\begin{aligned} X &= 2C_1 u^1 + 2C_2 u^2 + 2C_3 u^3 + C_4 (1 - (u^1)^2 - (u^2)^2 - (u^3)^2) + C_0; \\ T &= \sum_{i=1}^3 X^3 \frac{\partial X}{\partial u^i} u_{xx}^i - \frac{1}{2} \sum_{i=1}^3 X^3 \frac{\partial^2 X}{\partial u^i \partial u^i} + 6 \sum_{j \neq i} \frac{\partial X}{\partial u^i} u^j \frac{\partial X}{\partial u^j} u^i (u_x^i)^2 \\ &\quad + \sum_{1 \leq i < j \leq 3} X \frac{\partial^2 X}{\partial u^i \partial u^j} + 6 \sum_{i=1}^3 \frac{\partial X}{\partial u^i} u^j + \frac{\partial X}{\partial u^j} u^i - \sum_{i=1}^3 u_x^i u_x^j + \sum_{i=1}^3 \left[\frac{\partial X}{\partial u^i}; X \right] u_x^i + F_2(u^1; u^2; u^3); \end{aligned}$$

where $\Delta = 1 + (u^1)^2 + (u^2)^2 + (u^3)^2$ and

$$\begin{aligned} (39) \quad F_2 &= (-2[C_4; [C_1; C_4]] + 3C_1 r_4) (u^1)^5 + (3C_2 r_4 - 2[C_4; [C_2; C_4]]) (u^1)^4 u^2 \\ &\quad + (3C_3 r_4 - 2[C_4; [C_3; C_4]]) (u^1)^4 u^3 + (-4[C_4; [C_1; C_4]] + 6C_1 r_4) (u^1)^3 (u^2)^2 \\ &\quad + (-4[C_4; [C_1; C_4]] + 6C_1 r_4) (u^1)^3 (u^3)^2 + (-4[C_4; [C_2; C_4]] + 6C_2 r_4) (u^1)^2 (u^2)^3 \\ &\quad + (6C_3 r_4 - 4[C_4; [C_3; C_4]]) (u^1)^2 (u^2)^2 u^3 + (-4[C_4; [C_2; C_4]] + 6C_2 r_4) (u^1)^2 u^2 (u^3)^2 \\ &\quad + (6C_3 r_4 - 4[C_4; [C_3; C_4]]) (u^1)^2 (u^3)^3 + (-2[C_4; [C_1; C_4]] + 3C_1 r_4) u^1 (u^2)^4 \\ &\quad + (-4[C_4; [C_1; C_4]] + 6C_1 r_4) u^1 (u^2)^2 (u^3)^2 + (-2[C_4; [C_1; C_4]] + 3C_1 r_4) u^1 (u^3)^4 \\ &\quad + (3C_2 r_4 - 2[C_4; [C_2; C_4]]) (u^2)^5 + (3C_3 r_4 - 2[C_4; [C_3; C_4]]) (u^2)^4 u^3 \\ &\quad + (-4[C_4; [C_2; C_4]] + 6C_2 r_4) (u^2)^3 (u^3)^2 + (6C_3 r_4 - 4[C_4; [C_3; C_4]]) (u^2)^2 (u^3)^3 \\ &\quad + (3C_2 r_4 - 2[C_4; [C_2; C_4]]) u^2 (u^3)^4 + (3C_3 r_4 - 2[C_4; [C_3; C_4]]) (u^3)^5 \\ &\quad + (2C_4 r_3 + 2[C_3; [C_3; C_4]] + 3C_4 r_4 - 2C_4 r_1) (u^1)^4 \\ &\quad + (4[C_3; [C_3; C_4]] - 2C_4 r_2 - 2C_4 r_1 + 6C_4 r_4 + 4C_4 r_3) (u^1)^2 (u^2)^2 \end{aligned}$$

$$\begin{aligned}
& + (4[\mathbb{C}_3; [\mathbb{C}_3; \mathbb{C}_4]] + 2C_4r_3 + 6C_4r_4 - 2C_4r_1)(u^1)^2(u^3)^2 \\
& + (2C_4r_2 + 3C_4r_4 + 2[\mathbb{C}_3; [\mathbb{C}_3; \mathbb{C}_4]] + 2C_4r_3)(u^2)^4 \\
& + (2C_4r_2 + 4[\mathbb{C}_3; [\mathbb{C}_3; \mathbb{C}_4]] + 2C_4r_3 + 6C_4r_4)(u^2)^2(u^3)^2 + (3C_4r_4 + 2[\mathbb{C}_3; [\mathbb{C}_3; \mathbb{C}_4]]) (u^3)^4 \\
& + (2C_1r_4 - 4[\mathbb{C}_4; [\mathbb{C}_1; \mathbb{C}_4]] + 4C_1r_1)(u^1)^3 + (4[\mathbb{C}_4; [\mathbb{C}_2; \mathbb{C}_4]] + 2C_2r_4 + 4C_2r_1)(u^1)^2u^2 \\
& + (4[\mathbb{C}_4; [\mathbb{C}_3; \mathbb{C}_4]] + 4C_3r_1 + 2C_3r_4)(u^1)^2u^3 + (4[\mathbb{C}_4; [\mathbb{C}_1; \mathbb{C}_4]] + 2C_1r_4 + 4C_1r_2)u^1(u^2)^2 \\
& + (2C_1r_4 - 4[\mathbb{C}_4; [\mathbb{C}_1; \mathbb{C}_4]] + 4C_1r_3)u^1(u^3)^2 + (4[\mathbb{C}_4; [\mathbb{C}_2; \mathbb{C}_4]] + 4C_2r_2 + 2C_2r_4)(u^2)^3 \\
& + (4C_3r_2 + 2C_3r_4 - 4[\mathbb{C}_4; [\mathbb{C}_3; \mathbb{C}_4]]) (u^2)^2u^3 + (4[\mathbb{C}_4; [\mathbb{C}_2; \mathbb{C}_4]] + 4C_2r_3 + 2C_2r_4)u^2(u^3)^2 \\
& + (4[\mathbb{C}_4; [\mathbb{C}_3; \mathbb{C}_4]] + 4C_3r_3 + 2C_3r_4)(u^3)^3 + (2C_4r_1 + 4C_4r_3 + 4[\mathbb{C}_3; [\mathbb{C}_3; \mathbb{C}_4]]) (u^1)^2 \\
& + (4[\mathbb{C}_3; [\mathbb{C}_3; \mathbb{C}_4]] + 2C_4r_2 + 4C_4r_3)(u^2)^2 + (4[\mathbb{C}_3; [\mathbb{C}_3; \mathbb{C}_4]] + 6C_4r_3)(u^3)^2 \\
& + (2[\mathbb{C}_4; [\mathbb{C}_1; \mathbb{C}_4]] + 3C_1r_4)u^1 + (3C_2r_4 - 2[\mathbb{C}_4; [\mathbb{C}_2; \mathbb{C}_4]])u^2 \\
& + (3C_3r_4 - 2[\mathbb{C}_4; [\mathbb{C}_3; \mathbb{C}_4]])u^3 + 2C_4r_3 + 2[\mathbb{C}_3; [\mathbb{C}_3; \mathbb{C}_4]] + C_4r_4 \quad^3 + C^0:
\end{aligned}$$

Here $C^0; C_0; C_1; C_2; C_3; C_4$ are the generators of the W E algebra.

The W E algebra is equal to the direct sum $\mathfrak{g}(4) \oplus A$, where $\mathfrak{g}(4)$ is the Lie algebra given by the generators $C_1; C_2; C_3; C_4$ and the relations

$$(40) \quad [\mathbb{C}_i; [\mathbb{C}_i; \mathbb{C}_k]] - [\mathbb{C}_j; [\mathbb{C}_j; \mathbb{C}_k]] = (r_j - r_i)C_k \quad \text{for } i \notin k; j \notin k;$$

$$(41) \quad [\mathbb{C}_i; [\mathbb{C}_j; \mathbb{C}_k]] = 0 \quad \text{for } i \notin k; i \notin j; j \notin k;$$

and A is the two-dimensional abelian Lie algebra spanned by

$$C_0; \quad C^0 + r_3 + \frac{1}{2}r_4 C_4 + [\mathbb{C}_3; [\mathbb{C}_3; \mathbb{C}_4]]:$$

In particular, one has

$$[\mathbb{C}_i; C_0] = [\mathbb{C}_i; C^0 + r_3 + \frac{1}{2}r_4 C_4 + [\mathbb{C}_3; [\mathbb{C}_3; \mathbb{C}_4]]] = 0 \quad \text{for } i = 0; 1; 2; 3; 4:$$

3.3. The case $n = 5$. Similarly to the previous subsections one gets the following results.

$$X = 2C_1u^1 + 2C_2u^2 + 2C_3u^3 + 2C_4u^4 + C_5 - 1 - (u^1)^2 - (u^2)^2 - (u^3)^2 - (u^4)^2 - 1 + C_0:$$

$$\begin{aligned}
T = & \sum_{i=1}^4 \frac{\partial^4 X}{\partial u^i \partial x^x} u_x^i - \frac{1}{2} \sum_{i=1}^4 \frac{\partial^2 X}{\partial u^i \partial u^i} + 6 \sum_{j \neq i} \frac{\partial X}{\partial u^i} u^j \sum_{j \neq i} \frac{\partial X}{\partial u^j} u^j - 1 (u_x^i)^2 \\
& + \sum_{1 \leq i < j \leq 4} \frac{\partial^2 X}{\partial u^i \partial u^j} + 6 \sum_{i=1}^4 \frac{\partial X}{\partial u^i} u^j + \frac{\partial X}{\partial u^j} u^i - 1 u_x^i u_x^j + \sum_{i=1}^4 \frac{\partial^4 X}{\partial u^i} [X] u_x^i + F_2(u^1; u^2; u^3; u^4):
\end{aligned}$$

Here $= 1 + (u^1)^2 + (u^2)^2 + (u^3)^2 + (u^4)^2$ and

$$F_2 = P(u^1; u^2; u^3; u^4) - 3 + C^0;$$

where $P(u^1; u^2; u^3; u^4)$ is a cumbersome polynomial of degree 5 similar to (39) and $C^0, C_0, C_1, C_2, C_3, C_4, C_5$ are the generators of the W E algebra.

The W E algebra is equal to the direct sum $\mathfrak{g}(5) \oplus A$, where $\mathfrak{g}(5)$ is the Lie algebra given by the generators $C_1; C_2; C_3; C_4; C_5$ and relations (40), (41) for $i; j; k = 1; \dots; 5$, and A is the two-dimensional abelian Lie algebra spanned by

$$C_0; \quad C^0 + r_4 + \frac{1}{2}r_5 C_5 + [\mathbb{C}_4; [\mathbb{C}_4; \mathbb{C}_5]]:$$

In particular, we have

$$[C_i; C_0] = [C_i; C^0 + r_4 + \frac{1}{2}r_5 C_5 + [C_4; [C_4; C_5]]] = 0 \quad \text{for } i = 0; 1; \dots; 5:$$

The explicit structure of the obtained algebras is described in the next section.

4. Explicit structure of the algebras

Let F be the free Lie algebra generated by the letters $p_1; \dots; p_n$, $n \geq 3$. Denote by $g(n)$ the quotient of F over the relations

$$(42) \quad [p_i; [p_i; p_k]] - [p_j; [p_j; p_k]] = (r_j - r_i)p_k \quad \text{for } i \notin k; j \notin k;$$

$$(43) \quad [p_i; [p_j; p_k]] = 0 \quad \text{for } i \notin k; i \notin j; j \notin k:$$

The natural image of $p_i \in F$ in the quotient Lie algebra $g(n)$ is denoted by the same symbol p_i .

The results of Sections 3.1, 3.2, 3.3 can be summarized as follows.

Theorem 2. For $n = 3; 4; 5$ the W E algebra of system (16) is isomorphic to the direct sum of $g(n)$ and a two-dimensional abelian Lie algebra.

The next theorem describes the explicit structure of $g(n)$.

Theorem 3. For any $n \geq 3$ the mapping

$$(44) \quad \varphi : g(n) \rightarrow L(n); \quad p_i \mapsto Q_i;$$

is an isomorphism of Lie algebras. Here $L(n) = \mathfrak{so}_{n+1}$, Q is the infinite-dimensional Lie algebra defined in Section 3.

Proof. For $n = 3$ this theorem was proved in [19] for a different matrix representation of $L(n)$, and we will use a similar method.

It is easy to check that Q_i satisfy relations (42), (43), so mapping (44) is an epimorphism.

Define a filtration on $L(n)$ by vector subspaces L^m as follows

$$L^0 = 0; \quad L^1 = \langle Q_1; \dots; Q_n \rangle; \quad L^m = L^1 + \sum_{i+j=m} [L^i; L^j] \quad \text{for } m > 1;$$

$$L^0 \subset L^1 \subset L^2 \subset \dots \subset L(n) = L^i:$$

From (20), (21), (22) one gets that for all $k \in \mathbb{N}$

$$Q_1^{2s-1}; Q_{ij}^{2s}; \quad i, j, l = 1; \dots; n; \quad i < j; \quad s \leq k; \quad \text{form a basis of } L^{2k};$$

$$Q_1^{2s-1}; Q_{ij}^{2s-2}; \quad i, j, l = 1; \dots; n; \quad i < j; \quad s \leq k; \quad \text{form a basis of } L^{2k-1}:$$

This implies

$$(45) \quad \dim(L^m = L^{m-1}) = \begin{cases} n; & \text{if } m \text{ is odd;} \\ n(n-1)/2; & \text{if } m \text{ is even;} \end{cases}$$

Consider the similar filtration on $g(n)$ by vector subspaces g^m

$$g^0 = 0; \quad g^1 = \langle p_1; \dots; p_n \rangle; \quad g^m = g^1 + \sum_{i+j=m} [g^i; g^j] \quad \text{for } m > 1:$$

Clearly,

$$(46) \quad \varphi(g^m) = L^m:$$

Combining this with (45), we obtain that it remains to prove

$$(47) \quad \dim(\mathfrak{g}^m = \mathfrak{g}^{m-1}) \begin{cases} n; & \text{if } m \text{ is odd;} \\ n(n-1)/2; & \text{if } m \text{ is even;} \end{cases}$$

Indeed, if (47) holds then (45) and (46) imply that ϕ is an isomorphism.

For $n = 3$ statement (47) was proved in [19]. Below we suppose $n \geq 4$.

For each $k \geq 2$ set

$$\begin{aligned} P_{ij}^{2k} &= (\text{ad } p_i)^{2k-1}(p_j); \quad \text{for } i \neq j; \\ P_1^{2k-1} &= (\text{ad } p_2)^{2k-2}(p_1); \\ P_i^{2k-1} &= (\text{ad } p_1)^{2k-2}(p_i) \quad \text{for } i = 2, 3, \dots, n; \end{aligned}$$

We will use the following notation for iterated commutators

$$[a_1 a_2 \dots a_{k-1} a_k] = [a_1; [a_2; [\dots; [a_{k-1}; a_k] \dots]];$$

Also, for brevity we replace each generator p_i by the corresponding index i . So, for example,

$$\begin{aligned} [i[jjk]lk] &= [p_i; [p_j; [p_j; p_k]; [p_l; p_k]]]; \\ P_{ij}^{2k} &= [i; \underbrace{\dots}_{2k-1} j]; \quad P_1^{2k-1} = [1; \underbrace{\dots}_{2k-2} 1]; \\ P_i^{2k-1} &= [1; \underbrace{\dots}_{2k-2} i] \quad \text{for } i = 2, 3, \dots, n; \end{aligned}$$

For such an iterated commutator C of several p_i denote by $o(C)$ the number of p_i that appear in C . For example,

$$\begin{aligned} o([ij]) &= 2; \quad o([iij]) = 3; \quad o([iijjk]lk) = 7; \\ o(P_{ij}^{2k}) &= 2k; \quad o(P_i^{2k-1}) = 2k-1; \quad o(P_{ij}^{2k_1}; P_{i^0 j^0}^{2k_2}) = 2(k_1 + k_2); \end{aligned}$$

For two such iterated commutators $A; B$ we write $A \leq B$ if

$$A \leq B \iff \exists m \text{ such that } A \in \mathfrak{g}^m \text{ for } m = \max(o(A), o(B)) - 1;$$

In particular, $A \leq 0$ means that $A \in \mathfrak{g}^m$ for $m = o(A) - 1$.

Remark 4. Denote by $\bar{C}$ the image of C in the quotient vector space $\mathfrak{g}^{o(C)} = \mathfrak{g}^{o(C)-1}$. Since $[\bar{g}^i; \bar{g}^j] \subset \bar{g}^{i+j}$, we can consider the associated graded Lie algebra

$$\text{gr}(\mathfrak{g}) = \bigoplus_{i=0}^M \mathfrak{g}^i = \mathfrak{g}^{i-1};$$

So, $A \leq B$ if and only if $A = B$ in $\text{gr}(\mathfrak{g})$. In Lemma 1 below we essentially obtain some identities in the algebra $\text{gr}(\mathfrak{g})$.

From Lemma 1 by induction on k one obtains that

$$\begin{aligned} P_1^{2s-1}; P_{ij}^{2s}; \quad i, j, l = 1, \dots, n; \quad i < j; \quad s \leq k; \quad \text{span } \mathfrak{g}^{2k}; \\ P_1^{2s-1}; P_{ij}^{2s-2}; \quad i, j, l = 1, \dots, n; \quad i < j; \quad s \leq k; \quad \text{span } \mathfrak{g}^{2k-1}; \end{aligned}$$

which implies (47). This lemma is proved in the Appendix.

Lemma 1. If i, j, i^0, j^0 are distinct integers from $1, \dots, n$ then for all $k_1, k_2 \in \mathbb{Z}_+$ one has

$$(48) \quad [\underbrace{[i; \dots] j}_{2k_1} \underbrace{[i^0; \dots] j^0}_{2k_2}]' = 0; \quad \text{in particular; } [P_j^{2k_1+1}; P_j^{2k_2+1}]' = 0;$$

$$(49) \quad P_{ij}^{2(k_1+k_2+1)} \neq P_{ji}^{2(k_1+k_2+1)};$$

$$(50) \quad [P_{ij}^{2k_1}; P_{ij}^{2k_2+2}] = 0 \quad \text{for } k_1 = 1;$$

$$(51) \quad [P_i^{2k_1+1}; P_j^{2k_2+1}] = P_{ij}^{2(k_1+k_2+1)};$$

$$(52) \quad [P_i^{2k_1+1}; P_{ij}^{2k_2+2}] = P_j^{2(k_1+k_2)+3};$$

$$(53) \quad [P_i^{2k_1+1}; P_{i^0 j^0}^{2k_2+2}] = 0;$$

$$(54) \quad [P_{ij}^{2k_1}; P_{i^0 j^0}^{2k_2+2}] = 0 \quad \text{for } k_1 = 1;$$

$$(55) \quad [P_{ij}^{2k_1}; P_{i^0 j^0}^{2k_2+2}] = P_{jj^0}^{2(k_1+k_2+1)} \quad \text{for } k_1 = 1;$$

5. Miura type transformations

The definition of Miura type transformations was given in the introduction. Consider an evolutionary system of order $d = 1$

$$(56) \quad u_t^i = F^i(u^1; \dots; u^m; u_1^1; \dots; u_1^m; \dots; u_d^1; \dots; u_d^m); \quad i = 1; \dots; m; \quad u_k^i = \frac{\partial^k u^i}{\partial x^k};$$

Suppose that for this system we have a ZCR of the form

$$(57) \quad [D_x + M; D_t + N] = D_x(N) - D_t(M) + [M; N] = 0;$$

$$M = M(u^1; \dots; u^m); \quad N = (u^1; \dots; u^m; u_1^1; \dots; u_1^m; \dots; u_{d-1}^1; \dots; u_{d-1}^m)$$

with values in a Lie algebra g . According to Proposition 1, such ZCR is determined by a homomorphism from the WE algebra to g .

Consider a homomorphism from g to the Lie algebra of vector fields on an m -dimensional manifold W . Let $w^1; \dots; w^m$ be local coordinates in W and

$$(58) \quad (M) = \sum_{i=1}^m a^i(w^1; \dots; w^m; u^1; \dots; u^m) \frac{\partial}{\partial w^i};$$

$$(59) \quad (N) = \sum_{i=1}^m b^i(w^1; \dots; w^m; u^1; \dots; u^m; \dots; u_{d-1}^1; \dots; u_{d-1}^m) \frac{\partial}{\partial w^i};$$

Then relation (57) implies that the system

$$(60) \quad \frac{\partial w^i}{\partial x} = a^i(w^1; \dots; w^m; u^1; \dots; u^m);$$

$$(61) \quad \frac{\partial w^i}{\partial t} = b^i(w^1; \dots; w^m; u^1; \dots; u^m; \dots; u_{d-1}^1; \dots; u_{d-1}^m);$$

$$u_t^i = F^i(u^1; \dots; u^m; u_d^1; \dots; u_d^m); \quad i = 1; \dots; m;$$

is consistent. Suppose that from equations (60) one can express locally

$$(62) \quad u^i = c^i(w^1; \dots; w^m; \frac{\partial w^1}{\partial x}; \dots; \frac{\partial w^m}{\partial x}); \quad i = 1; \dots; m;$$

Substituting this to (61), we obtain an evolutionary system for $w^1(x; t); \dots; w^m(x; t)$ connected by a Miura type transformation (62) with system (56). This is the simplest case of a more general method to obtain Miura type transformations from ZCRs [13].

Let us apply this construction to system (16) and its ZCR (17), (18), where $s_1; \dots; s_n$ are again treated as complex parameters satisfying equations (2) and $S = (s^1; \dots; s^n)$ is given by formulas (15). This ZCR takes values in the Lie subalgebra $so_{n,1} \oplus gl_{n+1}(\mathbb{C})$ generated by the elements $A_i = E_{i,n+1} + E_{n+1,i}$, $i = 1; \dots; n$.

Consider the canonical action of the group $GL_{n+1}(\mathbb{C})$ on the projective space CP^n . It determines a homomorphism ρ from $gl_{n+1}(\mathbb{C})$ to the Lie algebra of vector fields on CP^n .

In the standard coordinates $w^1, \dots, w^n$ of an affine chart C^n of CP^n this homomorphism reads

$$\rho: E_{ij} \mapsto w^i \frac{\partial}{\partial w^j} - \sum_{l=1}^n w^l \frac{\partial}{\partial w^l} \quad \text{for } i, j = 1, \dots, n+1; w^{n+1} = 1.$$

In particular,

$$(63) \quad \rho(A_i) = \frac{\partial}{\partial w^i} - \sum_{l=1}^n w^l \frac{\partial}{\partial w^l}.$$

Clearly, vector fields (63) are tangent to the $(n-1)$ -dimensional submanifold

$$W = \{w^1, \dots, w^n \mid \sum_{j=1}^n (w^j)^2 = 1\} \subset C^n.$$

On some neighborhood $W^0 \subset W$ of the point $(w^1 = \dots = w^{n-1} = 0; w^n = 1) \in W$ we can take $w^1, \dots, w^{n-1}$ as local coordinates and

$$(64) \quad w^n = \frac{1}{\sqrt{1 - \sum_{j=1}^{n-1} (w^j)^2}}.$$

Thus we obtain the following homomorphism from so_{n+1} to the Lie algebra of vector fields on W^0

$$\begin{aligned} (A_i) = \rho(A_i)|_{W^0} &= \frac{\partial}{\partial w^i} - \sum_{l=1}^n w^l \frac{\partial}{\partial w^l}; \quad i = 1, \dots, n-1; \\ (A_n) = \rho(A_n)|_{W^0} &= \frac{1}{\sqrt{1 - \sum_{j=1}^{n-1} (w^j)^2}} \sum_{l=1}^n w^l \frac{\partial}{\partial w^l}. \end{aligned}$$

Applying this to (17), one gets

$$(65) \quad (M) = \sum_{i=1}^{n-1} s^i \frac{\partial}{\partial w^i} - \sum_{j=1}^n w^j \frac{\partial}{\partial w^j} \frac{\partial}{\partial w^i};$$

where one assumes (64), (15). Recall that the right-hand side of (60) is obtained from (58). Using (65), (15), (64), we get the following form of equations (60) in our case

$$(66) \quad \frac{\partial w^i}{\partial x} = \frac{1}{1 + \sum_{j=1}^{n-1} (u^j)^2} \left(2u^i \frac{\partial}{\partial x} - \sum_{j=1}^{n-1} u^j w^j \frac{\partial}{\partial w^j} - \sum_{j=1}^n w^j \frac{\partial}{\partial w^j} \right) \frac{\partial}{\partial w^i}; \quad i = 1, \dots, n-1;$$

Computing (N) by means of (18) and (65), from the coefficients of the vector field (59) we obtain equations (61) of the following form

$$(67) \quad \frac{\partial w^i}{\partial t} = \sum_{i=1}^{n-1} s_{xx}^i \frac{\partial}{\partial w^i} - \sum_{j=1}^n w^j \frac{\partial}{\partial w^j} \left(s_{xx}^j \frac{\partial}{\partial w^j} + \sum_{l=1}^{n-1} s_{xl}^j \frac{\partial}{\partial w^l} \right) + \left(r_1 + \frac{1}{2} + \frac{1}{2} (S; RS) + \frac{3}{2} (S_x; S_x) \right) \sum_{j=1}^n w^j \frac{\partial}{\partial w^j} \frac{\partial}{\partial w^i}; \quad i = 1, \dots, n-1;$$

where $S = (s^1, \dots, s^n)$ is given by (15) and w^n is equal to (64).

Denote by $a^i = a^i(w^1, \dots, w^{n-1}; u^1, \dots, u^{n-1})$ the right-hand side of (66). It is easily seen that the $(n-1) \times (n-1)$ -matrix $\| \partial a^i / \partial u^j \|$ at the point

$$(68) \quad w^1 = \dots = w^{n-1} = u^1 = \dots = u^{n-1} = 0$$

is equal to $\text{diag}(2^{-1}, \dots, 2^{-n-1})$. Suppose that

$$(69) \quad a_i \neq 0; \quad i = 1, \dots, n-1;$$

then, by the implicit function theorem, on a neighborhood of the point (68) from equations (66) we can express

$$(70) \quad u^i = c^i(w^1, \dots, w^{n-1}; \frac{\partial w^1}{\partial x}, \dots, \frac{\partial w^{n-1}}{\partial x}); \quad i = 1, \dots, n-1;$$

Substituting (70) to (67), one gets an evolutionary system of the form

$$(71) \quad w_t^i = H^i(w^1, \dots, w^{n-1}; w_{xxx}^1, \dots, w_{xxx}^{n-1}); \quad i = 1, \dots, n-1;$$

connected with system (16) by the Miura type transformation (70).

It remains to study the case when (69) does not hold. That is, $a_k = 0$ for some $1 \leq k \leq n-1$. Since in equations (2) one has $r_i \neq r_j$ for $i \neq j$, we obtain that $a_i \neq 0$ for $i \neq k$. Then at any point of the form

$$(72) \quad u^k = 1; \quad u^1 = w^1 = 0 \quad \forall 1 \leq k$$

one has

$$\frac{\partial a^i}{\partial u^j} = 0 \quad \forall i \neq j; \quad \frac{\partial a^i}{\partial u^i} = 2^{-i} \quad \forall i \neq k; \quad \frac{\partial a^k}{\partial u^k} = 2w^{n-1} \frac{P}{1 - (w^k)^2};$$

Therefore, at any point of the form (72) with $w^k \neq 0$, 1 the matrix $\| \partial a^i / \partial u^j \|$ is non-singular, and on a neighborhood of such point from equations (66) we can again get expressions of the form (70) and proceed as described above.

Appendix: proof of Lemma 1

We prove this by induction on $k_1 + k_2$. For $k_1 + k_2 = 0$ (that is, $k_1 = k_2 = 0$) the statements follow easily from (42) and (43). Assume that all the statements are valid for $k_1 + k_2 \leq m$ for some $m \in \mathbb{Z}_+$. We must prove them for $k_1 + k_2 = m + 1$.

Below l is an arbitrary integer such that $1 \leq l \leq n$, $l \neq i$, $l \neq j$. We use the notation from Section 4.

Proof of (48). First, note that by the induction assumption for $q \leq m$ one has

$$\| \dot{P}_i \dot{P}_j \|_{2q} = \| P_i^1 P_j^{2q} \|, \quad \| P_j^{2q+1} \| = \| P_i^1 P_j^{2q+1} \|, \quad \| P_j^{2q+2} \| = \| P_i^1 P_j^{2q+2} \|, \quad P_j^{2q+3};$$

Similarly, we obtain

$$(73) \quad \| \dot{P}_i \dot{P}_j \|_{2q}' = \| \dot{P}_i \dot{P}_j \|_{2q+2}' = \| \dot{P}_i \dot{P}_j \|_{2q+2}' = P_j^{2q+3} \quad \forall q \leq m;$$

Without loss of generality we can assume $k_2 \geq 1$ in (48). Using (73) and the Jacobi identity, one gets

$$(74) \quad \| \dot{P}_i \dot{P}_j \|_{2k_1} \| \dot{P}_i \dot{P}_j \|_{2k_2} = \| \dot{P}_i \dot{P}_j \|_{2k_1} \| \dot{P}_i \dot{P}_j \|_{2k_2-2} = \| \dot{P}_i \dot{P}_j \|_{2k_1} \| \dot{P}_i \dot{P}_j \|_{2k_1} + \| \dot{P}_i \dot{P}_j \|_{2k_1} \| \dot{P}_i \dot{P}_j \|_{2k_1} \\ = \| \dot{P}_i \dot{P}_j \|_{2k_1} \| \dot{P}_i \dot{P}_j \|_{2k_1} - 2 \| \dot{P}_i \dot{P}_j \|_{2k_1} \| \dot{P}_i \dot{P}_j \|_{2k_1};$$

To obtain the last line, we used the fact that

$$[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-2}' - [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-2}';$$

because $[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-2}' = 0$ by the induction assumption.

Since, by (73), $[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1}' = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1+2}$, from (74) one obtains

$$[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-2}' - [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1+2} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-2} = 2 [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-2}';$$

If $k_2 = 2$, applying the same procedure to $[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1+2} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-2}$ yields

$$[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-2}' - [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1+4} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-4} = 2 [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-2} - 2 [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1+2} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2-4}';$$

Thus applying this procedure several times to the first summand of the right-hand side, we obtain

$$(75) \quad [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2}' - [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+k_2)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2}' = \sum_{s=1}^{k_2} 2 [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)}';$$

By the induction assumption and (73), one has

$$[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)}' = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)+2}^{1, P_j} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)}^{1, P_i} = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)+2}^{2(k_1+s)-1, P_j} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)}^{1, P_i} = 0;$$

Therefore,

$$(76) \quad [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2}' - [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)}] = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)}];$$

Since, by the induction assumption and (73),

$$[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)}' = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)+2}^{1, P_j} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)+1}^{1, P_j} = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)+2}^{2(k_2-s)+1, P_j} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)+1}^{1, P_j} = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)}];$$

$$[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)}' = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s)}];$$

equation (76) implies

$$[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2}' - [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)}] = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+s-1)}] - [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_2-s)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+k_2)}];$$

Combining this with (75) yields

$$(77) \quad [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_1} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2}' - [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+k_2)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2}' = 2 [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+k_2)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2}';$$

For $k_1 = 0$ this equation implies $[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2(k_1+k_2)} [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2k_2}' = 0$. Combining this with (77), we obtain (48).

Proof of (49). By the induction assumption, (48), and (42),

$$[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2m+1}' = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2m+1}]; \quad [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2m}' = 0;$$

$$[\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2m}' = 0; \quad [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2m}' = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2m}]; \quad [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2m}' = [\![\begin{smallmatrix} \vdots \\ \vdots \\ \vdots \end{smallmatrix}\!]\!]_{2m}];$$

Using this and (43), one gets

$$\begin{aligned}
P_{ij}^{2(k_1+k_2+1)} &= [\underset{2m+3}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] = [\underset{2m+1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]' \quad [\underset{2m+1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]i' \\
[\underset{2m}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]j' & \quad [\underset{2m}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]i = [\underset{2m}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]iij' \quad [\underset{2m}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]iij = [\underset{2m}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]iijj = \\
[\underset{2m}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]iijj &= [\underset{2m}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]iijj' \quad [\underset{2m}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]jijj = P_{ji}^{2(k_1+k_2+1)} :
\end{aligned}$$

P roof of (50). By the Jacobi identity,

$$(78) \quad P_{ij}^{2k_1}; P_{ij}^{2k_2+2} = [\underset{2k_1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] [\underset{2k_2+1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] = [\underset{2k_1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] [\underset{2k_2}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] + [\underset{2k_1-1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] [\underset{2k_2}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] :$$

By the induction assumption and (73),

$$[\underset{2k_1-1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] [\underset{2k_2}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]' \quad P_{ij}^{2k_1} P_j^{2k_2+1} \quad P_i^{2(k_1+k_2)+1} :$$

Substituting this to (78) and using (48), we obtain

$$P_{ij}^{2k_1}; P_{ij}^{2k_2+2} \quad [\underset{2k_1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] [\underset{2k_2}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] + [\underset{2k_1-1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] P_i^{2(k_1+k_2)+1} \quad 0 :$$

P roof of (51). By (73) and the induction assumption of (53), for any $q_1; q_2 \in \mathbb{Z}_+$ such that $q_1 + q_2 \in 2(k_1 + k_2) - 1$ and $q_1 + q_2$ is odd one has

$$[\underset{q_1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] [\underset{q_2}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]' \quad 0 :$$

Using this and the Jacobi identity, we get

$$\begin{aligned}
[\underset{2k_1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] [\underset{2k_2}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]' & \quad [\underset{2k_1+1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] [\underset{2k_2-1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]' \\
[\underset{2k_1+2}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] [\underset{2k_2-2}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]' & \quad [\underset{2(k_1+k_2)}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] = [\underset{2(k_1+k_2)}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]i
\end{aligned}$$

Combining this with (73) and (49) yields

$$\begin{aligned}
P_i^{2k_1+1}; P_j^{2k_2+1} & \quad [\underset{2k_1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] [\underset{2k_2}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]' \\
[\underset{2(k_1+k_2)}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]i' & \quad [\underset{2(k_1+k_2)}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]i = P_{ji}^{2(k_1+k_2+1)} \quad P_{ij}^{2(k_1+k_2+1)} :
\end{aligned}$$

P roof of (52). Consider first the case $k_1 = 0$, then

$$P_i^{2k_1+1}; P_{ij}^{2k_2+2} = [\underset{2k_2+1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] :$$

Set $c = 1$ if $j \neq 1$ and $c = 2$ if $j = 1$. If $i = c$ then $[\underset{2k_2+1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}] = P_j^{2(k_1+k_2)+3}$ and the proof is complete.

Suppose that $i \neq c$, then using (49) one gets

$$P_i^{2k_1+1}; P_{ij}^{2k_2+2} = [\underset{2k_2+1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]' \quad [\underset{2k_2+1}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]i :$$

Since, by (48), $[\underset{2k_2}{\begin{smallmatrix} i & j \\ \vdots & \vdots \\ \vdots & \vdots \end{smallmatrix}}]i' = 0$, using (73) and $[[ij]c] = 0$ we obtain

$$\begin{aligned}
& \mathbb{P}_i^{2k_1+1}; \mathbb{P}_{ij}^{2k_2+2}]', \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]j] ', \quad [\underset{2k_2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[ij]j] ', \quad [\underset{2k_2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[ij]\underset{\{\underline{z}\}}{\mathbb{P}}i] = \\
& \quad \underset{2k_2}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{\{\underline{z}\}}{\mathbb{P}}[ij]i] = \underset{2k_2}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{\{\underline{z}\}}{\mathbb{P}}[ij]] ', \quad \underset{2k_2}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{\{\underline{z}\}}{\mathbb{P}}[ccj]] = \mathbb{P}_j^{2(k_1+k_2)+3}:
\end{aligned}$$

Now consider the case $k_1 = 1$. By (73), for any $1 \notin i, 1 \notin j$ we have

$$\mathbb{P}_i^{2k_1+1}; \mathbb{P}_{ij}^{2k_2+2}] = \mathbb{P}_{ij}^{2k_2+2}; \mathbb{P}_i^{2k_1+1}]', \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j]] \underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]:$$

Since due to the induction assumption of (53) one has $[\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j]]' = 0$, using (73) we get

$$\mathbb{P}_i^{2k_1+1}; \mathbb{P}_{ij}^{2k_2+2}]', \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j]] \underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]i] ', \quad \underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j]]i] = \underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_2+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j]] ', \quad [\underset{2(m+2)}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j]]:$$

In the consideration of the case $k_1 = 0$ we showed that for any $i \notin j$ $[\underset{2(m+2)}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j]]' = \mathbb{P}_j^{2m+5}$. (Recall that $m = k_1 + k_2 - 1$). Therefore,

$$[\underset{2(m+2)}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j]]' = \mathbb{P}_j^{2(k_1+k_2)+3}:$$

Proof of (53). Consider first the case $k_1 = 1$. Since $n = 4$, there is $1 \leq f_1, \dots, f_m$ such that $1 \notin i, 1 \notin i^0, 1 \notin j^0$. Then, by the induction assumption of (53),

$$[\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]]' = 0; \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]i]' = 0:$$

Using this and (73), one gets

$$(79) \quad \mathbb{P}_i^{2k_1+1}; \mathbb{P}_{i^0j^0}^{2k_2+2}] = \mathbb{P}_{i^0j^0}^{2k_2+2}; \mathbb{P}_i^{2k_1+1}]', \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]] \underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i]i] ', \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]i]i] ', \quad 0:$$

If we set $k_2 = 0$ then (79) implies that for any distinct integers $c_1; c_2; c_3; c_4 \geq 1; \dots; n$

$$(80) \quad [\underset{2m+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_1c_2]\underset{2m+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_3c_4]]' = 0:$$

Since $[\underset{2m+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_1c_2]\underset{2m+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_3c_4]]' = [\underset{2m+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_2c_1]\underset{2m+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_3c_4]]'$ due to (73), from (80) we obtain

$$(81) \quad [\underset{2m+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_1c_2]\underset{2m+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_3c_4]]' = 0:$$

By the Jacobi identity, (81), and (73),

$$(82) \quad [\underset{2m+3}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_1\underset{2m+3}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_2c_3c_4]]' = [\underset{2m+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_2c_1\underset{2m+2}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_3c_4]]' = [\underset{2m+3}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_2\underset{2m+3}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_1c_3c_4]]:$$

Also, property (49) implies

$$(83) \quad [\underset{2m+3}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_1\underset{2m+3}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_2c_3c_4]]' = [\underset{2m+3}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_1\underset{2m+3}{\underset{\{\underline{z}\}}{\mathbb{P}}}[c_3c_2c_4]]:$$

It remains to study the case $k_1 = 0$. Using (82) and (83), we get

$$\begin{aligned}
& \mathbb{P}_i^{2k_1+1}; \mathbb{P}_{i^0j^0}^{2k_2+2}] = [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]] ', \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]i] ', \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]i] ', \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]i] ', \\
& \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]i] ', \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]i] ', \quad [\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[i^0]\underset{2k_2+1}{\underset{\{\underline{z}\}}{\mathbb{P}}}[j^0]i] = \mathbb{P}_i^{2k_1+1}; \mathbb{P}_{i^0j^0}^{2k_2+2}]:
\end{aligned}$$

Therefore, $\mathbb{P}_i^{2k_1+1}; \mathbb{P}_{i^0j^0}^{2k_2+2}]' = 0$.

Proof of (54). By (53),

$$[i; P_{i^0 j^0}^{2k_2+2}]' = 0; \quad [j; P_{i^0 j^0}^{2k_2+2}]' = 0:$$

Obviously, this implies (54).

Proof of (55). By property (53), $[[i; \begin{smallmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{smallmatrix} j] j^0]' = 0$. Using this, equation (73), and (49), one obtains

$$\begin{aligned} [P_{ij}^{2k_1}; P_{ij^0}^{2k_2+2}]', \quad [P_{ij}^{2k_1}; P_{j^0 i}^{2k_2+2}] &= [[i; \begin{smallmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{smallmatrix} j] j^0]' \begin{smallmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{smallmatrix} i]' + \begin{smallmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{smallmatrix} j^0]' [i; \begin{smallmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{smallmatrix} j] j]' = \\ & \begin{smallmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{smallmatrix} j^0]' \begin{smallmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{smallmatrix} i]' + \begin{smallmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{smallmatrix} j^0]' \begin{smallmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{smallmatrix} j]' = P_{j^0 j}^{2(k_1+k_2+1)}, \quad P_{j j^0}^{2(k_1+k_2+1)}: \end{aligned}$$

Acknowledgements

The authors thank T. Skrypnyk and V. V. Sokolov for helpful discussions. Work of S.I. is supported by the NW OVENI grant 639.031.515. JvdL is partially supported by the European Union through the FP6 Marie Curie Grant (ENIGMA) and the European Science Foundation (MISGAM). S.I. thanks the Max Planck Institute for Mathematics (Bonn, Germany) for its hospitality and excellent working conditions during 02.2006–01.2007, when a part of this research was done.

References

- [1] M. Ju. Balakhnev. The vector generalization of the Landau-Lifshitz equation: Backlund transformation and solutions. *Appl. Math. Lett.* 18 (2005), 1363{1372.
- [2] A. V. Bocharov, V. N. Chetverikov, S. V. Duzhin, N. G. Khor'kova, I. S. Krasil'shchik, A. V. Samokhin, Yu. N. Torkhov, A. M. Verbovetsky, and A. M. Vinogradov. *Symmetries and Conservation Laws for Differential Equations of Mathematical Physics*. Amer. Math. Soc., Providence, RI, 1999.
- [3] C. Hoenselaers and W. K. Schief. Prolongation structures for Harry Dym type equations and Backlund transformations of CC ideals. *J. Phys. A: Math. Gen.* 25 (1992), 601{622.
- [4] R. Dodd and A. Fordy. The prolongation structures of quasipolynomial flows. *Proc. Roy. Soc. London Ser. A* 385 (1983), 389{429.
- [5] H. N. van Eck. A non-Archimedean approach to prolongation theory. *Lett. Math. Phys.* 12 (1986), 231{239.
- [6] L. Feher and I. Marshall. Extensions of the matrix Gelfand-Dickey hierarchy from generalized Drinfeld-Sokolov reduction. *Comm. Math. Phys.* 183 (1997), 423{461.
- [7] I. Z. Golubchik and V. V. Sokolov. Generalized Heisenberg equations on Z-graded Lie algebras. *Theoret. and Math. Phys.* 120 (1999), 1019{1025.
- [8] I. Z. Golubchik and V. V. Sokolov. A multicomponent generalization of the hierarchy of the Landau-Lifshitz equation. *Theoret. and Math. Phys.* 124 (2000), 909{917.
- [9] M. F. de Groot, T. J. Hollowood and J. L. M. Miramontes. Generalized Drinfeld-Sokolov hierarchies. *Comm. Math. Phys.* 145 (1992), 57{84.
- [10] L. D. Faddeev and L. A. Takhtajan. *Hamiltonian methods in the theory of solitons*. Springer-Verlag, Berlin, 1987.
- [11] S. Igonin. Coverings and fundamental algebras for partial differential equations. *J. Geom. Phys.* 56 (2006), 939{998.
- [12] S. Igonin. Analogues of coverings and the fundamental group in the category of PDEs. To appear.
- [13] S. Igonin. Miura type transformations and homogeneous spaces. *J. Phys. A: Math. Gen.* 38 (2005), 4433{4446.
- [14] S. Igonin and R. Martini. Prolongation structure of the Krichever-Novikov equation. *J. Phys. A: Math. Gen.* 35 (2002), 9801{9810.
- [15] I. S. Krasil'shchik and A. M. Vinogradov. Nonlocal trends in the geometry of differential equations. *Acta Appl. Math.* 15 (1989), 161{209.
- [16] I. M. Krichever and S. P. Novikov. Algebras of Virasoro type, Riemann surfaces and the structures of soliton theory. *Functional Anal. Appl.* 21 (1987), 126{142.

- [17] M. Marvan. On zero curvature representations of partial differential equations, in: *Differential Geometry and Its Applications*, Proc. Conf. Opava, Czechoslovakia, Aug. 24{28, 1992 (Silesian University, Opava, 1993), 103{122 (<http://www.emis.de/proceedings/5ICDGA/>).
- [18] M. Marvan. On the spectral parameter problem. Preprint at <http://arxiv.org/abs/0804.2031>
- [19] G. H. M. Roelofs and R. Martini. Polongation structure of the Landau-Lifshitz equation. *J. Math. Phys.* 34 (1993), 2394{2399.
- [20] S. Yu. Sakovich. Cyclic bases of zero-curvature representations: five illustrations to one concept. *Acta Appl. Math.* 83 (2004), 69{83.
- [21] W. F. Shadwick. The Backlund problem for the equation $\partial^2 z = \partial x^1 \partial x^2 = f(z)$. *J. Math. Phys.* 19 (1978), 2312{2317.
- [22] T. Skrypnyk and P. H. Od. Hierarchies of integrable equations associated with hyperelliptic Lie algebras. *J. Phys. A: Math. Gen.* 34 (2001), 11123{11137.
- [23] T. Skrypnyk. Quasigraded Lie algebras on hyperelliptic curves and classical integrable systems. *J. Math. Phys.* 42 (2001), 4570{4581.
- [24] T. Skrypnyk. Spin generalizations of Calogero and Neumann integrable systems. *J. Phys. A* 36 (2003), 4407{4416.
- [25] T. Skrypnyk. Deformations of loop algebras and integrable systems: hierarchies of integrable equations. *J. Math. Phys.* 45 (2004), 4578{4595.
- [26] H. D. Wahlquist and F. B. Estabrook. Polongation structures of nonlinear evolution equations. *J. Math. Phys.* 16 (1975), 1{7.

Sergey Igonin, Department of Mathematics, University of Utrecht, P.O. Box 80010,
3508 TA Utrecht, The Netherlands
E-mail address: igonin@mccme.ru

Johan van de Leur, Department of Mathematics, University of Utrecht, P.O. Box
80010, 3508 TA Utrecht, The Netherlands
E-mail address: J.W.vandeLeur@uu.nl

Gianni Manno, Università degli Studi di Milano-Bicocca, Dipartimento di Matematica
e Applicazioni, Via Cozzi 53, 20125 Milano, Italy
E-mail address: gianni.manno@unimib.it

Vladimir Trushkov, University of Pereslavl', Sovetskaya 2, 152020 Pereslavl'-Zalessky,
Yaroslavl region, Russia
E-mail address: vladimir@trushkov.pereslavl.ru