## BETTI NUMBERS AND INJECTIVITY RADII

## MARC CULLER AND PETER B. SHALEN

The theme of this paper is the connection between topological properties of a closed orientable hyperbolic 3-manifold M and the maximal injectivity radius of M. In [4] we showed that if the first Betti number of M is at least 3 then the maximal injectivity radius of M is at least log 3. By contrast, the best known lower bound for the maximal injectivity radius of M with no topological restriction on M is the lower bound of  $\operatorname{arcsinh}(\frac{1}{4}) = 0.24746...$  due to Przeworski [7]. One of the results of this paper, Corollary 4, gives a lower bound of 0.32798 for the case where the first Betti number of M is 2 and M does not contain a "fibroid" (see below). Our main result, Theorem 3, is somewhat stronger than this

The proofs of our results combine a result due to Andrew Przeworski [7] with results from [5] and [6].

The results of [5] and [6] were motivated by applications to the study of hyperbolic volume, and these applications were superseded by the results of [2]. The applications presented in the present paper do not seem to be accessible by other methods.

\* \* \*

As in [5], we define a book of *I*-bundles to be a compact, connected, orientable topological 3-manifold (with boundary) W which has the form  $W = \mathcal{P} \cup \mathcal{B}$ , where

- $\mathcal{P}$  is an *I*-bundle over a non-empty compact 2-manifold-with-boundary;
- each component of  $\mathcal{B}$  is homeomorphic to  $D^2 \times S^1$ ;
- the set  $\mathcal{A} = \mathcal{P} \cap \mathcal{B}$  is the vertical boundary of the *I*-bundle  $\mathcal{P}$ ; and
- each component of  $\mathcal{A}$  is an annulus in  $\partial \mathcal{B}$  which is homotopically non-trivial in  $\mathcal{B}$ .

(Note that this terminology differs slightly from that of [1], where it is the triple  $(W, \mathcal{B}, \mathcal{P})$  that is called a book of *I*-bundles.)

As in [5], we define a *fibroid* in a closed, connected, orientable 3-manifold M to be a connected incompressible surface F with the property that each component of the compact manifold obtained by cutting M along F is a book of I-bundles. Note that in defining a fibroid to be connected, we are following the convention of [5] rather than that of [6].

We define a function R(x) for x > 0 by

(1) 
$$R(x) = \frac{1}{2}\operatorname{arccosh}\left(\frac{e^{2x} + 2e^x + 5}{(\cosh\frac{x}{2})(e^x - 1)(e^x + 3)}\right).$$

1991 Mathematics Subject Classification. Primary 57M50; Secondary 57N10. Partially supported by NSF grants DMS-0504975 and DMS-0608567.

As in [3, Section 10], we define a function V(x) for x > 0 by

(2) 
$$V(x) = \pi x \sinh^2 R(x) = \frac{\pi x}{e^x - 1} \left( \frac{e^{2x} + 2e^x + 5}{2(e^x + 3)\cosh(x/2)} \right) - \frac{\pi x}{2}.$$

Thus, in a closed hyperbolic 3-manifold, if a geodesic of length  $\ell$  is the core of an embedded tube of radius  $R(\ell)$ , then this tube has volume  $V(\ell)$ .

The following result is implicit in [6], but we will supply a proof for the sake of comprehensibility.

**Proposition 1.** Let M be a closed hyperbolic 3-manifold. Suppose that there is an infinite subset  $\mathcal{N}$  of  $H_2(M;\mathbb{Z})$  such that every element of  $\mathcal{N}$  is represented by some connected, incompressible surface which is not a fibroid. Let  $\lambda$  be a positive number less than  $\log 3$ . Then either the maximal injectivity radius of M is at least  $\lambda/2$ , or M contains a closed geodesic C of length at most  $\lambda$  such that the maximal tube about C has radius at least  $R(\lambda)$  and volume at least  $V(\lambda)$ , where  $R(\lambda)$  and  $V(\lambda)$  are defined by (1) and (2).

Proof. The hypothesis implies in particular that  $H_2(M; \mathbb{Z})$  has infinitely many primitive elements, and so the first Betti number  $\beta_1(M)$  is at least 2. If  $\beta_1(M) \geq 3$ , then according to [4, Corollary 10.4], the maximal injectivity radius is at least  $\frac{1}{2} \log 3 > \lambda/2$ . We may therefore assume that  $\beta_1(M) = 2$ . Hence the quotient of  $H_1(M, \mathbb{Z})$  by its torsion subgroup is a free abelian group L of rank 2. We let  $h: \pi_1(M) \to L$  denote the natural homomorphism.

We distinguish two cases. First consider the case in which M contains a non-trivial closed geodesic C of some length  $\ell < \lambda$  such that the conjugacy class represented by C is contained in the kernel of h. Let T denote the maximal embedded tube about C. According to [3, Corollary 10.5] we have Vol  $T \ge V(\lambda)$ . If  $\rho$  denotes the radius of T, this gives

$$\pi \ell \sinh^2 \rho = \operatorname{Vol} T \ge V(\lambda) = \pi \lambda \sinh^2 R(\lambda) > \pi \ell \sinh^2 R(\lambda),$$

and hence  $\rho > R(\lambda)$ .

Thus the second alternative of the proposition holds in this case.

We now turn to the case in which no non-trivial closed geodesic of length  $< \lambda$  represents a conjugacy class contained in the kernel of h. Since M is closed, there are only a finite number  $n \geq 0$  of conjugacy classes in  $\pi_1(M)$  that are represented by closed geodesics of length < l. Let  $\gamma_1, \ldots, \gamma_n$  be elements belonging to these n conjugacy classes. Then  $\bar{\gamma}_i = h(\gamma_i)$  is a non-trivial element of L for  $i = 1, \ldots, n$ . Since L is a free abelian group of rank 2, there exists, for each  $i \in \{1, \ldots, n\}$ , a homomorphism  $\phi_i$  of L onto  $\mathbb{Z}$  such that  $\phi_i(\bar{\gamma}_i) = 0$ . Because  $\bar{\gamma}_i \neq 0$ , the epimorphism  $\phi_i$  is unique up to sign.

The epimorphism  $\phi_i \circ h \colon \pi_1(M) \to \mathbb{Z}$  corresponds to a primitive element of  $H^1(M; \mathbb{Z})$ , whose Poincaré dual in  $H_2(M; \mathbb{Z})$  we shall denote by  $c_i$ . Since the set  $\mathcal{N} \subset H_2(M; \mathbb{Z})$  given by the hypothesis of the theorem is infinite, there is an element c of  $\mathcal{N}$  which is distinct from  $\pm c_i$ for  $i = 1, \ldots, n$ . Since  $c \in \mathcal{N}$  it follows from the hypothesis that there is a connected incompressible surface  $S \subset M$  which represents the homology class c and is not a fibroid.

We now apply Theorem A of [5], which asserts that if S is a connected non-fibroid incompressible surface in a closed, orientable hyperbolic 3-manifold M, and if  $\lambda$  is any positive number, then either (i) M contains a non-trivial closed geodesic of length  $< \lambda$  which is homotopic in M to a closed curve in M - S, or (ii) M contains a hyperbolic ball of radius  $\lambda/2$ . In the present situation, with  $\lambda$  chosen as above, we claim that alternative (i) of the conclusion of Theorem A of [5] cannot hold.

Indeed, suppose that C is a non-trivial closed geodesic of length  $<\lambda$  with the properties stated in (i). Since C has length  $<\lambda$ , the conjugacy class represented by C contains  $\gamma_i$  for some  $i \in \{1, \ldots, n\}$ . Since C is homotopic to a closed curve in M-S it follows that the image of  $\gamma_i$  in  $H_1(M; \mathbb{Z})$  has homological intersection number 0 with c. Thus if  $\psi \colon \pi_1(M) \to \mathbb{Z}$  is the homomorphism corresponding to the Poincaré dual of c, we have  $\psi(\gamma_i) = 0$ . Now since C is the quotient of C is to subgroup, C factors as C is some homomorphism from C to C. Since C is primitive, C is surjective, and hence so is C. But we have C is C in view of the uniqueness that we observed above for C it follows that C is that C is that C in view of the uniqueness that we observed above for C it follows that C is that C is that C is a non-trivial closed geodesic of C in view of the uniqueness that we observed above for C it follows that C is a non-trivial closed geodesic of C in view of the uniqueness that we observed above for C in the contradicts our choice of C.

Hence (ii) must hold. This means that the maximal injectivity radius of M is at least  $\lambda/2$ . Thus the first alternative of the proposition holds in this case.

**Proposition 2.** Let M be a closed hyperbolic 3-manifold. Suppose that there is an infinite subset  $\mathcal{N}$  of  $H_2(M;\mathbb{Z})$  such that every element of  $\mathcal{N}$  is represented by some connected, incompressible surface which is not a fibroid. Then the maximal injectivity radius of M exceeds 0.32798.

*Proof.* We set  $\lambda = 2 \times 0.32798 = 0.65596$ .

According to Proposition 1, either the maximal injectivity radius of M is at least  $\lambda/2$ —so that the conclusion of the theorem holds — or M contains a closed geodesic C of length at most  $\lambda$  such that the maximal tube about C has volume at least  $V(\lambda)$ , where  $V(\lambda)$  is defined by (2). In the latter case, if R denotes the radius of T, we have

$$R \ge R(\lambda) = 0.806787...$$

Now according to [7, Proposition 4.1], the maximal injectivity radius of M is bounded below by

$$\operatorname{arcsinh}\left(\frac{\tanh R}{2}\right) > \operatorname{arcsinh}\left(\frac{\tanh 0.806787}{2}\right) > 0.32799.$$

This gives the conclusion of the theorem in this case.

**Theorem 3.** Let M be a closed hyperbolic 3-manifold. Suppose that there is an infinite set  $\mathcal{N}$  of primitive elements of  $H_2(M;\mathbb{Z})$  such that no element of  $\mathcal{N}$  is represented by a (connected) fibroid. Then the maximal injectivity radius of M exceeds 0.32798.

*Proof.* If  $\pi_1(M)$  has a non-abelian free quotient, then by [6, Theorem 1.3], the maximal injectivity radius of M is at least  $\frac{1}{2} \log 3 = 0.549...$  Now suppose that  $\pi_1(M)$  has no non-abelian free quotient. If  $\mathcal{N}$  is the set given by the hypothesis of Theorem 3, it now follows from

[6, Proposition 2.1] that every element of  $\mathcal{N}$  is represented by a connected incompressible surface, which by hypothesis cannot be a fibroid. Thus  $\mathcal{N}$  has the properties stated in the hypothesis of Proposition 2. The latter result therefore implies that the maximal injectivity radius of M exceeds 0.32798.

If M is a 3-manifold whose first Betti number is at least 2, then  $H_2(M; \mathbb{Z})$  has infinitely many primitive elements. If a non-trivial element of  $H_2(M; \mathbb{Z})$  is represented by a connected surface it must be primitive, since it has intersection number 1 with a class in  $H_1(M; \mathbb{Z})$ . Hence Theorem 3 implies:

**Corollary 4.** Let M be a closed hyperbolic 3-manifold. Suppose that the first Betti number of M is at least 2, and that M contains no non-separating fibroid. Then the maximal injectivity radius of M exceeds 0.32798.

## References

- [1] Ian Agol, Marc Culler, and Peter B. Shalen, Singular surfaces, mod 2 homology, and hyperbolic volume, I. arXiv:math.GT/0506396. To appear in Trans. Amer. Math. Soc.
- [2] Ian Agol, Peter A. Storm, and William P. Thurston, Lower bounds on volumes of hyperbolic Haken 3-manifolds, J. Amer. Math. Soc. **20** (2007), no. 4, 1053–1077 (electronic). With an appendix by Nathan Dunfield.
- [3] James W. Anderson, Richard D. Canary, Marc Culler, and Peter B. Shalen, Free Kleinian groups and volumes of hyperbolic 3-manifolds, J. Differential Geom. 43 (1996), no. 4, 738–782.
- [4] Marc Culler and Peter B. Shalen, Paradoxical decompositions, 2-generator Kleinian groups, and volumes of hyperbolic 3-manifolds, J. Amer. Math. Soc. 5 (1992), no. 2, 231–288.
- [5] \_\_\_\_\_, Volumes of hyperbolic Haken manifolds. I, Invent. Math. 118 (1994), no. 2, 285–329.
- [6] \_\_\_\_\_\_, Volumes of hyperbolic Haken manifolds. II, Proc. Amer. Math. Soc. **125** (1997), no. 10, 3059–3067.
- [7] Andrew Przeworski, Cones embedded in hyperbolic manifolds, J. Differential Geom. 58 (2001), no. 2, 219–232.

Department of Mathematics, Statistics, and Computer Science (M/C 249), University of Illinois at Chicago, 851 S. Morgan St., Chicago, IL 60607-7045

E-mail address: culler@math.uic.edu

Department of Mathematics, Statistics, and Computer Science (M/C 249), University of Illinois at Chicago, 851 S. Morgan St., Chicago, IL 60607-7045

E-mail address: shalen@math.uic.edu