Twisted speckle entities inside wavefront reversalm irrors.

A Yu O kulov

General Physics Institute of Russian Academy of Sciences Vavilova str. 38, 119991, Moscow, Russia

(Dated: June 09, 2009)

The previously unknown property of the optical speckle pattern reported. The interference of a speckle with the counter-propagating phase-conjugated speckle wave produces a random ly distributed ensemble of a twisted entities (ropes) surrounding optical vortex lines. These entities appear in a wide range of a random ly chosen speckle parameters inside the phase-conjugating m irrors regardless to an internal physical mechanism of the wavefront reversal. These numerically generated interference patterns are relevant to the Brillouin PC-m irrors and to a four-wave mixing PC-m irrors based upon laser trapped ultracold atom ic cloud.

PACS numbers: 42.30 M s 42.50.T x 42.65 H w 42.65 E s

Phase singularities of the optical beams attracted a substantial interest in a recent decades from the point of view of optical information processing [1]. Helical wavefronts had been shown to a ect a processes of the second harm onic generation [1], in age processing with a photorefractive m irrors [2], phase-conjugated re ection via nondegenerate four-wave mixing in a cold atom s cloud [3]. The photons with helical wavefunction had been shown to possess quantized angular momentum ~ per photon [4]. The optical nonlinearities are capable to transfer the angular m om entum from photons to an ensemble of ultracold atom s [3]. This e ect is considered as a possible tool for the light storage in addition to the slow light technique based on electrom agnetically induced transparency (EIT) [5, 6]. Recently the angularmomentum transfer to BEC cloud of sodium atoms via stimulated Raman scheme had been observed [7]. O f special interest is the nondegenerate four-wave mixing a cold atom ic cloud and phase conjugated (PC) re ection of the optical vortices (OV), i.e. Laguerre-Gaussian beams (LG) with helical wavefronts [3]. The re ection of the phase-conjugated LG photons from 107 cesium 10^{-3} K and reversal of the atoms cooled down to T orbital angular m om entum (OAM) had been interpreted as a consequence of internal macroscopic rotations inside atom ic cloud [8]. A fterwards these experim ental results were analyzed from a point of view of the angular momentum conservation for the incident and re ected photons from a Brillouin PC -mirror [9]. The goal of the current com m unication is to study the spiral interference patterns with the period of =2 around the nodes of optical speckle pattern. These coarse interference patterns tum in rotation the Brillouin medium [10] producing the acoustical vortices carrying OAM [11].

The traditional experim ental and num erical technique for the visualization of the optical phase singularities is based on m ixing of a wide quasi-plane wave reference optical beam with a speckle signal [12]. The dark spots (ze-

okulov@kapella.gpi.ru;

ros of am plitudes) appearing in intensity distribution are collocated with the helical ramps in phase distribution. The current com m unication reports a theoretical description of the alternative visualization technique which uses the interference of the speckle wave with the counterpropagating phase-conjugated one. In fact this technique is autom atically implemented in a wavefront reversing m irrors where a phase-conjugated wave produces near the bright spot the Bragg grating of dielectric perm ittivity of the form $\cos((k_p + k_s)z)$. In the vicinity of the dark line the modulation of the light intensity is more complex: the Bragg grating is transformed into spirals of the form $\cos((k_p + k_s)z + 2)$, where is the local azym uthalangle, ' is topological charge of the phase singularity, k_p ; k_s are wavenum bers of incident (pum p) and re ected (Stockes) waves respectively [9, 10]. In the real PC-m irrors the contrast of such gratings m ay be reduced due to the interference with a non-phase-conjugated com ponent of the radiation.

The novel feature compared to the previous ndings [12, 13] is that the ideal (or close to an ideal) PC m irror visualizes in the speckle patterns a peculiar optical entities, which m ight be called the ropes. Typically each such a rope is com posed of a 2-3 optical vortex lines (g.1). In contrast to the well studied straight optical vortex lines e.g. LG [1, 4], the OV in a speckle eld are self-twisted as it seen from g2. Having in m ind the well known fact [12], that OV appear and annihilate as a pairs of whirls with opposite circulations [9, 10] it is easy to conclude that at least two adjacent OV with opposite topological charges are needed to produce a rope (g.3). The mean length of each OV in Z direction is the Rayleigh range of a speckle pattern L_R $D^2 =$, where D is an average transverse size (in the plane X;Y) of the OV core, the wavelength = $2 = k_p$. The g.1 shows the num erically generated fragm ent of a speckle pattern which contains three ropes each composed of a set of a vortex lines. Let us describe the num erical procedure for generating the interference pattern of the two counter-propagating speckle elds.

The standard m odel of the phase conjugation via stim – ulated B rillouin scattering (SBS) is the B ragg re ection from the sound grating with period =2 moving with

Electronic address: URL:http://www.gpi.ru/~okulov



FIG. 1: Gray scale intensity plot for a small volume of a speckle pattern inside PC-m inform in (X;Z) -plane (Y = 23 m). The mean transverse size of a speckle is D 8 m. The mean longitudinal length of a speckle entity is of the order of the Rayleigh range L_R D^2 for 1 m. The size of the volume is 32 m in transverse (X;Y) directions and 128 m in longitudinal (Z) direction. The step of longitudinal modulation $\cos((k_p + k_s)z - 2)$ induced by the interference of counter-propagating pump and Stockes waves is enlarged here 8 times for visualization purposes. The characteristic - phase-jump is clearly visible in between adjacent Bragg's cos-m odulated roll patterns.

the speed of sound v_{ac} [13]. Due to the conservation of m om entum $p = -\tilde{\kappa_p} = -\tilde{\kappa_s} + -\tilde{\kappa_{ac}}$ the length of the wavevector of sound k_{ac} is equal to doubled length of wavevectorofpum p light k_p with an accuracy about 10⁵ [14]. The Doppler e ect de ness the small (of the order of 10⁵) frequency shift of the re ected (Stockes) wave $!_{ac} = 2!_p n v_{ac} = c$, where n is refractive index of medium, c is the speed of light in vacuum, $!_p;!_s$ is the frequency

of the pump and Stockes light respectively.

The evolution in space of the two counter-propagating paraxial laser beams inside B rillouin active medium obeys to M axwell equations with the cubic nonlinear polarization induced by electrostrictive e ect [13]. The linearly polarized pump eld E_p moves in a positive direction of Z axis, the relected Stockes eld E_s with the same polarization propagates in opposite direction (g.1). The acoustic eld Q is excited via electrostriction [13, 15]. The three-wave equations for the interaction of E_p , E_s with adiabatic elimination of the Q E_pE_s are:

$$\frac{@E_{p}(z;x;y;t)}{@z} + \frac{i}{2k_{p}} _{?} E_{p} = \frac{2!_{p}k_{a}^{2}}{32_{0}nc!_{ac}} E_{s}fE_{p};$$
(1)

$$\frac{@E_{s}(z;x;y;t)}{@z} = \frac{i}{2k_{s}} + E_{s} = \frac{2!_{s}k_{a}^{2}}{32_{0}nc!_{ac}} E_{p} f_{E_{s}};$$
(2)

where = (@ =@) is the electrostrictive coupling constant, $_0$ is the unpertubed density of medium [15]. Consider the phase conjugation with a random phase plate which produces the chaotic transverse modulation of the phase of the incident E_p with a characteristic size of the transverse inhom ogeneity of the order of 5 50 m [13]. In such a case the complex envelope amplitude of the inhom ogeneous speckle eld E_p at the entrance to PC-m irror is given as a multimode random eld [16]:

$$E_{p}(\boldsymbol{x}; z = 0) \quad E_{p}^{\circ} \qquad A_{j_{x}}; j_{y}$$

$$\stackrel{0 < j_{x}; j_{y} < N_{g}}{\sum_{j_{x} \neq j_{x}} p_{x}} \quad j_{x} + \frac{y_{y}}{p_{y}} \quad j_{y} + i_{j_{x}}; j_{y} g]; \quad (3)$$

where random phases $j_x; j_y$ are the random numbers from interval [0;], A $j_x; j_y$ are the real amplitudes of the spatial harm onics, $p_x; p_y$ are maximal transverse sizes (in X;Y-plane), r = (x; y) = (r;), $j_x; j_y$ are integers corresponding to N_g = (16;32;64) plane waves with random phases $j_x; j_y$. The random numbers $j_x = p_x = p_y^{\circ}$ and $j_y = p_y = p_y^{\circ}$ having the uniform distribution in the small vicinity of a 1 are responsible for a random tilt of a plane waves constituting the spatial Fourier spectrum of the light transmitted through a random phase plate. For the paraxial beam propagation the amplitude and phase structure of the com plex eld E_p have the following form [17] in an arbitrary Z-plane :

$$E_{p}(\mathbf{r}; z > 0) \quad E_{p}^{\circ} \exp(ik_{p}z) \qquad \exp[i_{j_{x}}; j_{y}] \\ \xrightarrow{0 < j_{x}} (j_{y} < N_{g}) \\ A_{j_{x}}; j_{y}} \exp[i2 f \frac{x}{p_{j_{x}}^{\circ}} + \frac{y}{p_{j_{x}}^{\circ}} + \frac{z}{p_{j_{y}}^{\circ}} + \frac{z}{2k_{p}} (\frac{j_{x}}{p_{j_{x}}^{\circ}} + \frac{j_{y}}{p_{j_{y}}^{\circ}})g];(4)$$

The interference pattern I(r;z) is responsible for the sound grating in a Brillouin PC-m irror. For the grating produced by a linear superposition of the random plane waves we have:



FIG.2: The plot of the light intensity maxim a at a given moment t in a small volume inside the PC-mirror. The four pairs of spiral interference patterns with opposite handedness are located random ly in space. Their diameters and directions are changed smoothly due to diraction. The step of longitudinalm odulation $\cos{(k_{\rm p}+k_{\rm s})z}=2$) is enlarged here 16 times.

$$I(x;z > 0) = \frac{1}{2}p(x;z) + E_s(x;z)f;$$
 (5)

where the phase-conjugated Stockes wave E_s is given by:

$$\begin{array}{c} \text{E}_{s}(\textbf{r};z>0) \quad \text{E}_{p}^{\circ} \exp(\text{i}k_{s}z) & \exp[\text{i}_{j_{x}};j_{y}] \\ \text{A}_{j_{x}};j_{y}} \exp[\text{i}2 \quad f\frac{x}{5}\frac{x}{p}\frac{y}{j_{x}} + \frac{y}{p}\frac{y}{j_{y}}\frac{y}{p} + \frac{z}{2k_{s}}(\frac{j_{x}^{2}}{p}\frac{z}{j_{x}}^{2} + \frac{j_{y}^{2}}{p}\frac{z}{j_{y}})g];(6) \end{array}$$

provided that $E_s = E_p$ (noiseless PC-m irror). Such an approach presumes the ideally perfect phase-conjugation with ultimate rejection R = 1 or interference of the non-interacting elds $E_s = E_p$ when the right hand sizes of Eq. (1,2) are equal to zero.

The intensity snapshots given by (5) were obtained num erically by summation of the series (4) and (6) on standard Intel platform with dual core 1.86 G hz processor and 1G bm em ory using standard educational software. The num erical simulations show that optical vortex lines intertwine and form ropes. The average length of a rope is of the order of Rayleigh range $D^2 = .$ This may be interpreted as a consequence of di ractive divergence within the angle =D (g.1). The longitudinal m odulation by the Bragg factor $\cos[(k_p + k_s)z = 2]$ is accompanied by an additional bending and twisting at characteristic wavelength $_{\rm K}$. The latter is of the order of a several tenth of optical , resem bling K elvin m odes of the vortex lines [18, 19]. The handedness of helical ropes changes random ly from one entity to another: the clockwise and counterclockwise ropes appear with the equal



FIG. 3: G ray scale intensity plot for the variational anzatz (7) substituted in (5). The 128 2048 points arrays are used for the intensity distributions (5) in X Z planes at xed Y = 64. The single sinusoidally m odulated LG_0^{-1} vortex line (see eq. (7)) is shown for M = 0:1 D (a) and M = D (b). Two LG_0^{-1} distant vortex lines LG_0^{-1} (c) with parallel (d) opposite (e) topological charges 'constitute the rope when the distance between them is reduced to zero (d,e). The step of longitudinal m odulation is enlarged here 32 times.

probability. Despite the statistical nature of a speckle pattern (g.1) the ropes are structurally stable: they appear at a di erent locations and have a variable sizes but their morphology is reproduced from a given statistical realization to another one.

The physical interpretation of these numerically observed patterns is given in the g.3 using the (5) with the electrical elds $E_p = E_s$ in the form of a two overlapping elementary LG₀¹ optical vortices:

$$E_{p} \exp (ik_{p}z (x_{1}^{2} + y_{1}^{2}) + i'_{1}) p \frac{p}{(x_{1}^{2} + y_{1}^{2})} + \exp (ik_{p}z (x_{2}^{2} + y_{2}^{2}) i'_{2}) p \frac{(x_{2}^{2} + y_{2}^{2})}{(x_{2}^{2} + y_{2}^{2})}; (7)$$

where the location of the vortex cores $(x_1D = x + M \cos(2 z_{=_K}); (y_1D = y + M \sin(2 z_{=_K}))$ and $(x_2D = x + M \cos(2 z_{=_K} + =2)); (y_2D = y + M \sin(2 z_{=_K} + =2))$ changes sinusoidally with period $_K$ along Z -axis, $_1$ and $_2$ are the local azim uthal angles around vortex cores. Such vortex-vortex (g.3d) and vortex-antivortex (g.3e) pair interference pattern is used here as a variational anzatz in eq.(5). The g.3 reproduces qualitatively a some features of the num erically generated speckle patterns (g.1).

In summary the nontrivial topology of the mul-

tiply connected optical speckle patterns was demonstrated num erically. Using the interference with the counter-propagating phase-conjugated speckle eld we have shown the hidden twisted geometry of the multimode wave eld composed of the random ly tilted plane waves [16]. Noteworthy the ropes exist without phaseconjugated counterpart: the re ection from PC-m irror makes these twisted entities visible due to the characteristic -phase-jump in between the adjacent Bragg's sinusoidally -m odulated roll patterns (g.1). Thus it seems probable that a rope entities exist in a wide class of a superpositional physical elds like an electrom agnetic or acoustical ones [11]. The other most evident examples are the blackbody radiation eld in a cavity or a monochromatic eld in a cavity with a rough mirrors. Taking into account the phenom ena of a so-called nonlinear superposition which take place for collisions of the optical solitons, vortices [20] and form ation of the

optical vortex lattices [21] one m ight expect the nding of a stable ropes in a nonlinear elds. The relevant exam – ples are photorefractive PC -m irrors [2], ultracold m atter, e.g. phase-conjugating PC -m irrors based on nondegenerate four-wave m ixing in a trapped gases [3, 8] and the situations of the slow [5, 6, 22] and the fast light [23, 24] propagation in a resonant m edium.

The reported result extends the set of the possible form s of the photon's wavefunction. In addition to the wellknown photon wavefunctions in the form of the elementary optical vortex lines, e.g. LG beam s which possess a helical wavefronts [1, 4], the twisted entities each composed of the several vortex lines o er a new form of the photon's wavefunction [25, 26].

A cknow ledgem ent. This work was supported in part by the Russian Fund of Basic Research G rant 08-02-01229.

- M.S.Soskin and M.V.Vasnetsov, "Progress in Optics", 42,pp.219-276, ed.E.W olf (Elsevier, Amsterdam) (2001).
- [2] A V M am aev, M Sa m an and A A Zozulya, PhysRev. A 56, R1713 (1997).
- [3] JW R. Jabosa and D. V. Petrov, PhysRev Lett. 83, 4967 (1999).
- [4] JLeach M JP adgett, S M B amett, S Franke-A mold and JC ourtial, Phys. Rev. Lett. 88, 257901 (2002).
- [5] M Fleischhauer, A Im am oglu and JP M arangos, Rev M od Phys. 77, 63 (2005).
- [6] A B M atsko, Y V R ostovtsev, M F leischhauer and M O Scully.PhysRevLett.86, 2006 (2001).
- [7] M F Andersen, C Ryu, P C lade, V Natarajan, A Vaziri, K Helmerson and W . D . Phillips, Phys. Rev. Lett. 97, 170406 (2006).
- [8] J.W R.Tabosa and A.Lezam a, J.P.hys B.40, 2809 (2007).
- [9] A Yu O kulov, JP hysB. 41, 101001 (2008).
- [10] A Yu O kulov, JETP Letters 88, pp. 487-491 (2008).
- [11] J-L.Thom as and R M archiano, Phys. Rev. Lett. 91, 244302 (2003).
- [12] N B Baranova, B Y Zeldovich, A V M am aev, N F P ilipetsky and V V Shkunov, JETP, 83, 1702 (1982).
- [13] B.Y. Zeldovich, N.F. Pilipetsky and V.V. Shkunov, "Principles of Phase Conjugation", (Springer-Verlag) (1985).

- [14] N G Basov, IG Zubarev, A B M ironov, S IM ikhailov and A Y O kulov, JETP 52, 847 (1980).
- [15] R W Boyd, K R zazew skiand P N anum, PhysRev.A 42, 5514 (1990).
- [16] A Yu O kulov, JM od O pt. 38, n.10, 1887 (1991).
- [17] A.Yu, Okulov, JOSA, B7, 1045 (1990).
- [18] M K lawunn and L Santos, New JPhys. 11, 055012 (2009).
- [19] L P P itaevskii, JE T P 13, 451 (1961).
- [20] Y S K ivshar and G P A grawal, "Optical Solitons: from Fibers to Photonic Crystals,", (A cademic Press, San Diego) (2003).
- [21] A.YuOkulov, JM od Opt. 55, n 2, 241 (2008).
- [22] M.S.Bigelow, N.N.Lepeshkin and R.W.Boyd, PhysRevLett. 90, 113903 (2003).
- [23] B D C lader and J H Eberly, JO SA, B 24, 916 (2007).
- [24] A Yu O kulov and A N O raevskii, Sov J Q uantum E lectron. 18, n 2, 233 (1988).
- [25] JE Sipe, PhysRev.A 52, 1875 (1995).
- [26] E M Lifshitz, L P Pitaevskii, and V B Berestetskii, "Quantum Electrodynamics", (Landau and Lifshitz Course of Theoretical Physics, Butterworth-Heineman, Oxford) Vol4 Ch.I x 6,8 (1982).