## Decay of Rabioscillations by dipolar-coupled dynam ical spin environments

V.V.Dobrovitski, A.E.Feiguin,  $^{2,3}$  R.Hanson, and D.D.Awschalom<sup>5</sup>

<sup>1</sup>Am es Laboratory, Iowa State University, Am es IA 50011, USA <sup>2</sup>Condensed Matter Theory Center, Department of Physics,

University of Maryland, College Park, MD 20742

<sup>3</sup>M icrosoft Station Q, University of California, Santa Barbara, CA 93106.

<sup>4</sup>K avli Institute of Nanoscience Delft, Delft University of Technology,

 $^5\mathrm{C}\,\mathrm{enter}$  for Spintronics and Quantum Computation,

University of California, Santa Barbara, California 93106, USA

(D ated: February 21, 2024)

We investigate the Rabi oscillations decay of a spin decohered by a spin bath whose internal dynam ics is caused by dipolar coupling between the bath spins. The decay form and rate as a function of the intra-bath coupling is studied analytically, and con rm ed num erically. The decay in general has neither exponential/G aussian or power-law form, and changes non-m onotonically with the intra-bath coupling, decelerating for both slow and fast baths. The form and rate of Rabi oscillations decay can be used to experim entally determ ine the intra-bath coupling strength for a broad class of solid-state system s.

PACS num bers: 76.30 D a, 03.65.Y z, 76.30.-v, 76.20.+ q

M easurem ent of the Rabioscillations decay is an important step in studying decoherence of quantum system s. For instance, extensive studies of Rabioscillations in superconducting qubits [1, 2, 3] greatly enhanced understanding of the decoherence caused by the bosonic baths and by the 1=f noise. The most promising directions for improvement were identied, and the decoherence time for superconducting qubits have been extended to microseconds range [4], placing them among most promising solid-state qubit candidates. Recently, much progress has been achieved in experimental im plem entation of long-living coherent Rabioscillations in various spin system s: m agnetic m olecules [5], NV im purity centers in diam ond [6, 7, 8], rare-earth ions in nonmagnetic host [9], quantum dots [10, 11, 12], to mention a few. Many of these systems are very attractive for basic studies of quantum spin coherence e ects, and also constitute prom ising candidates for quantum inform ation processing, coherent spintronics, or high-precision m aqnetom etry devices, provided that detailed understanding of the decoherence processes will be achieved.

A major decoherence source in these spin systems is the coupling of the central spin (e.g. the electron spin of the NV center in diam ond) to other spins present in the sample (environmental bath spins, e.g. the spins of nitrogen atoms in diam ond). Moreover, for many relevant spin systems, the direct coupling between the environmental spins is essential, producing internal dynamics of the bath. Here we explore theoretically the in uence of such a dynamical spin bath on the Rabioscillations of the central spin. A voiding the commonly used framework of generalized B both equations [13, 14, 15, 16], we are able to investigate the form and rate of decay as a function of the intra-bath coupling strength. We nd interesting behavior of the Rabi oscillations, which contradicts the expectations based on standard Red eld-type analysis: e.g., the slow bath leads to pronounced decay, while for the fast bath the decay rate vanishes but the Rabi frequency becom es renorm alized. We demonstrate how the form and rate of the Rabi oscillations decay can be used in experiment to characterize the intra-bath dynamics, and provide a rather simple recipe for analysis of experimental data.

Speci cally, we consider the situation of dilute dipolarcoupled bath: in a non-magnetic host crystal, a single centralspin of species S (e.g. belonging to a param agnetic in purity) is coupled to a bath of dilute spins of species I (e.g. belonging to other kind of param agnetic in purities or to nuclear spins). The coupling between the bath spins is non-negligible, and is caused by dipolar interactions. This situation encompasses a wide range of interesting solid-state spin systems, from Erions in CaW O4 studied in 1960s [17] to the NV centers in diam ond [6, 7, 8] which gained much attention very recently. M oreover, experim ental progress now makes possible detailed studies of Rabioscillations of an individual S spin [6, 8, 18]. We assume that a large magnetic eld  $B_z$  is applied along the z-axis (as in standard NMR/ESR settings), leading to Zeem an splittings  $!_S = {}_S B_z$  for the central spin S and  $!_{I} = {}_{I}B_{z}$  for the bath spins  $I_{k}$  ( $_{S}$  and  $_{I}$  are the gyrom agnetic ratios of the S and I spins, respectively). A lso, strong R abidriving eld H  $_{\rm R}$  is applied at the frequency  $!_{\rm S}$ . The di erence  $j!_{\rm S}$  $!_{I}$  j is much larger than any other energy scale, and after transform ation into rotating frame we obtain a standard secular H am iltonian for two dipolar-coupled spin species [13, 14]:

$$H = h_x S_x + A_k S_z I_k^z + C_{k1} (3I_k^z I_l^z I_k I_l)$$
(1)  

$$K = k_k S_z I_k^z + k_{k1} (3I_k^z I_l^z I_k I_l)$$
(1)

where  $S^{x;y;z}$  and  $I_k^{x;y;z}$  are the spin operators in the ro-

P.O. Box 5046, 2600 GA Delft, The Netherlands

tating frame, and  $h_x = H_R = 2$  is the rotating-frame R abidriving eld. The coupling constants  $A_k = \sum_{s \in I} [1 - 3(n_k^z)^2] = r_k^3$  are determined by the positions  $r_k$  of the bath spins  $I_k$  (k = 1; ::: N), where  $r_k = jr_k j$  and  $n_k = r_k = r_k$  (the origin of the coordinate frame coincides with the central spin). Sim ilarly, the intra-bath couplings  $C_{k1} = {}^2_I [1 - 3(r_{k1}^z)^2] = r_{k1}^3$  are determined by the vectors  $r_{k1} = r_k$  if. Note that the same H am iltonian (1) can be obtained without external eld  $B_z$  if the transition frequencies  $!_s$  and  $!_I$  are determined by the zero- eld splitting (e.g. due to anisotropic interactions). Sim ilarly, the assumption (used below) that S and  $I_k$  are spins 1/2 is not essential: for larger spins, we can consider each pair of levels as a pseudo-spin 1/2 [13, 14].

Initially the central spin is in the state "up", and the bath is in a maximally mixed state (unpolarized bath at high temperatures), i.e. the initial density matrix of the whole system is (0) = 2 <sup>N</sup> j "ih" j  $1_B$ , where  $1_B$  is the  $2^N$   $2^N$  identity matrix. This is appropriate for most experiments (for nuclear spins at temperatures above few nK, for electron spins | above tens of K). We calculate the time-dependent z-component of the central spin  $hS_z$  (t) = Tr (t) $S_z$ , where

(t) = exp ( iH t) (0) exp (iH t). In order to see wellpronounced long-living Rabioscillations, the driving eld should be large, so we assume that  $h_x$  is much larger than all other energy scales. We calculate the oscillations damping in the lowest order in  $1=h_x$ , treating the second (spin-bath coupling) and the third (bath internal H am iltonian) terms in Eq. 1 perturbatively, and excluding the bath internal H am iltonian  $H_B = \frac{1}{k_{fl}} B_{kl} (3I_k^z I_l^z I_k I_l)$ by the interaction representation transform ation. The resulting second-order H am iltonian

$$H_0 = h_x S_x + S_x B^2$$
 (t)=(2h<sub>x</sub>) (2)

where  $\hat{B}_{P}(t) = \exp(iH_{B}t)\hat{E}\exp(iH_{B}t)$ , and the operator  $\hat{B} = \int_{k}^{k} A_{k} I_{k}^{z}$ .

The evolution of  $\vec{B}$  (t) is complex, involving intricate correlations between bath spins. However, exact dynam ics of every single bath spin is not important, since hS<sub>z</sub> (t) i involves tracing over all bath spins. This is typical for m any spin-bath decoherence problem s, from m aqnetic resonance to quantum inform ation processing, and many approaches have been developed from the early days of NMR/ESR theory [19, 20] till very recently [21, 22, 23, 24, 25, 26, 27, 28]. All approaches involve a trade-o between quantitative rigour and qualitative understanding. Below, following the works [19, 20], we approximate the e ect of the bath by a random eld B (t), which is modeled as an O mstein-Uhlenbek process with the correlation function  $B (t) B (0) i = b^2 \exp \left( -R t \right)$  ,  $_{k}$  A  $_{k}^{2}$  , while the correlation where the dispersion b= decay rate R is determined by H  $_{\rm B}$  . This model m ay be oversimpli ed for complicated situations, e.g. the description of advanced control protocols requires m ore sophisticated treatm ent [24, 28]. However, our direct num erical simulations evidence that this model is quantitatively adequate for description of the Rabi oscillations decay, while providing a clear description of the physics underlying the Rabi oscillations decay, and allowing access to the regimes outside of the Bloch equations framework.

The H am iltonian (2) re ects a simple physical picture. The zero-order eigenstates of the H am iltonian (1) correspond to the gentral spin states j + i = 1 = 2[j"i + j#i] and j = 1 = 2[j"i - j#i], separated by a large R abi frequency  $h_x$ . Since  $h_x$  b and  $h_x$  R, the eld B (t) has no spectral components of noticeable m agnitude at the R abi frequency, and does not lead to transition between the states j + i and j i. The only relevant process is the pure dephasing, when the eld B destroys the initial phase relation between the states j + i and j i, leading to decay of  $hS_z$  (t) i, so that

 $hS_z$  (t)i = 1=2 hcos i = 1=2 Re hexp (i)i (3)

where h  $\,$  i denotes average over all possible realizations of B (t), and the phase

= 
$$h_x t + \frac{1}{2h_x} \int_{0}^{Z_t} B^2(s) ds = h_x t +$$
 (4)

is the total phase di erence between the states j + i and j i accumulated during time t, cf. the Ham iltonian l). The key quantity M (t) = hexp (i) i is an analytically computable G aussian path integral over the Omstein-Uhlenbeck process, which gives the answer

$$hS_{z} (t)i = 1=2 \operatorname{Re} M (t) \exp (ih_{x}t)]$$

$$M (t)]^{2} = \exp (Rt) [\cosh Pt + (R=P) \sinh Pt]$$

$$\frac{b^{2}}{h_{x}P} \exp (Rt) \sinh Pt; \quad (5)$$

where  $P = P \overline{R^2 2ib^2 R = h_x}$ .

A nalysis of R abi oscillations is often based on B lochtype equations [13] or its generalizations [15], derived for various systems from quantum optics [16] to solid state [15]. It is based on the Red eld-type approach [13, 14], taking into account only the term swhich are secular with respect to the Rabidriving h<sub>x</sub>S<sub>x</sub>. In addition to dephasing, these terms describe the actual transitions between the states *i* i and *j* i, which lead to a longitudinal relaxation (along the x-axis) of the central spin with the rate 1  $B^2 R = h_x^2$ . This rate is of second order in  $1 = h_x$ , and is determ ined by the spectral density of B (t) at the Rabi frequency  $h_x$ . Also, the generalized Bloch equations, having constant coe cients, always predict the decay to have a (multi)exponential form. In contrast, our results are not limited to the term s secular with respect to Rabidriving, and give the decay rate of the st order in 1=h<sub>x</sub>. The solution (5) predicts the decay which has, in general, no simple form (multiexponential, Gaussian, power-law, etc.) The fam iliar exponential decay occurs only at special values of b,  $h_x$ , and R, see below .

Thee ect of the bath internal dynam icsm ay be im portant even for slow baths, with R  $b^2 = h_x$ . In this lim it,  $2iR = h_x$ , see Eq. 5. At short times t P = b1=₽ j the bath behaves as static, and the Rabioscillations envelope exhibits non-exponential slow decay of the form  $1+\left(b^2t=h_x\right)^2$  )  $^{1=4}$  , in accordance with the exact results obtained earlier [10,25,26,27]. At long timest 1= إ j the decay has exponential form exp (  $bt R = 4h_x$ ), with the rate which decreases very slow ly with decreasing R. As a result, even for very slow bath the e ect of the bath dynam ics remains noticeable. In experiments, the above-described behavior of the Rabi oscillations decay can be detected by varying  $h_x$  (since the ratio of R to  $b^2 = h_x$  determ ines how fast/slow the bath is), and m akes it possible to estimate b and R .

A nother interesting feature of our results is that the decay of R abi oscillations changes non-m onotonously with R: it is fastest for R  $\dot{\mathcal{B}}=h_x$ , slowing down for both slow and fast baths. This is con rm ed by direct num erical simulations, see Fig. 1 and discussion below.

The regime of fast bath, with R  $b^2 = h_x$ , is even more interesting and unexpected. In this case

M (t) = exp 
$$i \frac{b^2 t}{2h_x} = \frac{b^4 t}{4h_x^2 R}$$
; (6)

and the R abi oscillations exhibit exponential decay with the rate  $_t = b^4 = (4h_x^2 R)$ , which vanishes quickly for large R. However, it does not mean that the e ect of the fast bath disappears. The bath still noticeably a ects the central spin, shifting its R abi frequency by  $b^2 = 2h_x$ , and only the decaying part of the bath e ect vanishes [29, 30]. This slowing of the R abi oscillations decay is con med by direct num erical simulations (Fig. 2).

On the other hand, for fast baths, the transitions between the states j+i and j i become important. W hile the dephasing between these states just shifts the Rabi frequency, the transitions lead to longitudinal relaxation along the x-axis with the rate 1 $b^2 R = h_v^2$ . This im – plies [13] exponential decay of R abi oscillations with the rate  $_1=2$ , which is comparable with the decay rate  $_t$ caused by the pure dephasing. Above, for sim plicity, we om itted the longitudinal relaxation (since it is of order  $1=h_{x}^{2}$ ), but we can include it using the Red eld-type approach: the answer (5) for  $S_z$  (t) just has to be multiplied by exp  $(_1t=2)$ . This explains how the system enters the generalized Bloch equation regime at R b: the dephasing e ect becom es negligible and the longitudinal relaxation becom es dom inant. In experim ent, this regim e can be identied by comparing l and t: they would be of the same order, changing as  $1=h_x^2$  for all su ciently large  $h_x$ .

To ensure that the physical picture above is adequate for realdipolar-coupled bath, we perform ed direct num erical simulations of the Rabioscillations decay. We place the central spin and N bath spins random ly, with uniform



FIG. 1: (Color online). Exact num erical simulations for N = 15 bath spins. (a) | Correlation function hB (0)B (t)i (norm alized to 1 at t = 0) obtained from direct simulations (black), see also [32]. Its tting (green) determ ines the bath parameters  $b_1$ ,  $b_2$ , and R for each value of the intra-bath coupling scale  $E_B$  (here  $E_B = 1$ ,  $b_1 = 0.62$ ,  $b_2 = 0.58$ , R = 0.095). (b) | Rabioscillations decay for  $h_x = 14.14$ , b = 0.85, and  $E_B = 1$ ; individual oscillations are not resolved. Num erical results (black) agree well with analytics (m agenta line, only the envelope shown). (c) Envelopes of simulated Rabidecay for  $E_B = 1$  (blue),  $E_B = 0:1$  (green), and  $E_B = 0$ (static bath, red); corresponding individual oscillations near t = 300 are shown on panel (d) by same colors. A nalytical results practically coincide with simulations, and are not shown. The decay rate changes nonmonotonically with  $E_B$ : the decay is slower for  $E_B = 1$  (blue), and  $E_B = 0$  (red) in comparison with  $E_{B} = 0:1$  (green). Att = 300, on panel (d), the oscillation amplitude for  $E_B = 0.1$  (green) is twice sm aller than that for  $E_B = 0$ .

density n = 1, inside a cube with the side  $(N + 1)^{1=3}$ , central spin being in the center of the cube. All interaction coe cients are calculated according to Eq. 1, using the actual coordinates of the spins. The value  $E_{SB} = S_{I}$ , which determ ines the strength of coupling between the central spin and the bath, is set to 1, thus de ning the energy and time scales for all quantities below. The value  $E_B = \frac{2}{1}$ , which governs the energy scale of intra-bath couplings, is varied, making the bath slower or faster. We simulate the dynamics of the system using two num ericalapproaches. For small N (e.g., N = 15 in Fig. 1) we exactly solve of the Schrodinger equation with the H am iltonian (1) via Chebyshev polynom ialexpansion [31]. The dipolar-coupled systems with N > 15 are dicult to m odel this way (due to exponentially increasing resources requirem ents), so we use the P-representation sam pling [31] for modeling larger baths (N = 159 in Fig. 2). To com pare num erical solutions with the analytics, we calculate the total coupling to a bath  $b = \begin{bmatrix} P & N \\ k = 1 \end{bmatrix} A_k^2 I^{1-2}$ directly from the positions of the spins (see Eq. 1 and below). The correlation decay rate R requires a separate



FIG.2: (Color online). Num erical simulations for N = 159bath spins. (a) Correlation function hB (0)B (t)i (norm alized to 1 at t = 0) obtained from P-representation simulations (black); its tting (green) determines the parameter R for each value of the intra-bath coupling scale  $E_{\,B}\,$  (here  $E_B = 1$  and R = 0.097). In spite of rather large statistical uctuations, the parameter R is determined precisely enough to be used in Eq. 5. (b) Rabioscillations decay for  $h_x = 15.0$ , b = 1.39, and  $E_B = 0.1$ ; individual oscillations are not resolved. Num erical results (black) agree well with analytics (m agenta line, only the envelope shown). (c) Envelopes of simulated Rabidecay for  $E_B = 0:1$  (blue),  $E_{B} = 1$  (green), and  $E_{B} = 10$  (red); the longitudinal decay for E<sub>B</sub> = 10 is taken into account. A nalytical results practically coincide with simulations, and are not shown. The decay rate decreases for faster baths: the decay is slowest for  $E_B = 10$  (red). Individual oscillations for  $E_B = 10$  near t = 20 are shown on panel (d) in red: their frequency is shifted by  $b^2=2h_x = 0.064$  from the value  $h_x = 15.0$  (oscillations with frequency  $h_x = 15.0$  are shown by dotted black line to dem onstrate the phase di erence accumulated since t = 0).

simulation: we calculate hB (0)B (t)i and nd R by tting it to a decaying exponent, see also [32]. Varying the system parameters in a wide range, we simulated baths with N = 15, 59, and 159 spins, and found good agreement between num errics and analytics; typical results are given in Figs. 1,2.

The resulting experimental recipe is rather simple. If the decay has a power-law form at shorter times, changing to exponent later, with the decay constants changing as  $1=h_x$  and the duration of the two regimes varying with  $h_x$ , then the bath is slow. If the R abi oscillations decay is exponential, with the rate changing as  $1=h_x^2$  and the frequency shift varying as  $1=h_x$ , then the bath is fast. The fast bath can be m ade slow by strong increase in  $h_x$ .

Sum m arizing, we studied the decay of the Rabi oscillations of the central spin interacting with a dipolarcoupled dynam ical spin bath. A pproxim ating the e ect of the bath as a random eld (O mstein-U hlenbeck process), we nd analytically the form of the decay. Validity of the approximation is conmed by direct numerical simulations. The oscillations decay has interesting features, such as non-monotonic variation of the decay rate with increasing the intra-bath coupling, and slowing down of the decay for fast baths. Studying the behavior of the Rabi oscillations may help in experimental characterization of the dynamical spin bath, and is wellwithin experimental reach e.g. for NV centers in diamond. W ork at Ames Laboratory was supported by the Department of Energy | Basic Sciences under Contract No. DE-AC 02-07C H 11358. We acknow ledge support from AFOSR (DDA.), FOM and NWO (RH.). A E F. acknow ledges support from the M icrosoft Corporation.

- [1] E. Paladino et al, Phys. Rev. Lett. 88, 228304 (2002)
- [2] Y.M.Galperin et al, Phys. Rev. Lett. 96, 097009 (2006)
- [3] J.C larke and F.K.W ilhelm, Nature 453, 1031 (2008)
- [4] G. Ithier et al, Phys. Rev. B 72, 134519 (2005); A.A. Houck et al, Phys. Rev. Lett. 101, 080502 (2008).
- [5] S.Bertaina et al., Nature 453, 203 (2008)
- [6] T.Gaebelet al, Nature Physics 2, 408 (2006).
- [7] L.Childress et al., Science 314, 281 (2006).
- [8] R. Hanson et al., Science 320, 352 (2008).
- [9] S.Bertaina et al., Nature Nano. 2, 39 (2007)
- [10] F. H. L. K oppens et al, Phys. Rev. Lett. 99, 106803 (2007)
- [11] J.R. Petta et al, Science 309, 2180 (2005)
- [12] R.Hanson et al, Rev.M od.Phys.79, 1217 (2007).
- [13] C. P. Slichter, Principles of Magnetic Resonance (Springer, Berlin, New York, 1990).
- [14] A. Abragam, Principles of Nuclear M agnetism (Oxford University Press, Oxford, New York, 1961).
- [15] E.Geva et al., J.Chem. Phys. 102, 8541 (1995).
- [16] C.Cohen-Tannoudjiand S.Reynaud, J.Phys.B 10, 365 (1977).
- [17] W .B.M im s, Phys. Rev. 168, 370 (1968)
- [18] H.J.M am in et al, Phys.Rev.Lett. 91, 207604 (2003).
- [19] P.W. Anderson and P.R.W eiss, Rev.M od. Phys. 25, 269 (1953); J.R.K lauder and P.W. Anderson, Phys. Rev. 125, 912 (1962).
- [20] R. Kubo, J. Phys. Soc. Japan 17, 1100 (1962); R. Kubo, M. Toda, and N. Hashitsum e, Statistical Physics II (Springer, Berlin, New York, 1998).
- [21] A.Garg, Phys. Rev. Lett. 74, 1458 (1995).
- [22] V. V. D obrovitski et al, Phys. Rev. Lett. 90, 210401 (2003).
- [23] W . A. Coish and D. Loss, Phys. Rev. B 72, 125337 (2005).
- [24] R.de Sousa and S.Das Sam a, Phys.Rev.B 68, 115322 (2003); W.M.W itzel and S.Das Sam a, Phys.Rev.B 74, 035322 (2006).
- [25] V.V.Dobrovitski et al, quant-ph/0112053; V.V.Dobrovitski et al, HAIT J.Sci.Eng. 1, 586 (2005).
- [26] F.M.Cucchiettietal, Phys.Rev.A 72, 052113 (2005).
- [27] J.M. Taylor and M.D. Lukin, Quant. Info. Proc. 5, 503 (2006).
- [28] W . Yao et al, Phys. Rev. Lett. 98, 077602 (2007); S.

Saikin et al, Phys. Rev. B 75, 125314 (2007).

- [29] This is in clear contrast with the motional narrowing phenomenon, when the free oscillations decay rate vanishes for fast bath [13, 14, 19, 20]. In the motional narrowing case, extremely fast bath does not a left the central spin at all.
- [30] This result can be obtained rather universally, for any non-pathological bath, starting from Eq.2 and assuming that the correlations hB (0)B (t)idecay to zero with a rate much faster than  $b^2 = h_x$ .
- [31] W . Zhang et al, J. Phys. Cond. M atter 19, 083202

(2007). P-representation sampling, although not exact, is expected to give accurate results for our problem at large N .

[32] For N = 15, due to nite-N e ects, the correlations decay to a nite value, so that the eld B (t), in addition to the O mstein-U hlepbeck component, has a static component (of order of  $1^{\pm}$  N), and the correlation function has the form hB (0)B (t)i =  $b_1^2 \exp((Rt) + b_2^2)$ , with restriction  $b_1^2 + b_2^2 = b^2$ . Simulations still agree with analytics very well.