Supernova N eutrinos D etection O n E arth*

GUO X in-Heng^{1;1)} HUANG M ing-Yang^{1;2)} YOUNG B ing-L in^{2;3;3)}

(College of Nuclear Science and Technology, Beijing Norm al University, Beijing 100875, China)
 (Department of Physics and Astronomy, Iowa State University, Ames, Iowa 5001, USA)
 (Technology, Chinas, Chinas,

3 (Institute of Theoretical Physics, Chinese A cadem y of Sciences, Beijing, China)

A bstract In this paper, we rst discuss the detection of supernova neutrino on Earth. Then we propose a possible method to acquire inform ation about $_{13}$ sm aller than 1.5 by detecting the ratio of the event num bers of di erent avor supernova neutrinos. Such an sensitivity cannot yet be achieved by the D aya B ay reactor neutrino experiment.

K ey words supernova, neutrino, collective e ects, M SW e ects, Earth m atter e ects, D aya B ay

PACS 14.60 Pq, 13.15.+ g, 25.30 Pt

1 Introduction

The supernova (SN) explosion is one of the most spectacular cosm ic events and a source of new physics $ideas^{[1, 2, 28, 29, 30]}$. Observable e ects of SN neutrinos in underground detectors have been a subject of intense investigation in astroparticle physics, both on general grounds and in relation to the SN event like 1987A. In particular, avor oscillation in the SN may shed light on the problem of neutrino masses and mixing by means of the associated matter e ects. Several neutrino laboratories, including the Daya Bay reactor neutrino underground laboratory^[3] which is under construction, can be used to detect possible neutrino events from an SN explosion and serve as the SN Earth Warning System. Hence theoretical prediction for the detection of SN neutrinos in the D aya Bay and other neutrino experiments is very desirable.

In the realistic case, when we detect neutrinos from a type II SN explosion at Daya Bay, there are three e ects (the collective e ects arising from neutrino-neutrino interactions^[4, 5, 6, 7, 8, 9, 10, 11, 12], the well-known Mikheyev-Sm imov-W olfenstein (MSW) e ects^[13, 14, 15, 16], and Earth matter e ects^[17, 18, 19]) need to be considered. Using the Landau-Zener form ula^[20, 21], the expression of the crossing probability, $P_{\rm H}$, which is the neutrino jump probability from mass eigenstate 1 to 3 at the high resonance region inside the SN, can be calculated^[22, 23, 24]. W ith the relation between P_H and the mixing angle ₁₃, we can predict the SN neutrino event numbers N as a function of ₁₃. Therefore, we can propose a possible method to acquire information about sm all ₁₃ through the detection of SN neutrinos^[19].

2 Detection of SN neutrinos on Earth

SN are extremely powerful explosions in the universe which term inate the life of som e stars^[1, 25, 26, 27]. They make the catastrophic end of stars more massive than 8 solarm asses leaving behind com pact rem nants such as neutron stars or black holes which m ay be observed. For historical reasons, SN are divided into two wide categories (type I and type II) characterized by the absence or presence of hydrogen lines. How ever, the most im portant physical characteristics is the mechanism that produces the SN, which distinguishes SN of type Ia from SN of type Ib, Ic, and II. This di erence becom es noticeable in the light spectrum some months after maximum lum inosity, when the ejecta become optically thin and the innerm ost regions become visible: the spectrum of SN Ia is dom inated by Fe emission lines, while SN Ib, Ic, and $\rm II$ show O and C em ission lines. From the point of view of neutrino physics, type 1b, Ic, and II SN are much m ore im portant than type Ia SN, sim ply because they

supported in part by N ational N atural Science Foundation of C hina (10535050, 10675022), the K ey P roject of C hinese M inistry of E ducation (106024) and the Special G rants from B eijing N orm al U niversity.

¹⁾ E -m ail:xhguo@ bnu.edu.cn

²⁾ E -m ail:hm y19151905@ m ailbnu.edu.cn

³⁾ E -m ail:young@ iastate.edu

produce a huge ux of neutrinos and antineutrinos of all avors.

The type II SN is thought to be generated by the core collapse of red (or blue as SN 1987A) giant star with a mass between about 8-9 and 40-60 solar masses. The total energy release (about 3 10^{53}) is approximately the gravitational binding energy of the core. It generates intensive neutrinos which take away about 99% of this total energy. The explosion itself consumes about 1% of this total energy. The vast amount of neutrinos are produced in two bursts. In the rst burst which lasts for only a few milliseconds, electron neutrinos are generated via the electron capture by nuclei $e + N (Z;A)! N (Z 1;A) + _{e}$ and the inverse betadecay $e + p! n + \dots$. In the second burst which lasts longer, neutrinos of all avors (and with being e, ,) are produced through the electron-positron + , electron-nucleon pair annihilation $e + e^+$! brem sstrahlung e + N ! e + N + + , nucleonnucleon brem sstrahlung N + N ! N + N + + ,plasm on decay ! + , and photoannihilation +e ! e + + [25, 26, 27]. When SN neutrinos of a de nite avorare produced they are approximately in an e ective m ass eigenstate due to the extrem ely high m atter density environm ent. W hile they propagate outward to the surface of the SN they could experience collective e ects^[4, 5, 6, 7, 8, 9, 10, 11, 12] and M SW e ects^[13, 14, 15, 16]. A fler travelling the cosm ic distance to reach Earth, the arriving neutrinos are mass eigenstates, which then oscillate in avors while going through Earth m atter. Therefore, we also have to consider Earth matter e ects^[17, 18, 19] when we com pute the event num bers of the various avors of neutrions.

LetP represent the collective e ects of neutrinoneutrino interactions which is a stepwise avor conversion probability of neutrinos at a critical energy $E_{\rm c}$. In order to obtain a simple expression for P for the neutrino and P for the antineutrino, we take a constant matter density and box-spectra for both the neutrino and antineutrino^[8, 9, 10]. An analysis of the collective e ects in the case of three avors has been m ade in R ef. [11], and it allow s us to characterize the collective oscillation e ects and to write down the avor spectra of the neutrino and antineutrino arriving at Earth. Following Ref. [11], we have P = P = 1for the norm all hierarchy; and P = 1, while (

$$P = \begin{cases} 1 & (E < E_{c}); \\ 0 & (E > E_{c}); \end{cases}$$
(1)

for the inverted hierarchy, where E $_{\rm C}$ = 7M eV $^{\rm [11,\,12]}$.

There are two M SW resonance regions: the high resonance region and the low resonance region. Let us denote the probability that the neutrinos jump from one m ass eigenstate to another at the high (low) resonance layer by $P_{\rm H}$ ($P_{\rm L}$). U sing the Landau-Zener form ula^[20, 21], one can obtain that

$$P_{H} = \frac{\exp(\frac{1}{2}F) \exp[\frac{1}{2}(\frac{F}{\sin^{2}13})]}{1 \exp[\frac{1}{2}(\frac{F}{\sin^{2}13})]}; \quad (2)$$
$$= \frac{jm_{13}^{2}j\sin^{2}2_{13}}{2E} \frac{\sin^{2}2_{13}}{\cos^{2}13} \frac{1}{jd\ln N_{e} = dr_{Hes}};$$

where N_e is the electron density and F can be calculated by Landau's method^[22]. Using the SN matter density pro le C $(10 \text{ cm} = \text{r})^3$ 10⁹ g=cm³ where C is a structure constant between 1 and 15^[28, 29, 30] and F 1 (sm all ₁₃), we can obtain an simple expression of P_H that^[19, 22, 23, 24]

$$P_{H} = \exp \left(\frac{10^{10} \text{M eV}}{12} \frac{\sin^{3} 2_{13}}{\cos^{2} 2_{13}} + \frac{\sin^{2} 2_{23}}{\cos^{2} 2_{13}} + \frac{j \text{m}^{2} 2_{23}}{1 \text{eV}^{2}} \right)$$
(3)

where j m $^2_{32}$ j= 2:6 10 3 eV 2 . Similarity, we can calculate the expression of the crossing probability at the low resonance region inside the SN, P_L. How ever, due to the large angle solution of the neutrino mixing, P_L 0.



Fig. 1. Illustration of the path of the SN neutrino reaching the detector underground in Earth. D is the location of the detector, is the incident angle of the neutrino, O is the center of Earth, L is the distance the neutrino travels through Earth, and x is the distance of the neutrino to the center of Earth.

Suppose a neutrino reaches the detector at the incident angle (see Fig. 1). Then the distance that neutrino travels through Earth is

$$L = (R_{E} + h)\cos + \frac{R_{E}^{2}}{R_{E}^{2}} (R_{E} - h)^{2}\sin^{2}; \quad (4)$$

Т

where h (0.4km for the D aya B ay experiment) is the depth of the detector in and $R_E = 6400$ km is the radius of E arth. Let x be the distance that the neutrino travels into E arth, then the distance of the neutrino to the center of E arth, x, is given by

$$x = \frac{P}{(R_E + h)^2 + (L_x)^2 + 2(R_E - h)(L_x)\cos}$$
:

Let P_{ie} be the probability that a neutrino m ass eigenstate $_i$ enters the surface of E arth and arrives at the detector as an electron neutrino $_e$, one obtains^[18, 19]

$$P_{2e} = \sin^2_{12} + \frac{1}{2} \sin^2 2_{12} \int_{x_0}^{x_{\rm f}} dx V(x) \sin_{x!x_{\rm f}}^{m}; (5)$$

where $_{12} = 32.5$, the potential V (x) that the electron neutrino experiences in Earth is $\frac{p}{2G_F}$ (x)=(m_p+m_n) and $\frac{m}{a!b}$ is dened as

$$\begin{array}{rcl} & & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & \\ & & & \\ & & & & \\$$

where " $(x) = 2EV(x)=4m_{21}^2$ and (x) is the matter mass density inside Earth^[31].

In the following, we calculate the event numbers N (i) of SN neutrinos that can be observed through various reaction channels "i". This is done by integrating over the neutrino energy E of the product of the target number N_T, the cross section of each channel (i), and the neutrino ux function at the detector F^{D} (E)=4 D²,

N (i) = N_T dE (i)
$$\frac{1}{4 D^2}$$
 F³; (6)

where stands for the neutrino or antineutrino of a given avor, and D (10 kpc in the present discussion) is the distance between the SN and Earth. A fler a straightforward calculation, the uxes at the detector can be obtained^[11, 12] (x = i):

$$F_{e}^{D(N)} = P_{2e}P_{H}F_{e}^{(0)} + (1 P_{2e}P_{H})F_{x}^{(0)};$$

$$F_{e}^{D(N)} = (1 P_{2e})F_{e}^{(0)} + P_{2e}F_{x}^{(0)};$$

$$2F_{x}^{D(N)} = (1 P_{2e}P_{H})F_{e}^{(0)} + (1 + P_{2e}P_{H})F_{x}^{(0)};$$

$$2F_{x}^{D(N)} = P_{2e}F_{e}^{(0)} + (2 P_{2e})F_{x}^{(0)};$$

$$(7)$$

for the norm all ierarchy (4 m $_{31}^2 > 0$), and

$$F_{e}^{D(I)} = \begin{pmatrix} P_{2e}F_{e}^{(0)} + (1 P_{2e})F_{x}^{(0)}; & (E < E_{C}) \\ F_{e}^{(0)}; & (E > E_{C}) \end{pmatrix}$$

$$F_{e}^{D(I)} = P_{H} (1 P_{2e})F_{e}^{(0)} + (1 + P_{2e}P_{H} P_{H})F_{x}^{(0)}; \\ (1 P_{2e})F_{e}^{(0)} + (1 + P_{2e})F_{x}^{(0)}; & (E < E_{C}) \end{pmatrix}$$

$$F_{e}^{(0)} + F_{x}^{(0)}; & (E > E_{C}) \end{pmatrix}$$

$$F_{e}^{(0)} + F_{x}^{(0)}; & (E > E_{C}) \end{pmatrix}$$

$$F_{e}^{(1)} = (1 + P_{2e}P_{H} P_{H})F_{e}^{(0)} + (1 + P_{2e}P_{H} P_{H})F_{e}^{(0)}; \qquad (1 + P_{2e}P$$

for the inverted hierarchy (4 m $_{31}^2$ < 0). In Eqs. (7) and (8), F ⁽⁰⁾ is the tim e-integrated neutrino energy spectrum of avor in vacuum which can be described by the Ferm i-D irac distribution

$$F^{(0)}(E) = \frac{L^{(0)}}{F_{3}T^{4}} \frac{E^{2}}{\exp(E = T) + 1}; \quad (9)$$

where T $\,$ is the tem perature of the neutrino $^{\scriptscriptstyle [32,\ 33]}$

$$_{e} = 3$$
 4M eV; $T_{e} = 5$ 6M eV;
 $T_{x} = 7$ 9M eV; (10)

is the pinching param eter of the spectra to represent the deviation from being exactly therm $al^{32, 33]}$

3 5; 2:0 2:5; 0 2; (11)

 $L^{(0)}$ is the the lum inosity, and F_{j} is dened by

$$F_{j} = \int_{0}^{Z_{1}} \frac{x^{j}}{\exp(x) + 1} dx:$$

In the next section, using the relation between the event number of SN neutrinos detected at D aya B ay, N , and the m ixing angle $_{13}$, we will propose a possible m ethod to acquire inform ation about $_{13}$ sm aller than 1:5 .

3 A cquire information about 13 sm aller than 1:5 at D aya B ay

It can be seen from the above section that, using Eqs. (1), (3), (5), (6), (7), and (8), we can obtain the relation between the event number of di erent avor SN neutrinos, N, and the mixing angle $_{13}$. Let R be the ratio of the event number of $_{e}$ over that of $_{e}$, one can obtain R as a function of $_{13}$. Therefore, we can propose a possible method to acquire some e inform ation about the mixing angle $_{13}$ by detecting SN neutrinos. In this section, we will just consider the process of the neutrino-carbon reactions, since our method is not suitable for the inverse beta-decay and the neutrino-electron reactions in the D aya B ay experiment. The details are discussed at length in R ef. [19].

The Daya Bay Collaboration uses LAB as the main part of the liquid scintillator and the total detector mass is about 300 tons. LAB has a chem ical composition including C and H. In our calculation, the ratio of the numbers of C and H, $N_c = N_H$, is about 0.6. Therefore, the total numbers of target

protons N_T^(C) = 1:32 10^{31} . For the neutrino-neutrino reactions, the elective cross sections are as follow s^[34]:

h
$$({}^{12}C (_{e};e) {}^{12}N)i= 1.85 \quad 10^{43} \text{ cm}^{2};$$

h $({}^{12}C (_{e};e^{+}) {}^{12}B)i= 1.87 \quad 10^{42} \text{ cm}^{2};$ (12)

for the charged-current capture, and

h
$$\binom{12}{e}$$
C)i=1:33 10⁴³ cm²;
h $\binom{12}{e}$ C)i=6:88 10⁴³ cm²; (13)
h $\binom{x}{x}\binom{x}{x}^{12}$ C)i=3:73 10⁴² cm²; x=;;

for the neutral-current capture. Using Eqs. (1), (3), (5), (6), (7), (8), (12) and (13), we plot the ratio R of the event number of $_{\rm e}$ over that of $_{\rm e}$ which can be detected at D aya B ay as a function of the m ixing angle $_{13}$. The result is shown in Fig. 2.



Fig. 2. The ratio of the event number of e to that of e, R, as a function of the mixing angle ¹³ in the channel of neutrino-carbon reactions at the D aya B ay experiment. The incident angle is (a) = 30; (b) = 90; (c) = 93; (d) = 150. The solid curves correspond to the norm alhierarchy (m ax), the dashed curves correspond to the inverted hierarchy (m ax), the dot-dashed curves correspond to the norm al hierarchy (m in), the dotted curves correspond to the inverted hierarchy (m in), where "m ax" ("m in") corresponds to the maximum (m inim um) values of T and .

References

- 1 K otake K, Sato K, Takahashi K, R ept. P rog. P hys., 2006, 69:971 | 1143
- 2 Dighe A S, Physics potential of future supernova neutrino observation, arX iv hep-ph/0809.2977, Neutrinos from a core collapse supernova, arX iv hep-ph/0712.4386
- 3 [Daya Bay] GUO X H et al., A precision m easurement of

It can be seen from Fig. 2 that the uncertainties of R due to T and are not large. For $_{13}$ 1.5, R is very sensitive to $_{13}$. However, while $_{13} > 1.5$, R is nearly independent of $_{13}$. Therefore, when $_{13}$ is smaller than 1.5, we may restrict the mixing angle $_{13}$ in a small range and get information about mass hierarchy by detecting the ratio of event num – bers of SN neutrinos even though there are still som e uncertainties due to the incident angle , the mass hierarchy 4 m $_{31}^2$, and the structure coe cient C of the SN density function.

At the D aya B ay experiment, the sensitivity of $\sin^2 2_{13}$ will reach 0.01, i.e., to determ ine $_{13}$ down to about 3. Therefore, if the actual value of $_{13}$ is smaller than 3, the D aya B ay experiment can only provide an upper limit for $_{13}$. However, if an SN explosion takes place during the operation of D aya B ay, roughly within the cosm ic distance considered here, it is possible to reach a much smaller value of $_{13}$ through the ratio of the event num bers of di erent avor SN neutrinos in the channel of neutrinocarbon reactions as discussed above. It is interesting to note that because of the multi detectors set up, experiments such as the D aya B ay have an internal coincidence check for SN neutrino events.

4 Summary and discussion

In this paper, we rst discuss the detection of SN neutrinos on Earth. Since neutrino avor conversions inside the SN depend on the neutrino mixing angle $_{13}$, we give a possible method to acquire information about $_{13}$ sm aller than 1.5 by detecting SN neutrinos at D aya B ay.

We let the parameters in the neutrino energy spectra (the temperatures and the pinching parameters) vary in some reasonable ranges. In fact, the simulations from the two leading groups, the Livermore group^[35] and the Garching group^[36, 37], led to parameters which agree within about 20-30%. How ever, their central values of SN parameters are dierent.

the neutrino m ixing angle 13 using reactor antineutrinos at D aya B ay, arX iv hep-ex/0701029

- 4 DUAN H Y, Fuller G M, QIAN Y Z, Phys. Rev. D, 2006, 74:123004
- 5 DUAN H Y, Fuller G M, Carlson J et al. Phys. Rev. D, 2006, 74: 105014
- 6 DUAN H Y, Fuller G M, Carlson J et al. Phys. Rev. D, 2007, 75: 125005

- 7 DUAN H Y, Fuller G M, Carlson J, Comp. Scie. & Disc., 2008, 1:015007
- 8 Hannestad S,Ra elt G G,SiglG et al.Phys.Rev.D,2006, 74:105010
- 9 Ra elt G G, Sm imov A Y, Phys. Rev. D, 2007, 76:081301
- 10 Ra elt G G, Sm imov A Y, Phys. Rev. D, 2007, 76: 125008
- 11 Dusgupta B, Dighe A, Phys. Rev. D, 2008, 77: 113002
- 12 Chakraborty S, Choubey S, Dasgupta B et al. J. Cosm ol. A stropart. Phys., 2008, 09:013
- 13 Wolfenstein L, Phys. Rev. D, 1978, 17: 2369 2374
- 14 W olfenstein L, Phys. Rev. D, 1979, 20: 2634 2375
- 15 M ikheyev S P, Sm imov A Y, Sov. J. Nucl. Phys., 1985, 42: 913 | 917
- 16 M ikheyev S P, Sm imov A Y, Nuovo C in ento C, 1986, 9: 17 | 26
- 17 Dighe A S, Sm imov A Y, Phys. Rev. D, 2000, 62:033007
- 18 GUO X H, YOUNG B L, Phys. Rev. D, 2006, 73: 093003
- 19 GUO X H, HUANG M Y, YOUNG B L, Realistic Earth m atter e ects and a method to measure sm all ₁₃ in the detection of supernova neutrinos, arX iv:0806.2720 [hep-ph]
- 20 Zener C, Proc. R. Soc. London A, 1932, 137:696 702
- 21 Landau L D, Lifshitz E M.Quantum Mechanics: Nonrelativistic Theory.New York: Pergamon, 1977.199 237
- 22 Kuo T K, Pantaleone J, Rev. M od. Phys., 1989, 61: 937 979
- 23 Lunardini C, Sm imov A Y, J. Cosm ol. A stropart. Phys., 2003, 06:009

- 24 Kachelrie M, Strum ia A, Tom as R et al. Phys. Rev. D, 2002, 65: 073016
- 25 G iuntiC, K im C W .Fundam entals of N eutrino P hysics and A strophysics. New York: Oxford, 2007.511 539
- 26 Shapiro S L, Teukolsky S A.B lack Holes, W hite Dwarfs, and Neutron Stars.New York: W iley, 1983.512 542
- 27 Mohapatra R N, PalP B.Massive Neutrino in Physics and Astrophysics, 3nd ed. Singapore: World Scientic, 2004. 340 358
- 28 Brown G E, Bethe H A, Baym G, Nucl. Phys. A, 1982, 375:481 532
- 29 Bethe H A, Rev. M od. Phys., 1990, 62:801 866
- 30 Janka H T, Astron. Astrophys., 2001, 368: 527 560
- 31 D ziewonski A M, Anderson D L, Phys. Earth Planetary Interiors, 1981, 25: 297 | 356
- 32 Janka H T, H illebrandt W , A stron. A strophys., 1989, 224: 49| 56
- 33 Janka H T, Hillebrandt W, Astron. Astrophys. Suppl. Ser., 1989, 78: 375 397
- 34 Cadonati L, Calaprice F P, Chen M C, A stropart. Phys., 2002 16, 361–372
- 35 TotaniT, Sato K, Dalhed H E et al. A strophys. J., 1998, 496:216 | 225
- 36 KeilM T, Ra elt G G, Janka H T, Astrophys. J., 2003, 590:971 | 991
- 37 K eilM T, Supernova neutrino spectra and applications to avor oscillations, arX iv astro-ph/0308228