# R adiative Transfer M odeling of the W inds & C ircum stellar Environm ents of H ot A nd C ool M assive Stars

A.Lobel

RoyalObservatory of Belgium, Ringlaan 3, B-1180 Brussels, Belgium

A bstract. We present modeling research work of the winds and circum stellar environments of a variety of prototypical hot and cool massive stars using advanced radiative transfer calculations. This research aims at unraveling the detailed physics of various mass-loss mechanisms of lum inous stars in the upper portion of the H-R diagram. Very recent 3-D radiative transfer calculations, com bined with hydrodynamic simulations, show that radiatively-driven winds of OB supergiants are structured due to large-scale density- and velocity- elds caused by rotating bright spots at the stellar equator. The mass-loss rates com puted from m atching D iscrete Absorption C om ponents (DAC s) in IUE observations of HD 64760 (B Ib) do not reveal appreciable changes from the rates of unstructured (sm ooth) wind models.

Interm ediate yellow supergiants (such as the yellow hypergiant C as, F G Ia0), on the other hand, show prom inent spectroscopic signatures of strongly increased m ass-loss rates during episodic outbursts that cause dram atic changes of the stellar photospheric conditions. Long-term high-resolution spectroscopic m on itoring of cool hypergiants near the Yellow E volutionary Void reveals that their m ass-loss rates and w ind-structure are dom inated by photospheric enuptions and large-am plitude pulsations that im part m echanicalm om entum to the circum stellar environm ent by propagating acoustic (shock) waves.

In massive red supergiants, how ever, clear evidence for mechanical wave propagation from the sub-photospheric convection zones is lacking, despite their frequently observed spectroscopic and photom etric variability. Recent spatially resolved HST-STIS observations inside Betelgeuse's (M Iab) very extended chrom osphere and dust envelope show evidence of warm chrom ospheric gas far beyond the dust condensation radius of radiative transfer models. M odels for these long-term spectroscopic observations demonstrate that the chrom ospheric pulsations are not spherically symmetric. The STIS observations point to the importance of mechanical wave propagation for heating and sustaining chrom ospheric conditions in the extended winds of red supergiants.

#### 1. Introduction

The winds and circum stellar environments of massive stars are unique laboratories for studying the detailed physics of stellar mass-loss mechanisms. In many lum inous hot and cool supergiants large spectroscopic variability is observed on time-scales of weeks to years. Long-term spectroscopic monitoring of massivestar variability over the past three decades therefore has provided a wealth of information about the driving physics and active structuring mechanisms in these stars with very extended winds. This paper provides a brief overview of important research results for a hot, intermediate, and cool supergiant obtained with advanced radiative transferm odeling of the spectroscopic variability.



Figure 1. Panel left: 3-D non-LTE radiative transfer t with W ind3D to the DACs in Si iv 1395 compared to IUE observations of HD 64760 (m iddle panel). Panel right: Density contrast of the hydrodynam ic input m odel computed with Zeus3D. Two unequally bright spots produce large-scale w ind-velocity structures and density enhancements. The size of the over-plotted arrows indicates the magnitude of the velocity deceleration with respect to the sm ooth unperturbed wind. The bright spots cause rotating density waves in the equatorial wind of the hot supergiant (see text).

## 2. 3-D M odeling the W ind of H ot Supergiant H D 64760 (B Ib)

In recent years it has become clear that mass-loss rates of hot massive stars are considerably overestimated because the clumping of their winds has not been taken into account. Understanding the physical nature of these wind clumping processes, determining the amount of wind clumping, and revising the total stellar mass-loss rates are therefore hot topics in current massive star research. Detailed information about the wind structure can be obtained with accurate numerical models. We develop a parallel computer code W ind 3D which solves the 3-D transport of radiation in the scattering winds of massive stars. It calculates the detailed shape of resonance lines that form in very extended winds (e.g., P Cygnipro les) in non-LTE.

We apply W ind 3D for the detailed modeling of DACs observed in the Siiv resonance lines of HD 64760 (B0.5 lb), a key massive hot star. The study provides new evidence that the DACs are in fact large-scale spiraling densityand velocity-structures in the equatorial wind. These wind structures are due to rotating hotspots at the stellar surface. The bright spots do not co-rotate with the stellar surface, but lag ve times behind the fast surface rotation. A hydrodynam icm odel with two spots of unequal brightness and size on opposite sides of the equator, with opening angles of 20 5 and 30 5 diam eter, and that are 20 5% and 8 5% brighter than the stellar surface, respectively, provides the best t to the observed DACs (Fig. 1). The mass-bass rate of the structured wind model does not exceed the rate of the spherically symmetric sm ooth wind m odel by more than 1% (Lobel & B lom me 2008).



Figure 2. Panel left: V isual brightness changes observed in C as between 1993 and 2003 (upper panel) are compared to the radial velocity variability (bwer panel) of an unblended photospheric absorption line of Fe i 5572. Panelright: D ynam ic spectra of H (left) and Fe i 5572 (right). O bservations are marked with left-hand tickmarks. The Fe i line strongly blue-shifts during the large outburst of m id 2000 when the atm osphere cools by 3000 K (see text).

### 3. The Millennium Outburst of Yellow Hypergiant Cas (F G Ia0)

Yellow hypergiants are post-red supergiants, rapidly evolving toward the blue supergiant phase (de Jager 1998). They are among the prime candidates for progenitors of Type II supernovae in the Galaxy. In July 2000 we observed the form ation of new TiO bands in F-type hypergiant Cas during a strong V-brightness decrease of 1<sup>m</sup> :4 (Fig. 2). Synthetic spectrum calculations revealthat  $T_e$  decreased by at least 3000 K, from 7250 K to ' 4250 K, and the spectrum became comparable to an early M -type supergiant. The TiO bands signal the form ation of a cool circum stellar gas shell with  $T_{gas}$  < 4000 K due to the supersonic expansion of the photosphere and upper atm osphere. We observe a shell expansion velocity of  $35 \ 2 \ \text{km} \ \text{s}^{1}$  from the T iO bands. From the synthetic spectrum ts to the bands we compute an exceptionally large massloss rate of M-' 5.4 10 <sup>2</sup> M yr <sup>1</sup>, (5% of M in 200 d). It is therefore one of the largest stellar outburst events that has directly been monitored so far (Lobelet al. 2003). The right-hand panel of Fig. 2 shows dynam ic spectra of H and Fe i 5572 observed between 1993 and late 2003. The radial velocity curves of the H absorption core and the photospheric Fe i lines (white dashed lines) reveal a velocity strati ed dynam ic atm osphere. A strong collapse of the upper H atm osphere and the lower photosphere precedes the outburst event during the large-am plitude pre-outburst pulsation cycle of 1999.

Based on radiative transport calculations G orlova et al. (2006) presented a model for the near-IR CO emission emerging from cool atmospheric layers in the immediate vicinity of Cas' photosphere. The gas kinetic temperature minimum in the model results from a periodical pulsation-driven shock wave.



Figure 3. Panels left: Large spectral resolution observations of the M g iih & k and Balmer H lines (boldly drawn lines) reveal long-term chrom ospheric variability in O ri. Best non-LTE ts (thin drawn lines) constrain the gas kinetic tem perature and electron density structure of the inner chrom osphere. The m odel of the photosphere below the chrom osphere has  $T_{\rm e}$  = 3500 K . (bottom panel). Panels right: The chrom ospheric em ission line spectrum (upper panel) is well-reproduced using only iron lines (dashed drawn line), and a m odel  $T_{\rm gas}$  m axim um ' 5550 K . The lower panel shows the best t (thin line) to the optical spectrum around strong -system band-heads of T io .

#### 4. M odeling C hrom ospheric D ynam ics of R ed Supergiant B et elgeuse

1-D m odels of the therm odynam ic and dynam ic structure of 0 ri's photosphere and inner chrom osphere (to 10 R) are developed in Lobel & Dupree (2000). M odern m ethods in radiative transfer m odeling venture w ell beyond outdated approaches that, for example, can only consider escape probabilities and radio uxes. The latter ux-integrated models completely fail to produce fundam ental chrom ospheric lines (e.g., Balmer H, and Mg ii, Ca ii, K i resonance lines, etc.). A courate m odels (1-D orm ulti-D) for these in portant spectral lines are required for unraveling the wind-driving mechanism and related heating physics in the outer atm ospheres of cool stars. The transport of radiation does not discrim inate between so-called warm and cool gas components'. In transfer calculations (irrespective of the num ber of dim ensions), the m odel tem perature and electron density structures (Ne) must properly excite chrom ospheric absorption (e.g., H, H) and emission lines, including the central self-reversals of optically thick em ission lines. The model in Fig. 3 shows the radialm ean values of  $T_{gas}$  and N  $_{e}$ up to 10 R, based on detailed ts to the classical chrom ospheric indicators observed in 0 ri. It correctly produces the observed near-UV emission line spectrum. A model without a gas kinetic temperature structure increasing to

5550 K in the inner chrom osphere cannot excite the subordinate H -transition of 300 m A. The entire absorption line would become invisible with respect to strong TiO absorption in the cooler photosphere (Lobel & Dupree 2000). The Mgiih & k em ission lines would also vanish. The fully intensity-saturated selfscattered cores of these broad lines (about the strongest chrom ospheric em ission lines observed in any cool star) result from a very large column of (warm)



Figure 4. Panels left: Detailed non-LTE t (thin line in upper panel) to the spatially resolved Sii 2516 em ission line observed in Oriat 75 m as with HST-STIS (boldly drawn line). The radial structures of the model are shown in the lower panels.  $T_{gas}$  is in kK units, and V wind and V turb in km s<sup>1</sup>. Panels right: 3-D mean ow velocities are inferred from detailed ts (thin drawn lines in upper panel) to the asym metric shape of the Sii line (bold lines). The reversal of the em ission line component intensity reveals simultaneous upand down ow across the front hem isphere in 1998 September.

chrom ospheric plasm a that dom inates the overall gas kinetic tem perature- and density-structure of the inner chrom osphere. A much steeper outwardly decrease of N<sub>e</sub> (i.e., by an order of magnitude over 2 R) yields too large decrements of totalwidths computed for spatially resolved chrom ospheric em ission lines. Large variations of chrom ospheric em ission line widths are consistently not observed in spatially resolved raster scans across the inner chrom osphere with HST-STIS in 1998 1999. The macro-broadening velocity is supersonic (9 1 km s<sup>1</sup>), and observed to be highly isotropic and hom ogeneous (Lobel & Dupree 2001) across the inner chrom ospheric disk.

Detailed transfer ts to the tem poral variability and spatial intensity reversals observed in the em ission line components of Sii 2516 reveal that the inner chrom osphere oscillates asymmetrically with simultaneous up-and down ow (4 km s<sup>1</sup>) across the front hem isphere (Fig. 4). Interestingly, Freytag & Hofner (2008) recently presented 3-D hydrodynamic models with large asymmetric movements of the photosphere and strong (M<sub>rad</sub>>2) radially propagating shock waves. Strong shock waves are an e cient mechanism for transporting mechanical momentum very far from the stellar surface. They can support the co-existence of dust-sustaining low-tem perature conditions (T<sub>dust</sub>' 600 K) and (possibly shock-excited) chrom ospheric em ission lines observed with STIS far inside the circum stellar dust envelope (Fig. 5; Lobel 2005) up to 3<sup>00</sup> (> 100 R).

# 5. Conclusions

M echanical wave action (provided by circum stellar density waves, propagation of pressure waves, and astrophysical turbulence) due to atm ospheric pulsation is



Figure 5. Panel left: Relative aperture positions of spatially resolved spectroscopic observations with HST-STIS in 2002-2003 compared to a 9.8 m in age (MMT-MIRAC2) of Ori/s inner circum stellar dust envelope (adapted from H inz et al. 1998). Panel right: The Mg ii k resonance em ission line observed across the inner and upper chrom osphere with the 63 200 m as aperture compared to a false color near-UV in age (HST-FOC). The line is scaled to equal intensities and reveals in portant pro le shape changes of the central self-scattering core, signaling outward acceleration of the chrom ospheric w ind.

a very important aspect for studying the detailed physics of active structuring, acceleration, and heating mechanisms of the extended winds and circum stellar environments of hot and cool massive stars. Recent advances in quantitative spectroscopy that combine 3-D radiative transferm ethods with multi-D hydrodynamic and sem i-empiric modeling provide powerfulnew methods for pinpointing these mechanisms.

A cknow ledgm ents. The contributions from many Cas observers over the past 15 years are gratefully acknow ledged. This research is based in part on data of O ri obtained with the NASA/ESA Hubble Space Telescope, collected at the STSCI, operated by AURA Inc., under contract NAS5-26555. Financial support has been provided by STSCI grant HST-GO-09369.01. The Belgian Federal Science Policy is gratefully acknow ledged for nancial support for 3-D radiative transferm odeling research of massive stars.

#### References

- de Jager, C. 1998, A & A Rev., 8, 145
- Freytag, B., & Hofner, S. 2008, A & A, 483, 571
- Gorbva, N., Lobel, A., Burgasser, A.J., Rieke, G.H., Ilyin, I., & Stau er, J.R. 2006, ApJ, 651, 1130
- Hinz, PM , Angel, J.R.P., Ho mann, W .F., et al. 1998, Nat, 395, 251
- Lobel, A. 2005, in ESA SP-560, 13th Cambridge W orkshop on Cool Stars, Stellar System s, and the Sun, ed. F. Favata et al., (Noordwijk: ESA Publications Division), 327
- Lobel, A., & Blomme, R. 2008, ApJ, 678, 408
- Lobel, A. et al. 2003, ApJ, 583, 923
- Lobel, A., & Dupree, A.K. 2001, ApJ, 558, 815
- Lobel, A., & Dupree, A.K. 2000, ApJ, 545, 454