

# A THIRD GIANT PLANET ORBITING HIP 14810

J. T. WRIGHT

Department of Astronomy, 226 Space Sciences Building, Cornell University, Ithaca, NY 14853  
 jtwright@astro.cornell.edu

D. A. FISCHER

Department of Physics and Astronomy, San Francisco State University, San Francisco, CA 94132

ERIC B. FORD, D. VERAS, J. WANG

Department of Astronomy, University of Florida, 211 Bryant Space Science Center, P.O. Box 112055, Gainesville, FL 32611-2055

G. W. HENRY

Center of Excellence in Information Systems, Tennessee State University, 3500 John A. Merritt Blvd., Box 9501, Nashville, TN 37209

G. W. MARCY, A. W. HOWARD<sup>1</sup>

Department of Astronomy, 601 Campbell Hall, University of California, Berkeley, CA 94720-3411

AND

JOHN ASHER JOHNSON<sup>2</sup>

Institute for Astronomy, University of Hawai'i, Honolulu, HI 96822

*Draft version February 12, 2022*

## ABSTRACT

We present new precision radial velocities and a three-planet Keplerian orbit fit for the  $V = 8.5$ , G5 V star HIP 14810. We began observing this star at Keck Observatory as part of the N2K Planet Search Project. Wright et al. (2007) announced the inner two planets to this system, and subsequent observations have revealed the outer planet planet and the proper orbital solution for the middle planet. The planets have minimum masses of 3.9, 1.3, and 0.6  $M_{\text{Jup}}$  and orbital periods of 6.67, 147.7, and 952 d, respectively. We have numerically integrated the family of orbital solutions consistent with the data and find that they are stable for at least  $10^6$  yr. Our photometric search shows that the inner planet does not transit.

*Subject headings:* planetary systems — stars: individual (HIP 14810)

## 1. INTRODUCTION

The first multiple exoplanet system detected around a normal star<sup>3</sup> was the triple system  $\nu$  Andromadae (Butler et al. 1999). Today, over 30 systems comprising more than one planet are known (Wright et al. 2009), including seven triple systems, two quadruple systems ( $\mu$  Arae (Pepe et al. 2007) and GJ 581 (Mayor et al. A&A submitted)) and the quintuple system 55 Cancri (Fischer et al. 2008).

Individual multiplanet systems offer insights into the dynamical evolution of planetary systems that singleton systems cannot. For instance, Ford, Lystad, & Rasio (2005) showed that  $\nu$  Andromadae bears the scars of strong planet-planet scattering events preserved in their planets' orbital parameters. Other systems show evidence of migration and eccentricity pumping through mean-motion resonances (MMRs), which may be the signpost of convergent migration in multi-planet systems (e.g. Kley, Peitz, & Bryden 2004). Multiplanet systems

with planet-planet interactions strong enough to be detected at current RV precision (as in GJ 876, Rivera et al. 2005) allow for measurement of the inclination of the system, providing true planet masses.

Comparison of multiplanet systems as an ensemble to singleton systems provides observational constraints to theories and models of the early dynamical evolution and migration history of planetary systems. Wright et al. (2009) showed that the while the eccentricity distribution of planets in multi-planet systems is similar to that of apparently singleton systems, their semimajor axis distributions differ significantly. The concentration of planets with orbital periods near 3-days seen in the single-planet systems is absent in multi-planet systems, as is the sharp jump beyond 1 AU in planet frequency. The fact that these features (the 3-day pileup and the 1 AU jump) are functions planetary multiplicity strongly suggests that planet-planet interactions play a key role in both migration and the origin of eccentricities.

## 2. THE TENTH TRIPLE SYSTEM

The announcement of HIP 14810  $d$  herein marks the tenth system with three or more detected planets and only the sixth known to host three or more giant

<sup>1</sup> Townes Fellow, Space Sciences Laboratory

<sup>2</sup> NSF Postdoctoral Fellow

<sup>3</sup> Prior to this, Wolszczan & Frail (1992), detected the first planets outside the solar system: three extraordinary planets orbiting the pulsar PSR 1257+12.

( $M \sin i > 10 M_{\text{Earth}}$ ) planets.<sup>4</sup> The minimum masses and orbital periods of the planets in this systems are similar to those of  $\nu$  Andromedae, but with the inner and outermost components reversed. For the three planets of HIP 14810 we find:  $M \sin i = 3.9, 1.2$ , and  $0.6 M_{\text{Jup}}$ , and  $P = 6.67, 147.7$ , and  $952$  d, respectively. We find modest but significant eccentricities for all 3 components (0.14, 0.16, and 0.17, respectively).

Table 1 contains a summary of the stellar properties of HIP 14810 (= BD+20 518), which sits at 53 pc ( $\pi = 18.7 \pm 1.3$ , van Leeuwen 2009) and has  $V = 8.5$ . We have performed an LTE analysis of our template spectra for HIP 14810 and derived its mass and radius using the methods described in Valenti & Fischer (2005). Although HIP 14810 is a solar mass star ( $M = 0.99 M_{\odot}$ ), its metallicity ( $[\text{Fe}/\text{H}] = +0.26$ ) and evolutionary status ( $\Delta M_V = 0.63$  mag, as calculated in Wright (2004)) give it a spectral type of G5. Its low rotation ( $v \sin i \sim 0.5 \pm 0.5 \text{ km s}^{-1}$ ) and Ca II H & K activity levels ( $S = 0.16$ , measured with the methods described in Wright et al. (2004)) are consistent with it being an old star (age  $\sim 8$  Gyr). This combined with its relatively small distance from the main sequence make it a particularly good radial velocity target, since it is expected to exhibit very low levels of jitter (Wright 2005).<sup>5</sup>

### 3. VELOCITIES AND ORBITAL SOLUTION

We began observations of HIP 14810 in 2005 as part of the N2K survey (Fischer et al. 2005) at Keck Observatory using HIRES (Vogt et al. 1994) and our usual iodine technique (Butler et al. 1996) to achieve typical internal (random) errors of  $0.8\text{--}1.4 \text{ ms}^{-1}$ . The presence of the innermost planet and the large residuals to its orbital fit inspired the California Planet Search consortium to continue regular observations of this system at Keck.

Wright et al. (2007) announced<sup>6</sup> the inner two planets of HIP 14810, though with a rather poor fit for the  $c$  component due to the poor phase coverage and the unaccounted-for effects of the  $d$  component. Also complicating the fit was what is now obviously a spurious data point, acquired during early dusk when significant contamination from the Solar spectrum likely produced an erroneous radial velocity measurement. We have applied a more rigorous data retention scheme (based solely on the measured internal errors<sup>7</sup>, not deviations from a fit) to the data set presented in Table 2. These velocities and uncertainties supersede our previously published values for this star, as we continue to refine our data reduction pipeline (Wright et al. 2009). Note that the times

given in the table are in heliocentric Julian days, and the quoted errors are our internal (random) errors, with no “jitter” included.

By 2007, residuals to a 2-planet fit clearly showed coherent structure indicative of an outer companion. As Figure 1 shows, by late 2008, these residuals appeared to describe one complete orbit of a  $P \sim 950$  d planet with modest eccentricity ( $e \sim 0.2$ ). We have performed a Monte Carlo false alarm probability (FAP) analysis of the complete set of residuals to determine the likelihood that an orbital fit of this quality could have been arrived at by chance. This method is very similar to the FAP analyses in Butler et al. (2006); Wright et al. (2007); Butler et al. (2009); Howard et al. (2009), and we refer the reader to those works for more details. After binning the data in 24-hour intervals and subtracting the best 2-planet fit, we redrew these residuals 1000 times (that is, we kept the times of observation the same for each trial, but at each time assigned a new velocity and velocity uncertainty pair randomly drawn from the entire set, with replacement). We then added the nominal orbital solution for the  $c$  component back into these new residuals and performed a thorough search for the best-fit 2-planet orbital solution to each of these 1000 realizations of the data. We measure the FAP as the fraction of these realizations for which we find a solution superior or equivalent in fit quality to the nominal solution.

Note that this is an extremely conservative test of the FAP of the new planet,  $d$ , because we have not restricted the parameter search for the FAP trials to long period or low eccentricity planets. The best-fit solutions found for the artificial data sets are thus often at short periods ( $P < 3\text{d}$ ) and/or high eccentricities  $e > 0.7$  (see the discussion in Butler et al. 2009). Nonetheless, even with this large parameter space available, we find that none of our 1000 realizations produced a better fit than the actual data, yielding an FAP  $< 0.1\%$ .

We have fit the data using the publicly available multi-planet RV-fitting IDL package `RV_FIT_MP`, described in Wright & Howard (2009). In Table 3 we present the best 3-planet Keplerian (kinematic) fit, which yields r.m.s. residuals of  $2.3 \text{ ms}^{-1}$ , and we plot the fit and velocities in Figure 1. We have observed just over one complete orbit for the outer component, HIP 14810 $d$ , and so its orbital parameters are sensitive to the assumption that there are no additional, external planets detectably influencing the velocities. In particular, it is possible that such an additional planet could be contributing a nearly linear trend to the observed radial velocities, a trend which might be absorbed into the  $d$  component’s orbital parameters as an inflated eccentricity. Any such degeneracy will be broken in the near future as the  $d$  component completes its second orbit. Continued observation of this system will thus reveal the presence of any additional detectable planets and allow for further analysis of their interactions (e.g. Ford 2005; Wright et al. 2008).

### 4. DYNAMICAL MODELING AND STABILITY ANALYSIS

We have performed long-term numerical integrations to test for stability of the orbital solution here, under the assumption that there are only three planets in the system. For these long-term stability tests, we applied direct n-body integrations to 100 orbital solutions consis-

<sup>4</sup> The others are  $\nu$  Andromedae, 55 Cnc,  $\mu$  Arae, HD 69830 (Lovis et al. 2006), and HD 37124 (Vogt et al. 2005).

<sup>5</sup> We expect, based on hundreds of similar stars,  $2 \text{ m s}^{-1}$  of jitter, consistent with the residuals to our best fit. This value is somewhat lower than predicted by the formulae in Wright (2005) because that work included instrumental sources of noise associated with the HIRES CCD detector in place prior to Aug 2004. The new detector has significantly better charge transfer properties, and apparently contributes a negligible amount to our overall error budget.

<sup>6</sup> A preliminary orbit for the  $b$  component also appears in the *Catalog of Nearby Exoplanets* (Butler et al. 2006).

<sup>7</sup> We derive radial velocity measurements from each of many independent “chunks” of spectrum for each observation. Our *internal error* for a given observation is determined from the variance of these velocities (Wright 2005).




FIG. 1.— Radial velocity curves for the HIP 14810 triple system.




FIG. 2.— *Top Panel:* The 1099 photometric observations of HIP 14810 in the combined Strömgren  $(b + y)/2$  passband, acquired with the T11 0.8m APT over four observing seasons and plotted modulo the 6.673855-day orbital period of the inner planet HIP 14810b. Phase 0.0 corresponds to the predicted time of mid transit. A least-squares sine fit at the orbital period yields a semi-amplitude of only  $0.00008 \pm 0.00006$  mag. *Bottom Panel:* The photometric observations of HD 14810 near the predicted time of transit replotted with an expanded scale on the abscissa. The solid curve shows the depth (0.01 mag) and duration ( $\pm 0.0125$  phase units) of a central transit, computed from the orbital elements and the planetary and stellar properties. The uncertainty in the predicted transit time is smaller than the diameter of the plotted points in the lower panel. The brightness levels of HIP 14810 within and outside the transit window agree to 0.00013 mag. Therefore, transits of HIP 14810b are ruled out for all reasonable densities.

TABLE 1  
STELLAR PROPERTIES OF HIP 14810

Parameter	Value
Spectral Type	G5 V
RA	$03^{\text{h}}11^{\text{m}}14^{\text{s}}.230$
Dec.	$+21^{\circ}05'50''.49$
B-V	0.78
V	8.52
Distance (pc)	$52.9 \pm 4.1$
$M_V$	4.9
$T_{\text{eff}}$ (K)	$5485 \pm 44$
$\log g$ [ $\text{cm s}^{-2}$ ]	$4.220 \pm 0.06$
[Fe/H]	$+0.26 \pm 0.03$
$v \sin i$	$0.54 \pm 0.5 \text{ km s}^{-1}$
Mass ( $M_{\odot}$ )	$0.99 \pm 0.04$
Radius ( $R_{\odot}$ )	$1.0 \pm 0.06$
S	0.16
$\log R'_{\text{HK}}$	-5.01
$\Delta M_V$ (mag)	0.64

tent with the RV data, assuming an edge-on orientation.<sup>8</sup> We integrated for at least  $10^8$  years using the hybrid integrator in *Mercury* (Chambers 1999). For the majority of each integration, *Mercury* uses a mixed-variable symplectic integrator (Wisdom & Holman 1991) with a time step approximately equal to a hundredth of the Keplerian orbital period calculated at a semi-major axis equal to the pericenter distance of the closest planet. During close encounters, *Mercury* uses a Bulirsch-Stoer integrator with an accuracy parameter of  $10^{-10}$ . We identified each set of initial conditions as an unstable system if: 1) two planets collide, 2) a planet is accreted onto the star (astrocentric distance less than 0.005 AU), or 3) a planet is ejected from the system (astrocentric distance exceeds 100 AU).

All of our simulations proved stable, and we manually verified that in all cases the final orbits were qualitatively similar to the initial conditions. We also ran an additional set of simulations for inclined (but coplanar) orientations of the system (i.e. smaller values of  $\sin i$ , and thus higher true planet masses and larger planet-planet interactions) and find the system to be stable for all tested scenarios with  $i > 3^{\circ}$ .

In addition, we performed dynamical analyses of three planet solutions using ensembles of initial conditions generated via Bayesian posterior sampling methods (e.g. Ford 2006), both ignoring and including the planet-planet interactions. We found that the superposition of Keplerians approximation is a very good approximation for the nominal edge-on configuration. As the orbits approach face-on, the planet-planet interactions eventually become significant for inclinations of a few degrees. While we only considered coplanar configurations, we expect that our results are likely representative for relative inclinations of up to 20 or 30 degrees.

## 5. PHOTOMETRIC OBSERVATIONS

<sup>8</sup> We have estimated the uncertainties in the orbital parameters in Table 3 using a variant of the bootstrapping described in Butler et al. (2006). We generate a suite of plausible data sets and calculate the best-fit solution for each realization. We selected 100 of these realizations as the basis for this stability analysis, as in Wright et al. (2009).

TABLE 2  
RADIAL VELOCITIES FOR HIP 14810

HJD JD-2440000	Velocity m s <sup>-1</sup>	Uncertainty m s <sup>-1</sup>
13693.76627	-147.5	1.1
13694.83716	-489.3	1.2
13695.91489	-242.3	1.1
13723.79085	146.4	1.0
13724.69303	304.8	1.2
13746.81748	-13.1	1.1
13747.85573	-450.0	0.9
13748.73690	-448.5	1.2
13749.74186	-86.0	1.1
13751.90068	342.3	1.0
13752.80978	225.5	0.8
13753.69384	-96.2	1.1
13753.81261	-154.0	1.0
13753.90329	-195.5	1.2
13775.83626	-256.0	0.9
13776.81286	107.0	1.2
13777.72301	332.0	1.2
13778.72061	398.1	1.1
13779.74412	220.8	1.2
13961.13028	-293.8	1.1
13962.13341	-426.9	1.0
13981.97179	-492.7	1.1
13982.94950	-217.1	1.2
13983.98363	137.6	1.0
13984.09915	169.9	1.2
13984.98803	333.0	1.1
13985.10437	343.6	1.2
14023.96949	88.2	1.3
14047.98188	-396.5	1.2
14129.79605	-208.9	0.9
14130.74873	125.2	0.9
14131.84182	334.8	1.2
14138.80623	336.4	1.2
14307.13007	92.8	1.0
14336.07711	-436.4	1.0
14344.03162	34.5	1.0
14345.14559	306.4	0.9
14396.82488	-108.0	1.0
14397.91112	254.0	1.0
14398.88586	411.3	1.1
14399.89293	364.3	1.4
14428.00013	-184.5	1.0
14428.87298	-470.7	1.0
14492.75897	356.1	0.9
14544.74216	260.4	1.1
14674.09015	219.2	1.2
14718.05894	146.8	1.1
14723.08578	-434.2	1.0
14725.92426	348.5	1.2
14727.05509	271.3	1.0
14727.97892	-45.6	1.1
14777.99836	67.5	1.1
14805.83497	358.4	1.1

We acquired 1099 good photometric observations of HIP 14810 during four observing seasons spanning 1180 days between 2005 November and 2009 February with the T11 0.8m automated photometric telescope (APT) at Fairborn Observatory. The T11 APT and its two-channel Strömgren *b* and *y* photometer are very similar to the T8 0.8m APT and precision photometer described in Henry (1999).

The measurements of HIP 14810 were made differentially with respect to the comparison star HD 18404 ( $V = 5.80$ ,  $B - V = 0.42$ , F5 IV). We combined the Strömgren *b* and *y* differential magnitudes into a single  $(b + y)/2$  passband to improve the precision of each measurement. Henry (1999) gives further details on the

TABLE 3  
ORBITAL ELEMENTS FOR EXOPLANETS IN THE HIP 14810 SYSTEM

Property	<i>b</i>	<i>c</i>	<i>d</i>
Per ( <i>d</i> )	6.673855(19)	147.730(65)	952(15)
$T_0$ (JD-2440000)	13694.5980(70)	14672.2400(73)	14317.1980(73)
$e$	0.14270(94)	0.164(12)	0.173(37)
$\omega$ ( $^\circ$ )	159.32(38)	329.0(2.5)	286(19)
$K$ (ms <sup>-1</sup> )	424.48(44)	50.01(46)	12.03(49)
$M \sin i$ ( $M_{\text{Jup}}$ )	3.88(32)	1.28(10)	0.570(52)
<i>a</i> (AU)	0.0692(40)	0.545(31)	1.89(11)
r.m.s. (ms <sup>-1</sup> )		2.3	
$\chi^2_\nu$		1.01	
jitter (ms <sup>-1</sup> )		2	
$N_{\text{obs}}$		53	

NOTE. — For succinctness, we express uncertainties using parenthetical notation, where the least significant digit of the uncertainty, in parentheses, and that of the quantity are to be understood to have the same place value. Thus, “0.100(20)” indicates “0.100 $\pm$ 0.020”, “1.0(2.0)” indicates “1.0  $\pm$  2.0”, and “1(20)” indicates “1  $\pm$  20”.

acquisition, reduction, and calibration of the APT data.

The standard deviation of a single observation from the mean of the entire dataset is 0.00158 mag, which closely matches the typical measurement precision with this APT. Periodogram analysis of the full dataset finds no significant periodicities between one and several hundred days. In particular, least-squares sine fits to the 6.673855- and 147.73-day orbital periods of the inner (*b*) and middle (*c*) planets yield semi-amplitudes of only  $0.00008 \pm 0.00006$  and  $0.00021 \pm 0.00006$  mag, respectively, confirming that stellar activity is not the cause of the radial-velocity variations at these two periods. Although we do not expect radial velocity variations due to long period variations in stellar activity in such old stars (Wright et al. 2008), we have also searched for photometric variations at the period of the outer (*d*) planet ( $P = 952$  d). While our dataset is not yet long enough to make this determination definitively, we see no suggestion of photometric variations at this period so far.

In the top panel of Figure 2, we plot the APT brightness measurements of HIP 14810 against phases computed from the 6.673855-day orbital period of the inner planet and a time of mid transit, JD 2,453,693.5856  $\pm$  0.0022, predicted from the orbital elements in Table 3. The observations near the predicted time of transit are replotted with an expanded horizontal scale in the bottom panel of Figure 2. The solid curve shows the depth (0.01 mag) and duration ( $\pm 0.0125$  phase units) of a central transit of planet *b*, computed from the orbital elements and the planetary and stellar properties. The precisely determined orbit of HIP 14810*b* (Figure 1, top panel) translates into an uncertainty in the mid-transit time that is smaller than the plotted points in the bottom of Figure 2. The mean of the 120 observations within the transit window is  $2.81423 \pm 0.00015$  mag; the mean of the 979 observations outside the window is  $2.81410 \pm 0.00005$  mag. Thus, the brightness levels inside and outside of the predicted transit window agree to within 0.00013 mag, allowing us to rule out transits of the *b* component for all reasonable densities.

## 6. CONCLUSIONS

HIP 14810 is orbited by at least 3 giant planets, the outermost of which (HIP 14810 *d*) has completed just

over one orbit since we began monitoring it at Keck Observatory as part of the N2K project in 2005. This makes it only the sixth system known to host more than 2 giant ( $M \sin i > 10 M_{\text{Earth}}$ ) planets. In retrospect, the previously published orbital solution for the middle (c) component was hampered by a spurious data point, poor phase coverage and the unaccounted-for effects of the d component. Continued monitoring will reveal the existence of any additional outer planets.

We have performed a dynamical analysis for a suite of orbital solutions assuming no fourth planet, and find the system to be stable at nearly any inclination ( $i < 3^\circ$ ). We have performed a photometric survey and find that the b component does not transit.

The work herein is based on observations obtained at the W. M. Keck Observatory, which is operated jointly by the University of California and the California Institute of Technology. The Keck Observatory was made possible by the generous financial support of the W.M. Keck Foundation. We wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to

have the opportunity to conduct observations from this mountain.

J.T.W. received support from NSF grant AST-0504874. G.W.M. received support from NASA grant NNG06AH52G, and D.A.F. from NASA grant NNG05G164G and the Cottrell Science Scholar Program. G.W.H. acknowledges support from NASA, NSF, Tennessee State University, and the State of Tennessee through its Centers of Excellence program. A.W.H. gratefully acknowledges support from a Townes Postdoctoral Fellowship at the UC Berkeley Space Sciences Laboratory. E.B.F. acknowledges the support of NASA RSA 1326409, and E.B.F. and D.V. acknowledge support of NASA grant NNX09AB35G. J.A.J. is supported by NSF grant AST-0702821.

We thank the anonymous referee for a helpful review. We acknowledge the University of Florida High-Performance Computing Center for providing computational resources and support that have contributed to the results reported within this paper. This research has made use of NASA's Astrophysics Data System. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France

*Facility* Keck:I

#### REFERENCES

- Butler, R. P., Marcy, G. W., Williams, E., McCarthy, C., Dosanjh, P., & Vogt, S. S. 1996, *PASP*, 108, 500
- Butler, R. P., Marcy, G. W., Fischer, D. A., Brown, T. M., Contos, A. R., Korzennik, S. G., Nisenson, P., & Noyes, R. W. 1999, *ApJ*, 526, 916–927
- Butler, R. P., Wright, J. T., Marcy, G. W., Fischer, D. A., Vogt, S. S., Tinney, C. G., Jones, H. R. A., Carter, B. D., Johnson, J. A., McCarthy, C., & Penny, A. J. 2006, *ApJ*, 646, 505–522
- Butler, R. P., Howard, A. W., Vogt, S. S., & Wright, J. T. 2009, *ApJ*, 691, 1738–1743
- Chambers, J. E. 1999, *MNRAS*, 304, 793–799
- Fischer, D. A., Laughlin, G., Butler, P., Marcy, G., Johnson, J., Henry, G., Valenti, J., Vogt, S., Ammons, M., Robinson, S., Spear, G., Strader, J., Driscoll, P., Fuller, A., Johnson, T., Manrao, E., McCarthy, C., Muñoz, M., Tah, K. L., Wright, J., Ida, S., Sato, B., Toyota, E., & Minniti, D. 2005, *ApJ*, 620, 481–486
- Fischer, D. A., Marcy, G. W., Butler, R. P., Vogt, S. S., Laughlin, G., Henry, G. W., Abouav, D., Peek, K. M. G., Wright, J. T., Johnson, J. A., McCarthy, C., & Isaacson, H. 2008, *ApJ*, 675, 790–801
- Ford, E. B. 2005, *AJ*, 129, 1706–1717
- Ford, E. B. 2006, In *New Horizons in Astronomy*: Frank N. Bash Symposium, S. J. Kannappan, S. Redfield, J. E. Kessler-Silacci, M. Landraiu, and N. Drory, eds., volume 352 of *Astronomical Society of the Pacific Conference Series*, pp. 15–+
- Ford, E. B., Lystad, V., & Rasio, F. A. 2005, *Nature*, 434, 873–876
- Henry, G. W. 1999, *PASP*, 111, 845–860
- Howard, A. W., Johnson, J. A., Marcy, G. W., Fischer, D. A., Wright, J. T., Henry, G. W., Giguere, M. J., Isaacson, H., Valenti, J. A., Anderson, J., & Piskunov, N. E. 2009, *ApJ*, 696, 75–83
- Kley, W., Peitz, J., & Bryden, G. 2004, *A&A*, 414, 735–747
- Lovis, C., Mayor, M., Pepe, F., Alibert, Y., Benz, W., Bouchy, F., Correia, A. C. M., Laskar, J., Mordasini, C., Queloz, D., Santos, N. C., Udry, S., Bertaux, J.-L., & Sivan, J.-P. 2006, *Nature*, 441, 305–309
- Pepe, F., Correia, A. C. M., Mayor, M., Tamuz, O., Couetdic, J., Benz, W., Bertaux, J.-L., Bouchy, F., Laskar, J., Lovis, C., Naef, D., Queloz, D., Santos, N. C., Sivan, J.-P., Sosnowska, D., & Udry, S. 2007, *A&A*, 462, 769–776
- Rivera, E. J., Lissauer, J. J., Butler, R. P., Marcy, G. W., Vogt, S. S., Fischer, D. A., Brown, T. M., Laughlin, G., & Henry, G. W. 2005, *ApJ*, 634, 625–640
- Valenti, J. A., & Fischer, D. A. 2005, *ApJS*, 159, 141–166
- van Leeuwen, F. 2009, *A&A*, 497, 209–242
- Vogt, S. S., Allen, S. L., Bigelow, B. C., Bresee, L., Brown, B., Cantrall, T., Conrad, A., Couture, M., Delaney, C., Epps, H. W., Hilyard, D., Hilyard, D. F., Horn, E., Jern, N., Kanto, D., Keane, M. J., Kibrick, R. I., Lewis, J. W., Osborne, J., Pardeilhan, G. H., Pfister, T., Ricketts, T., Robinson, L. B., Stover, R. J., Tucker, D., Ward, J., & Wei, M. Z. 1994, In *Proc. SPIE Instrumentation in Astronomy VIII*, David L. Crawford; Eric R. Craine; Eds., Volume 2198, p. 362, pp. 362–+
- Vogt, S. S., Butler, R. P., Marcy, G. W., Fischer, D. A., Henry, G. W., Laughlin, G., Wright, J. T., & Johnson, J. A. 2005, *ApJ*, 632, 638–658
- Wisdom, J., & Holman, M. 1991, *AJ*, 102, 1528–1538
- Wolszczan, A., & Frail, D. A. 1992, *Nature*, 355, 145–147
- Wright, J. T. 2004, *AJ*, 128, 1273–1278
- Wright, J. T. 2005, *PASP*, 117, 657–664
- Wright, J. T., & Howard, A. W. 2009, *ApJS*, 182, 205–215
- Wright, J. T., Marcy, G. W., Butler, R. P., & Vogt, S. S. 2004, *ApJS*, 152, 261–295
- Wright, J. T., Marcy, G. W., Fischer, D. A., Butler, R. P., Vogt, S. S., Tinney, C. G., Jones, H. R. A., Carter, B. D., Johnson, J. A., McCarthy, C., & Apps, K. 2007, *ApJ*, 657, 533–545
- Wright, J. T., Marcy, G. W., Butler, R. P., Vogt, S. S., Henry, G. W., Isaacson, H., & Howard, A. W. 2008, *ApJ*, 683, L63–L66
- Wright, J. T., Upadhyay, S., Marcy, G. W., Fischer, D. A., Ford, E. B., & Johnson, J. A. 2009, *ApJ*, 693, 1084–1099