

Families of Canonical Transformations by Hamilton-Jacobi-Poincaré equation. Application to Rotational and Orbital Motion

S. Ferrer and M. Lara^y

April 2, 2024

Abstract

The Hamilton-Jacobi equation in the sense of Poincaré, i.e. formulated in the extended phase space and including regularization, is revisited building canonical transformations with the purpose of Hamiltonian reduction. We illustrate our approach dealing with orbital and attitude dynamics. Based on the use of Whittaker and Andoyer symplectic charts, for which all but one coordinates are cyclic in the Hamilton-Jacobi equation, we provide whole families of canonical transformations, among which one recognizes the familiar ones used in orbital and attitude dynamics. In addition, new canonical transformations are demonstrated.

1 Introduction

Transformation of variables, including or not the change of time used as evolution parameter, is a basic tool in almost any problem of physics. A recent illustration of the usefulness of this technique, which original application goes back to Laplace and Euler, may be found in [19]. Changes of variables belong to Geometry and, therefore, do not depend on the physical system to which they may be applied. However, transformations are commonly derived for specific purposes and, hence, they are tightly related to the particular problem they are applied to.

A renewed interest on transformation of variables tailored to specific problems has recently emerged in three areas. On one hand there is a continuous search for more efficient numerical integrators demanded in many applied fields, ranging from research groups endowed with powerful computing facilities, to research involved in very long periods of time (see [5, 16] and references therein). On the other hand the Hamilton-Jacobi (H-J) equation is receiving new attention in extended phase space, connecting this technique with the variational principles of Lagrangian and Hamiltonian Mechanics (see [31] and references therein). Finally a third area, which has its roots in Poincaré's [30] fundamental

^DDepartamento de Matemática Aplicada, Universidad de Murcia, 30071 Espinardo, Spain

^yReal Observatorio de la Armada, 11110 San Fernando, Spain

problem of dynamics, deals with Hamiltonian systems of the form $H = H_0 + H_1$ where H_0 denotes an integrable system, and H_1 is considered a perturbation. The H-J equation is built from H_0 by searching for a generating function such that the new canonical variables satisfy some requirements of perturbation theories (see, for instance, [37, 17]). The content of this paper focuses in this last area. Finally note that the H-J approach has been used by many authors, along the history of this equation, looking for variables leading to separability. We do not consider that problem as such, although it might be included.

There is a long tradition in Classical and Celestial Mechanics | nowadays a common ground in many other fields | of using both canonical transformations and changes of independent variable dubbed as regularization. The H-J equation is the usual approach, in which the canonical transformation is obtained from either a principal or a characteristic function [18].

The characteristic function approach does not rely on the vanishing of the new Hamiltonian. It only requires a non-zero transformed Hamiltonian that is cyclic in all variables but depends on one or several momenta. The adequate selection of the new Hamiltonian will make the transformation successful for specific purposes. Thus, for instance, if one wants the transformed Hamiltonian to retain the topology of the original problem, probably several momenta must be retained in the new Hamiltonian. On the contrary, useful transformations for perturbation theory may look for transformed Hamiltonians that depend only on one momenta, thus simplifying the subsequent application of perturbation methods (which, in turn, have limited application to specific solutions as, for instance, quasiperiodic motion).

Sentences like "after several trials", which are frequent in the literature, might produce the impression that succeeding in the choice of the reduced Hamiltonian is a matter of educated guess or alquimia. However, the new Hamiltonian does not need to be specified since the beginning and, quite on the contrary, it may be handled formally throughout the process of computing the transformation [35]. Therefore, Hamiltonian problems solved by the H-J equation give rise to whole families of canonical transformations from which the selection of a particular one requires only the materialization of the new Hamiltonian¹.

We propose to obtain canonical transformations from a general form of a H-J equation that is formulated in the extended phase space. In principle, the only requirement imposed to the Hamiltonian function (6) is that H_0 has to be integrable. More precisely, the two families studied are presented using symplectic charts where the unperturbed part denotes a 1-DOF. In other words, we only deal with canonical transformations derived from integrable Hamiltonians that are cyclic in all but one variable, and perform two specific applications. In attitude dynamics, we find a family of transformations that

¹To make more clear our point from [35] (p. 412), where E denotes the Hamiltonian in the new variables, we quote "We are still free to choose the functional form of E . A convenient (and conventional) choice is $E(p_0; p_1; p_2) = \frac{1}{2} m^{-2} (2p_0^2)$. With this choice the momentum p_0 has dimensions of angular momentum, and the conjugate coordinate is an angle." As we will show this is not the only way of reasoning. Indeed, we arrive to the same choice when we impose the function called Kepler equation as one of the expressions defining the transformation. In other words, when the mean anomaly is chosen as one of the new coordinates. As Delaunay showed, in doing so he avoided the presence of mixed secular terms in this theory of the Moon. For a recent study on these variables from a geometric point of view the reader should consult [7]

orders for complete reduction of the Euler-Poinsot problem. In this case, we check that Sadov's [32, 33], Kinoshita's [24], Depri's [12], and the recent Fukushima's [17] reduced Hamiltonians are just particular cases of our family. Notably, our approach provides explicit transformations from Sadov's and Kinoshita's elements to Andoyer variables [3], thus overcoming them a major objection to both sets of elements of being constructed through implicit relations. Besides, we show that other selections of the new Hamiltonian provide new canonical transformations that may help in simplifying perturbation algorithms in attitude dynamics.

In orbital dynamics we obtain three different families of isochronal canonical transformations | derived from Henon's isochronal central potential [21, 6] | each one linked with a different form of the regularizing function. From our families we recover a variety of canonical transformations in the literature, ranging from historic ones like Delaunay [8], Levi-Civita [25, 26], or Hill [22], to the recent transformation of Yanguas [37]. In addition, we show that the set of Delaunay elements yields the simplest transformation of the isochronal family that reduces the Hamiltonian of the two body problem to one momentum and, besides, preserves the original form of the Kepler equation.

The two charts used enjoy similar characteristics. Both express the unperturbed part of the Hamiltonian families as genuine 1-D O F systems due to the central and axial symmetries that both possess. They belong to the category of superintegrable systems, but not maximally superintegrable (see Fasso [14] and Ortega and Ratiu [29]); only for particular choices of the parameters both systems fall into this last category. Moreover, from the computational point of view, these charts introduce great simplification in intermediary expressions which, in turn, lead to a deep insight about each family. Many authors still rely on Eulerian variables when dealing with rotational dynamics (see for instance Marsden and Ratiu [28]), and spherical variables for orbital type motions (see Goldstein et al. [18], Jose and Saletan [23] and Susman and Wisdom [35]).

In spite of the fact that the two families considered are independent and consequently, they are studied in a totally self-contained manner, we make both part of the paper because this allows to obtain a better insight in the way we approach the H-J formalism. A final comment is due. Although both families have parameters, we have made the analysis for the generic case only. With respect to the orbital case, as an illustration, we have added details when we restrict to Keplerian systems, showing the connection of the new variables with the anomalies. We would like to make clear that we are not doing a review of regularizations. In other words, it is left for the reader to try other canonical transformations within the general scheme presented here.

2 A General Form of the Hamilton-Jacobi Equation

We only deal with Hamiltonians of the type

$$K(x_0; x; X_0; X; \epsilon) = (X_0 + H); \quad (1)$$

where $x = (x_1; \dots; x_n)$ are coordinates and $X = (X_1; \dots; X_n)$ conjugate momenta; x_0 is the independent variable and X_0 its conjugate momenta in the extended phase space formulation, in which we restrict to the manifold $K = 0$; ϵ is a vector of parameters, and

the Hamiltonian H as well as the "regularizing factor" may depend on all or some of the parameters defining :

$$= (x_0; x; X_0; X); \quad H = H(x_0; x; X); \quad (2)$$

A dash in the place of a variable is used to remark that the corresponding variable is not present.

Hamilton equations are

$$\begin{aligned} \frac{dx_0}{d} &= \frac{\partial K}{\partial X_0}; & \frac{dX_0}{d} &= \frac{\partial K}{\partial x_0}; \\ \frac{dx_i}{d} &= \frac{\partial K}{\partial X_i}; & \frac{dX_i}{d} &= \frac{\partial K}{\partial x_i}; \quad (i = 1; \dots; n) \end{aligned} \quad (3)$$

where t is the evolution parameter of the flow. Note that the first of the previous equations reads

$$\frac{dx_0}{d} = \quad (4)$$

which tells the function ought to verify that > 0 in its domain. Moreover, in the case of conservative systems $H = H(x; X)$, X_0 is an integral, and the manifold $K = 0$ may be also seen as $X_0 = H = \text{constant}$.

We are interested in canonical transformations

$$(x_0; x; X_0; X) \xrightarrow{T} (y_0; y; Y_0; Y) \quad (5)$$

in the sense of Poincare. More precisely, we look for transformations such that they simplify Hamiltonian systems defined by functions H which can be written as

$$H = H_0 + H_1 \quad (6)$$

where $H_0 = H_0(x; X)$ defines an integrable system, and $H_1 = H_1(x_0; x; X)$ is a perturbation. Specially, we focus on canonical transformations such that the new Hamiltonian $K = K_0 + K_1$ satisfies

$$K_0 = (X_0 + H_0) = (; ; Y_0; Y); \quad (7)$$

ie. the full reduction of the unperturbed part is carried out. This Eq. (7) is what we refer as the variant of Poincare to the H-J equation; the classical case chooses $(; ; Y_0; Y) = 0$.

The transformations are defined by

$$X_i = \frac{\partial W}{\partial x_i}; \quad Y_i = \frac{\partial W}{\partial Y_i}; \quad i = 0; \dots; n \quad (8)$$

derived from a generator $W = W(x_0; x; Y_0; Y)$ that is a complete solution of the generalized H-J equation

$$\frac{\partial W}{\partial x_0} + H_0(x; X) + \frac{\partial W}{\partial x} \frac{\partial W}{\partial x_0} = (Y_0; Y); \quad (9)$$

Thus, the Hamiltonian K in the new variables will take the form

$$K = K_0 + K_1 = \quad + H_1 \quad (10)$$

where H_1 and \quad are expressed in the new variables. Note that in what follows we take $H_1 = 0$.

In this paper we limit to generators of the form

$$W = \sum_{0 \leq i < n-1}^X x_i Y_i + R(x_n; Y_0; Y); \quad (11)$$

and regularizing function $\quad = (x_n; X_0; X_1; \dots; X_{n-1})$. Hence, from Eq. (9) we may write

$$H_0(x; Y_1; \dots; Y_{n-1}; \frac{\partial R}{\partial x_n}) = \frac{(Y_0; Y_1; \dots; Y_n)}{(x_n; Y_0; Y_1; \dots; Y_{n-1})} Y_0; \quad (12)$$

Depending on the form of H_0 , Eq. (12) may be solved for $\partial R / \partial x_n$ and, therefore, R is computed from a quadrature, which solution will depend on the choices made for \quad and \quad . Other possibilities are under investigation [15]. Note that, in fact, there is no reason why we should impose on H_0 to be cyclic in x_0 . What we have presented above, properly adapted, remains valid if we lift that constraint. This is referred in the literature as nonautonomous systems; the driven oscillator, the relativistic particle, etc are just simple examples within that category (for more recent systems of interest see Struckmeyer [31]). In the families we will study below the possible presence of x_0 occurs in the perturbing part.

Apart from the general case of Eq. (12), transformations non-based on the homogeneous formalism ($\quad = 1, \quad = Y_0 + \quad$), adopt the simpler formulation

$$H_0(x_n; Y_1; \dots; Y_{n-1}; \frac{\partial R}{\partial x_n}) = (Y_1; \dots; Y_n); \quad (13)$$

In the two types of Hamiltonian systems we study below, instead of $(x_0; x; X_0; X)$ we will start each Section considering briefly the two symplectic charts on which our study relies. In doing so we will use the standard notations, although there are more than one: Andoyer variables [3] for rotational motion and Whittaker (polar-nodal) variables [36] for orbital dynamics.

With respect to the new variables $(y_0; y; Y_0; Y)$, we will use $(d; \quad; v; u; D; \quad; \quad; U)$ for the rotational families and $(f; g; h; u; F; G; H; U)$ for the orbital ones. For the different sets within each family, we do not find necessary to distinguish among them, like using subindices, etc. Each one distinguishes itself by the expression taken by K_0 .

3 Rigid body transformations

Within the frame set up by Eqs (7) and (8) we will consider first canonical transformations derived from free rigid body Hamiltonian function. As it is well known, meanwhile Euler of variables continues to be the ones used in classical and recent books (see [18], [28]) at the same time in research papers is already customary to consider those systems formulated

in Andoyer variables. More precisely, the Hamiltonian of the free rigid body takes the form

$$H = H(\alpha; \beta; \gamma; M; N; a_i) = \frac{1}{2} (a_1 \sin^2 \alpha + a_2 \cos^2 \alpha) (M^2 + N^2) + \frac{1}{2} a_3 N^2; \quad (14)$$

where $0 < a_3 < a_2 < a_1$ are parameters. Then, according to our notation Eq. (1), we deal with the Hamiltonian

$$K = \frac{h}{T} + \frac{1}{2} (a_1 \sin^2 \alpha + a_2 \cos^2 \alpha) (M^2 + N^2) + \frac{1}{2} a_3 N^2 \quad (\alpha; \beta; \gamma; M; N; T); \quad (15)$$

where $(\alpha; \beta; \gamma; M; N)$ are Andoyer variables [3], and $M^2 + N^2$ making the Hamiltonian, (14), strictly positive and $\frac{1}{2} a_3 M^2 + H = \frac{1}{2} a_1 M^2$. As we said in what follows we consider only the generic case $a_3 < a_2 < a_1$. We do not need to give more details on those variables here. The interested reader may consult the classical note of Depierre [9] or in more geometric terms Fassò [13].

3.1 The general structure of the transformation

We look for canonical transformations

$$(\alpha; \beta; \gamma; t; M; N; T) \xrightarrow{T} (d; \gamma; u; D; \gamma; U) \quad (16)$$

that convert (15) in a certain function $\Phi(\gamma; \gamma; U)$ depending only on momenta. The transformation will be defined through a function $S = S(\alpha; \beta; \gamma; D; \gamma; U)$ in mixed variables such that

$$d = S_D; \quad \gamma = S_\gamma; \quad \gamma = S_\gamma; \quad u = S_U; \quad \gamma = S_\gamma; \quad M = S_M; \quad N = S_N; \quad T = S_T; \quad (17)$$

where we use the notation $S_x = \partial S / \partial x$. Thus, from (15) we set the H-J equation

$$\frac{h}{2} (a_1 \sin^2 \alpha + a_2 \cos^2 \alpha) S^2 + S^2 + \frac{1}{2} a_3 S^2 + S_T^2 \quad (\alpha; \beta; \gamma; S; S; S_T) = \Phi(\gamma; \gamma; U) \quad (18)$$

and S is chosen in separate variables as

$$S = U t + \gamma + W(\gamma; D; \gamma; U) \quad (19)$$

because t, γ , and γ are cyclic in (15). Then,

$$W = \int_0^\gamma Q(\gamma; D; \gamma; U) d\gamma; \quad (20)$$

where

$$Q = \frac{a_1 \sin^2 \alpha + a_2 \cos^2 \alpha}{a_1 \sin^2 \alpha + a_2 \cos^2 \alpha} \frac{A}{a_3}; \quad A = \frac{2}{2} - U^2; \quad (21)$$

and \overline{Q} must be real for all γ ; therefore, $\overline{Q} < (\frac{1}{2} a_2^{-2} + U)$.

If we assume that the regularizing factor depends only on the coordinate z , the transformations (17) are

$$d = \frac{1}{p} D I_3; \quad (22)$$

$$= \frac{1}{p} + I_1 + \frac{2U}{2} I_2 - \frac{2}{2} - \frac{1}{p} I_3; \quad (23)$$

$$= \frac{1}{p} + I_3; \quad (24)$$

$$u = t - \frac{1}{p} I_2 + \frac{1}{p} U I_3 \quad (25)$$

$$N = \frac{q}{Q}; \quad (26)$$

$$T = U; \quad \frac{1}{p} = \frac{1}{p}; \quad M = \frac{1}{p}; \quad (27)$$

where

$$I_1 = \int_0^z \frac{q}{Q} d; \quad I_2 = \int_0^z \frac{Q_A}{p Q} d; \quad I_3 = \int_0^z \frac{Q_A}{p Q} d; \quad (28)$$

3.2 The case $\mu = 1$

We only deal with transformations with $\mu = 1$. Therefore, $I_2 = I_3$.

Note that a restriction of Andoyer variables is $|N| \leq M$, which, because of (26), further constrains the values of the new Hamiltonian to be into the positive interval $2(\frac{1}{2}a_3^2; \frac{1}{2}a_2^2)$. In addition, one should be aware that the sign of $\frac{q}{Q}$ must be taken in accordance with the sign of N .

Again, the transformation depends on the integration of two quadratures, which closed form solution requires well known changes of variables. Thus, introducing the parameters²

$$m = \frac{a_1 - a_2}{a_1 - a_3}; \quad f = \frac{a_1 - a_2}{a_2 - a_3}; \quad (29)$$

and the functions $\mathcal{D} = \mathcal{D}(z; U; a_1; a_2; a_3)$, $m = m(z)$, defined by

$$\mathcal{D} = \frac{a_1 - a_2}{2(a_1 - a_3)U} z^2; \quad (30)$$

$$m = \frac{1}{1 - \frac{a_1 - a_2}{2(a_1 - a_3)U} z^2} = \frac{a_1 - a_2}{2(a_1 - a_3)U} \frac{z^2 a_3}{a_2 - a_3}; \quad (31)$$

the quadratures in (28) are solved to give

$$I_1 = \frac{m F(z) - (m + f) \mathcal{D}(z; f, m)}{f}; \quad (32)$$

$$I_2 = \frac{F(z)}{2(a_1 - a_3)U} : \quad (33)$$

² For the sake of linking with recent literature we used Fukushima's notation [17]. Depierre [12] calls a_3 , $\frac{0}{3}$, and $m = k_3^2$, and Sadov [32] names m and $f = \frac{1}{2}$.

where $F(\phi|m)$ is the elliptic integral of the first kind of modulus m which amplitude ϕ is unambiguously defined through

$$\cos \phi = \frac{\sin \psi}{\sqrt{1 - m \sin^2 \psi}}; \quad \sin \phi = \frac{\sqrt{1 - m \cos^2 \psi}}{\sqrt{1 - m \sin^2 \psi}}; \quad d\phi = \frac{\sqrt{1 - m \cos^2 \psi}}{\sqrt{1 - m \sin^2 \psi}} d\psi; \quad (34)$$

On the other hand, $(\phi; f|m)$ is the elliptic integral of the third kind of modulus m , characteristic f , and amplitude ϕ . Alternatively, one can write

$$I_1 = \frac{s}{f} (1 - m \sin^2 \phi) (\phi; \sqrt{m}); \quad (35)$$

where the new amplitude ϕ is unambiguously defined through

$$\cos \phi = \frac{\cos \psi}{\sqrt{1 - m \sin^2 \psi}}; \quad \sin \phi = \frac{\sqrt{1 - m \sin^2 \psi}}{\sqrt{1 - m \sin^2 \psi}}; \quad d\phi = \frac{\sqrt{1 - m \sin^2 \psi}}{\sqrt{1 - m \sin^2 \psi}} d\psi; \quad (36)$$

If we further introduce

$$= \frac{1}{\sqrt{a_1 a_2 a_3}} \frac{\sqrt{a_1 a_2}}{\sqrt{a_1 a_2 a_3}} = \frac{\sqrt{a_1 a_2}}{a_1 a_2} = \frac{1}{a_2 a_3} \frac{s}{f}; \quad (37)$$

then, the transformation equations above, (22)-(26), adopt the compact form

$$d = -\frac{1}{D} F(\phi; m) \quad (38)$$

$$= -\frac{h}{(a_3 - a_2)} F(\phi; m) + (a_1 - a_3) (\phi; f \sqrt{m}) \quad (39)$$

$$= -\frac{U}{2} F(\phi; m) - \frac{U}{2 a_2} (\phi; \sqrt{m}) \quad (40)$$

$$= \frac{1}{D} F(\phi; m) = \frac{1}{D} d \quad (41)$$

$$u = t + \frac{1}{D} F(\phi; m) = t + \frac{1}{D} d; \quad (42)$$

$$N = \frac{\sqrt{1 - m \cos^2 \psi}}{\sqrt{1 - m \sin^2 \psi}} = \frac{r}{1 - m \sin^2 \psi}; \quad (43)$$

where we maintain two expressions for N to ease comparisons with transformations in the literature. Remark that r remains arbitrary. Therefore, (37)-(43) define a family of canonical transformations that provide complete reduction of the Euler-Poinsot problem. Note, besides, from (42) that reduced Hamiltonians of the form $H = U + \frac{1}{D} F(\phi; m)$ are required for transformations preserving the time scale.

Despite the family of transformations above has been obtained from a specific dynamical system, the rigid-body problem, once it has been derived it may be applied to any problem we wish. However, successful transformations for different problems are closely dependent on the selection of r . Thus, for instance, if we are to apply one of the transformations of the family (38)-(43) above to reduce the rigid body dynamics or to study

a perturbed rigid body, we should consider specific facts. Because neither \mathbf{p} nor T are altered by the transformation it seems natural to choose $\mathbf{p} = U + (D; \mathbf{j})$. Then, u is the time, \mathbf{p} , U , and D , remain constant, and

$$\mathbf{p} = \mathbf{p}_0 + \mathbf{p}_1 u; \quad d = d_0 + D u: \quad (44)$$

Because the super-integrability of the Euler-Poinsot problem the closure of a generic trajectory is a two-torus [4], Therefore, \mathbf{p} should depend both on D and \mathbf{j} if we want to retain the topology of the problem. Specific selections of \mathbf{p} depending only on D whose appearance is mandatory for equation (38) to be defined will produce just periodic solutions. Even particular selections $\mathbf{p} = (D; \mathbf{j})$ may reduce the solutions to just periodic orbits. For instance, if we impose

$$\mathbf{p} = \mathbf{j} D; \quad (45)$$

i, j , integer, any computed $\mathbf{p} = U + (iD + j \mathbf{j})$ with arbitrary \mathbf{j} will provide only periodic solutions.

We discuss here some other possible choices of \mathbf{p} . Without any application in mind, one criteria might be 'to simplify' the coefficients of the variables in the transformation.

Thus, for instance, $\mathbf{p} = D$ in (37) implies

$$\mathbf{p} = U + \frac{D^2}{2} \mathbf{a}_1 - \frac{D^2}{a_2 a_3} \mathbf{j}; \quad (46)$$

where \mathbf{p} has dimensions of momentum and D of inverse of momentum of inertia. This Hamiltonian simplifies the transformation equations, (38)-(43), to

$$d = \frac{D^2}{a_2 a_3} F(\mathbf{j}; m) \quad (47)$$

$$= \frac{D^2}{a_2 a_3} + D \mathbf{a}_1 \cdot \mathbf{a}_2 \frac{D^2}{a_2 a_3} (\mathbf{j}; \mathbf{j}_n) \quad (48)$$

$$= \quad (49)$$

$$u = t; \quad (50)$$

$$N = D \frac{1}{\sqrt{t}} \frac{m \sin^2}{(a_1 a_3)(a_2 a_3)}; \quad (51)$$

To discuss other simplifications, we find it convenient to introduce

$$s_1 = \frac{U}{D}; \quad s_2 = 2 \frac{U}{D} - \frac{1}{D}; \quad s_3 = \frac{1}{D}; \quad (52)$$

and consider the possible choices of \mathbf{p} that make s_i ($i = 1; 2; 3$) independent of the momenta.

3.2.1 Case $s_1 = s_{1,0}$ and $s_3 = s_{3,0}$ constant.

Then, from the first and third of (52) and from (37), we get

$$s_{3,0} D = \frac{s_{3,0}}{s_{1,0}} p \frac{a_2}{a_3} \frac{q}{a_1^2} \frac{1}{2(D+U)}; \quad (53)$$

and, therefore

$$U = \frac{a_1^2}{2} - \frac{a_2 a_3}{2 s_{1,0}^2} [s_{3,0} + (D+U)]^2; \quad (54)$$

with arbitrary.

Because the topology of the problem, we can add the condition $s_{3,0} = 0$, and therefore $s_{3,0} = 0$. In addition, if we want to preserve the time scale (42), we restrict to the case

$$U = \frac{a_1^2}{2} - \frac{a_2 a_3}{2 s_{1,0}^2} (D)^2; \quad (55)$$

Sadov's transformation. The choice

$$s_{1,0} = \frac{s \frac{a_1 a_3}{a_2 a_3} D}{a_2 a_3}$$

gives

$$U = \frac{a_1^2}{2} - \frac{a_1 a_3}{a_1} \frac{D}{2} = \frac{a_1^2}{2} - \frac{a_1 a_3}{a_1} \frac{D^2}{2 + D^2}; \quad (56)$$

that is Sadov's [32] reduced Hamiltonian in which $D^2 = f$ and

$$D^2 = \frac{1}{D^2} - 1; \quad (57)$$

Kinoshita's case. If we instead choose

$$s_{1,0} = \frac{a_1 (a_1 + a_2 - 2a_3)}{(a_2 - a_3) (a_1 + a_2)} D; \quad (58)$$

we find

$$U = \frac{2a_1}{a_1 + a_2} \tilde{H}; \quad \tilde{H} = \frac{1}{2} \frac{a_1 + a_2}{2} D^2 + \frac{1}{b} D^2; \quad \frac{1}{b} = a_3 - \frac{1}{2} (a_1 + a_2) < 0; \quad (59)$$

where \tilde{H} is Kinoshita's reduced Hamiltonian [24, see Eq. (10') on p. 433]. Alternatively, the scaling may be avoided in our general formulation by choosing a regularizing factor $= 2a_1 = (a_1 + a_2)$ independent of the variables.

Thus, the major objection to Sadov's and Kinoshita's elements of being related to Andoyer canonical variables through implicit transformations is easily circumvented using the general transformation (37)–(43) above particularized for either Sadov's or Kinoshita's Hamiltonians, Eq. (56) and (59) respectively.

Fukushima's proposal. If we choose now

$$s_{1,0} = \frac{s}{a_2} \frac{a_1}{a_3} p^2 D^2; \quad (60)$$

we get Fukushima's proposal [17]

$$= U + \frac{1}{2} a_1 D^2; \quad (61)$$

that, depending on less than the required momenta, constrain the topology of the transformed Hamiltonian to periodic solutions only. Nevertheless, this way of proceeding may be perfectly adequate for a perturbation theory, as shown in [37].

TR-type mapping. In a similar way to Fukushima, we propose to choose

$$s_{1,0} = \frac{s}{a_2} \frac{a_1}{a_3} (D); \quad (62)$$

leading to

$$U = a_1 D + \frac{1}{2} D^2 \quad (63)$$

that is formally equal to the reduced Hamiltonian of the TR-mapping [34, 11] given in Eq. (125) below. It is quadratic, depends on two momenta — as it should be for the more general case of a super-integrable three degrees of freedom problem with four independent integrals [4] — and the momenta keep the dimensions of the original problem because the Hamiltonian is multiplied by the inverse of the momentum of inertia a_1 .

Then, the reduced system has four elements $(\dot{q}; D; \dot{p}; \dot{q})$ and two variables that evolve linearly with time

$$q = q_0 + a_1 D t; \quad d = d_0 + a_1 (D) t;$$

the condition $D = \text{rational}$ providing the subset of periodic solutions.

3.2.2 Case $s_2 = s_{2,0}$ and $s_3 = s_{3,0}$ constant

From the second and third of (52) we get

$$s_{2,0} D = \frac{s_{2,0}}{s_{3,0}} U = 2 \frac{U}{D} \quad (64)$$

where $s_{3,0}$ is arbitrary.

A simple choice satisfying the condition in (64) could be $U = U + \frac{1}{2} D^2 (s_{2,0})$, for instance. Particular cases in the literature make $s_{2,0} = 0$. Thus, Depierre and Elie [12] choose

$$U = \frac{1}{2} D^2; \quad (65)$$

Other solutions could be, for instance,

$$U = \frac{1}{2} D^2; \quad U = \frac{1}{2} D^2; \quad (66)$$

4 Orbital Canonical transformations

What we have said at the beginning of the previous Section for canonical variables in rotational dynamics, it appears again in orbital variables. Thus, meanwhile Hamiltonian particle dynamics still happens to be introduced in spherical canonical variables (see for instance [6], [18], [23], [35]), at the same time in space research another canonical set of variables is widely in use; we refer to nodal-polar variables, already considered for planetary theories almost a century ago [22]. The main difference of both set of variables is that nodal-polar variables carries out a double reduction (axial and central symmetries) of Kepler type systems, meanwhile spherical variables only deals with the axial symmetry.

As in the search of orbital canonical transformations we build up our H-J equation based on nodal-polar variables, we briefly give a description about them. Taking a reference frame $(O; e_1; e_2; e_3)$, they are introduced in a natural way associated with the "instantaneous plane of motion" whose characteristic vector is $x \times X$. The nodal-polar variables $(r; \theta; \phi)$ are defined as follows: the variable r is $|x|$; the angle θ called the ascending node, is defined by $\cos \theta = \frac{e_1 \cdot (x \times X)}{|x \times X|}$ and $\sin \theta = \frac{e_2 \cdot (x \times X)}{|x \times X|}$, where \hat{n} is the unit vector defined by $\hat{n} = \frac{x \times X}{|x \times X|}$; the variable ϕ is a polar angle in the plane of motion giving the position of x reckoned from the vector \hat{n} . The variables $(R; N; T)$ are the corresponding conjugate momenta. For more details and explicit expressions of the transformation $(x; y; z; X; Y; Z) \rightarrow (r; \theta; \phi; R; N; T)$, called by some authors Whittaker transformation, see for instance [10].

4.1 The general structure of the transformations

Given the function

$$H = \frac{1}{2} R^2 + \frac{1}{r^2} + V(r) + T(r; \theta; \phi; R; N; T) \quad (67)$$

where $(r; \theta; \phi; R; N; T)$ are Hill or polar-nodal variables in the extended phase space and V only depends on distance, we look for a canonical transformation

$$(r; \theta; \phi; R; N; T) \rightarrow (f; g; h; u; F; G; H; U)$$

that converts (67) in a certain function $\mathcal{H}(F; G; H; U)$ depending only on the momenta.

The transformation will be defined through a generating function in mixed variables $S = S(r; \theta; \phi; F; G; H; U)$ such that

$$f = S_F; \quad g = S_G; \quad h = S_H; \quad u = S_U; \quad R = S_R; \quad \theta = S_\theta; \quad N = S_N; \quad T = S_T \quad (68)$$

Then, from (67) we set the H-J equation

$$\frac{1}{2} S_R^2 + \frac{1}{r^2} + V(r) + S_T(r; S_\theta; S_N; S_T) = \mathcal{H}(F; G; H; U) \quad (69)$$

Because t , ϕ , and θ are not present in (67), the generating function may be chosen in separate variables

$$S = U t + H(\phi) + G(\theta) + W(r; F; G; H; U) \quad (70)$$

Then,

$$\frac{1}{2} W_r^2 + \frac{G^2}{r^2} + V(r) + U = \frac{(F; G; H; U)}{(r; G; H; U)} \quad (71)$$

that can be solved for W by a simple quadrature:

$$W = \int_{r_0}^r \frac{dr}{Q(r; F; G; H; U)} \quad (72)$$

where

$$Q = 2 \frac{(F; G; H; U)}{(r; G; H; U)} - 2V(r) - \frac{G^2}{r^2} - 2U = 0 \quad (73)$$

Therefore, the transformation is

$$f = F I_3; \quad (74)$$

$$g = G I_1 + \int_{r_0}^r \frac{G}{2Q} dr; \quad (75)$$

$$h = H I_3 + \int_{r_0}^r \frac{H}{2Q} dr; \quad (76)$$

$$u = U I_2 + \int_{r_0}^r \frac{U}{2Q} dr; \quad (77)$$

$$R = W_r = \frac{1}{Q}; \quad (78)$$

$$= G; \quad N = H; \quad T = U; \quad (79)$$

where

$$I_1 = \int_{r_0}^r \frac{1}{Q} dr; \quad I_2 = \int_{r_0}^r \frac{dr}{Q}; \quad I_3 = \int_{r_0}^r \frac{dr}{Q}; \quad (80)$$

The case $\beta = (r)$. Transformations are clearly simplified when we choose the regularizing factor independent of the momenta G, H , and U . Thus, if $\beta = (r)$, we have

$$G = H = U = 0; \quad (81)$$

and the rightmost terms on the right side of (74)-(77) vanish, which turn these equations into

$$f = F I_3; \quad (82)$$

$$g = G I_1 + \int_{r_0}^r \frac{G}{2Q} dr; \quad (83)$$

$$h = H I_3; \quad (84)$$

$$u = U I_2 + \int_{r_0}^r \frac{U}{2Q} dr; \quad (85)$$

Note that $\beta = 1$ and $\beta = r^2$ make $I_2 = I_3$ and $I_3 = I_1$, respectively. We plan to discuss these choices in detail below, focusing on a two-parametric family $V = V(r; \beta)$ of radial potentials.

4.2 Families of the isochronal potential

In what follows we only deal with the transformation (82)–(85) and, in addition, limit our study to the specific case of Henon's isochronal potential [21, 6]

$$V = \frac{p}{b + \sqrt{b^2 + r^2}}; \quad (86)$$

where b and p are parameters. A relevant, particular case of the isochronal is the Keplerian potential $V = -r$ that occurs for $b = 0$.

For the isochronal potential (86) we find convenient to write (73) as

$$Q = \frac{r^2 + b^2}{r^2} Q = 0; \quad Q = \frac{p}{r^2 + b^2} \sqrt{\frac{2}{r^2 + b^2} + \frac{1}{a}}; \quad (87)$$

where r , p , and a , are certain functions of the momenta and parameters, which only will be determined after the regularizing parameter ϵ has been chosen. Then,

$$Q = p \left(\frac{1}{s} - \frac{1}{s_1} - \frac{1}{s_2} - \frac{1}{s} \right); \quad (88)$$

where s_1 ($s = \sqrt{\frac{p}{r^2 + b^2}}$) s_2 are the two possible roots of the conic $Q = 0$:

$$s_{1,2} = \frac{p}{1 \pm \frac{p}{a}} = \frac{a(1 \pm e^2)}{1 \pm e} = a(1 \pm e); \quad e^2 = 1 - \frac{p}{a} < 1; \quad (89)$$

4.2.1 Introducing auxiliary variables

The roots $s_{1,2}$, the extreme values of s , make natural the introduction of the auxiliary variables θ and ϕ , defined through

$$\sqrt{\frac{p}{r^2 + b^2}} = a(1 - e \cos \theta); \quad \sqrt{\frac{p}{r^2 + b^2}} = a \sin \phi; \quad (90)$$

$$\sqrt{\frac{p}{r^2 + b^2}} = \frac{p}{1 + e \cos \theta}; \quad \sqrt{\frac{1}{r^2 + b^2}} = \frac{e}{p} \sin \phi; \quad (91)$$

in order to ease the solution of quadratures in (80), which we find convenient to write

$$I_1 = \int_{r_0}^r \frac{r^2 + b^2}{r^2} \sqrt{\frac{1}{r^2 + b^2}} d\sqrt{\frac{p}{r^2 + b^2}}; \quad (92)$$

$$I_2 = \int_{r_0}^r \frac{1}{\sqrt{Q}} d\sqrt{\frac{p}{r^2 + b^2}}; \quad (93)$$

$$I_3 = \int_{r_0}^r \frac{r^2 + b^2}{\sqrt{Q}} d\sqrt{\frac{1}{r^2 + b^2}} = \int_{r_0}^r \frac{1}{\sqrt{Q}} d\sqrt{\frac{p}{r^2 + b^2}}; \quad (94)$$

For given values of the regularizing factor ϵ , the quadratures above, (92)–(94), may be integrated in closed form without need of specifying the new Hamiltonian, which remains as a formal function of the momenta $\mathbf{p} = (F; G; H; U)$.

Details on the families of canonical transformations generated by the cases $\epsilon = 1$, $\epsilon = b + \sqrt{b^2 + r^2}$, and $\epsilon = r^2$ are given below. Besides, for the Keplerian case $b = 0$ we recover Delaunay's [8], Levi-Civita's [25, 26], and Hill's [22] canonical transformations, as particular examples of our families. An elegant alternative for the derivation of these historic transformations may be found in [2].

4.3 Case I: Delaunay's family of transformations $\epsilon = 1$.

From (73) and (87) we obtain

$$\epsilon = 1; \quad a = \frac{G^2}{2(U)}; \quad p = \frac{G^2}{2} + 2b \frac{b^2}{a}; \quad (95)$$

Then, introducing the change of (90) in (93), I_2 is easily solved to give

$$I_2 = I_3 = \frac{1}{8(U)^3} (e \sin \varphi); \quad (96)$$

Analogously, the change of (91) is introduced in (92) to solve the quadrature I_1 . We obtain

$$I_1 = \frac{1}{2G} \frac{p}{G^2 + 4b}; \quad (97)$$

where we have introduced two new auxiliary variables φ_1, φ_2 , defined by means of the trigonometric relations [37]

$$\tan \frac{\varphi_1}{2} = \frac{\sqrt{\frac{1+e-b/a}{1-e-b/a}} \tan \frac{\varphi}{2}}{\frac{1+e-b/a}{1-e-b/a}}; \quad (98)$$

$$\tan \frac{\varphi_2}{2} = \frac{\sqrt{\frac{1+e+b/a}{1-e+b/a}} \tan \frac{\varphi}{2}}{\frac{1+e+b/a}{1-e+b/a}}; \quad (99)$$

Substitution of I_1 and $I_2 = I_3$, given by (97) and (96), respectively, in (82)-(85) provides the family of canonical transformations of the isochronal potential to which we attach the name of Delaunay.

Among the variety of possible choices of the new Hamiltonian, a simplifying option is to take $H = U + (F/G; H; b)$, that makes u to remain as the original time. In addition, the choice $H = 0, F = G = 1/(2)^{3/2}$, may be solved for φ , to give

$$\varphi = U \frac{1}{2} \frac{2}{(F+G)^2} \quad (100)$$

that maximally simplifies the transformation equations:

$$f = e \sin \varphi; \quad (101)$$

$$g = \frac{1}{2} \frac{G}{G^2 + 4b} \frac{2}{2} + f; \quad (102)$$

$$h = \varphi; \quad (103)$$

$$u = t; \quad (104)$$

while retaining the topology of the original Hamiltonian. Delaunay's selection $H = 0, G = 0$ leads to

$$\varphi = U \frac{2}{2F^2} \quad (105)$$

that depends on fewer momenta and, therefore, constrains the topology of the original system to periodic solutions only | which may be adequate for a perturbation study like [37].

In the Keplerian case $b = 0$, $V = -r$, the selection of the new Hamiltonian $= (U; F)$ does not constrain the range of solutions and the specific selection of (105) provides the popular Delaunay transformation that, taking into account that $\alpha_1 = \alpha_2 = 0$, is

$$f = e \sin \varphi; \quad g = \varphi; \quad h = \varphi; \quad u = t; \quad (106)$$

where the most extended notation writes $L = F, \quad \alpha = f$.

4.4 Case II: Levi-Civita's family of transformations $= b + \sqrt{b^2 + r^2}$

Now, we write

$$= +; \quad a = \frac{+}{2U}; \quad p = \frac{G^2}{+} + 2b \frac{b^2}{a}; \quad (107)$$

The change (91) is used to integrate I_1 , (92), and the change (90) to integrate I_2 and I_3 , (93) and (94). We get

$$I_1 = \frac{1}{2G} \sqrt{\frac{2}{G^2 + 4b(+)}}; \quad (108)$$

$$I_2 = \sqrt{\frac{+}{8U^3}} (e \sin \varphi); \quad (109)$$

$$I_3 = \sqrt{\frac{b^2}{2U}} \sqrt{\frac{b^2}{G^2 + 4b(+)}}; \quad (110)$$

where α_1, α_2 , are the same auxiliary variables defined in (98) and (99) respectively. Then, substitution of I_1, I_2, I_3 , (108)-(110), in (82)-(85) provides the family of canonical transformations of the isochronal potential to which we tie the name of Levi-Civita.

The, Keplerian case $b = 0$, $\alpha_1 = \alpha_2 = 0$, gives

$$f = \sqrt{\frac{1}{2U}}; \quad (111)$$

$$g = + \sqrt{\frac{1}{2U}}; \quad (112)$$

$$h = + \sqrt{\frac{1}{2U}}; \quad (113)$$

$$u = t \sqrt{\frac{+}{8U^3}} (e \sin \varphi) + \sqrt{\frac{1}{2U}}; \quad (114)$$

The canonical transformation still remains undefined. We note in (111) that $f = \sqrt{\frac{1}{2U}}$, a partial differential equation that may be solved to give

$$= \sqrt{\frac{h}{2U}} F + C_1(G; H; U); \quad (115)$$

where C_1 is an arbitrary function of G, H , and U . The simple choice $C_1 = 1$ provides the "first" Levi-Civita [25] transformation³

$$f = \frac{1}{2} \arctan \left(\frac{h}{g} \right); \quad g = \frac{1}{2} \left(\frac{p}{U} - \frac{h}{g} \right); \quad h = \frac{1}{2} \left(\frac{p}{U} + \frac{h}{g} \right); \quad u = t - \frac{p}{8U^3} (f - e \sin f); \quad (116)$$

Other possibility is to force $\frac{p}{U} = (1 + e) = (2U)$ so that only periodic oscillations are introduced in the time scale (114). Then,

$$\frac{p}{U} = 2U + C_2(G; H); \quad (117)$$

with C_2 arbitrary. The combination of both conditions (115) and (117) leads to

$$\frac{p}{2U} = F + \frac{p}{2U} + C(G; H); \quad (118)$$

with C arbitrary. If we further take $C = 0$, we get

$$f = \frac{1}{2} \arctan \left(\frac{h}{g} \right); \quad g = \frac{1}{2} \left(\frac{p}{U} - \frac{h}{g} \right); \quad h = \frac{1}{2} \left(\frac{p}{U} + \frac{h}{g} \right); \quad u = t + \frac{F}{2U} e \sin f; \quad (119)$$

that is the famous "second" Levi-Civita [26] transformation.

4.5 Case III: Hill's family of transformations: $\frac{p}{U} = r^2$

We get now

$$\frac{p}{U} = r^2; \quad a = \frac{G^2}{2U}; \quad p = \frac{G^2}{2} + 2b \frac{b^2}{a}; \quad (120)$$

The required quadratures are solved with the same changes of variables as before, giving

$$I_1 = I_3 = \frac{p}{2} \frac{1}{G^2} \frac{1}{2}; \quad \frac{p}{2} \frac{2}{G^2} \frac{2}{2 + 4b}; \quad (121)$$

$$I_2 = \frac{p}{8U^3} (f - e \sin f); \quad (122)$$

which substitution in (82)-(85) provides the family of canonical transformations of the isochronal potential to which we assign the name of Hill.

Focusing on the Keplerian case $b = 0$, $\frac{p}{U} = \frac{1}{2}$, a simplification choice that makes $f = 0$ is to select $\frac{p}{U}$ in such a way that $\frac{p}{U} I_1 = 1$, then

$$\frac{1}{2} G^2 = (F + C)^2; \quad (123)$$

where $C = C(G; H; U)$ is an arbitrary function. Then,

$$F = (F + C); \quad G = G (F + C) C_G; \quad H = (F + C) C_H; \quad U = (F + C) C_U; \quad (124)$$

³Note that Levi-Civita proceeds by scaling r by $1 = \frac{p}{2U}$, or choosing $\frac{p}{U} = r^2$ in our notation. This is equivalent to scaling our new Hamiltonian by the same factor, and yields to Levi-Civita's original Hamiltonian $H = F$ while the transformation (116) remain identical

The trivial choice $C = 0$ makes $g = \frac{1}{2}F$, $h = \frac{1}{2}G$, and $u = t - \frac{1}{8U^3}(\frac{1}{2}F \sin f)$.

Choosing C as a linear combination of them om enta, simply ad s to the angle variables. Speci ally, when we set $C = -\frac{1}{8U^3}G$ we obtain⁴

$$h = \frac{1}{2}F - \frac{1}{8U^3}G \quad (125)$$

and recover the TR-m apping [34, 11]

$$f = \frac{1}{2}F; \quad g = \frac{1}{2}G; \quad h = \frac{1}{2}F - \frac{1}{8U^3}G; \quad u = t - \frac{1}{8U^3}(\frac{1}{2}F \sin f): \quad (126)$$

5 Conclusions

A deep insight in the computation of canonical transformations is obtained through a general formulation of the Hamilton-Jacobi equation in the extended phase space that, besides, includes a regularizing function. As far as the transformed Hamiltonian may remain formal, one can obtain families of canonical transformations instead of single sets of canonical variables. Then, particular transformations that meet the user's required characteristics are systematically computed from the solution of partial differential equations that may enjoy very simple solutions.

Our formulation shows that different transformations in the literature belong to the same families. Furthermore, it reveals fundamental features of the transformations. Thus, among the worked examples, we identify families having as members known canonical changes of variables that originally were defined through implicit relations.

Acknowledgements

Partial support is recognized from projects MTM 2006-06961 (S.F.) and AYA 2009-11896 (M.L.) of the Government of Spain, and a grant from Fundacion Seneca of the autonomous region of Murcia.

References

- [1] Abraham, R., Marsden, J.E., 1985, *Foundations of Mechanics*, Perseus Books, Cambridge Mass.
- [2] Andoyer, M.H., 1913, "Sur l'Anomalie Excentrique et l'Anomalie Vraie comme Elements Canoniques du Mouvement Elliptique d'apres M.M. T. Levi-Civita et G.W. Hill," *Bulletin astronomique*, Vol. 30, pp. 425-429.
- [3] Andoyer, M.H., 1923, *Cours de Mecanique Celeste*, t. 1, Gauthier-Villars et cie, Paris, p. 57.

⁴Note that Hill proceeds by scaling r by $1 = \frac{1}{2}G - \frac{1}{2}F$, or choosing $r^2 = (G - \frac{1}{2}F)$ in our notation. This is equivalent to scaling our new Hamiltonian by $G - \frac{1}{2}F$, what yields to Hill's Hamiltonian $H = F$ while the transformation (126) remains unaltered.

- [4] Benettin, G., 2004, "The elements of Hamiltonian perturbation theory," in *Hamiltonian Systems and Fourier Analysis: New Prospects for Gravitational Dynamics*, D. Benest and C. Froeschle (ed.), pp. 1{98. Cambridge Scientific Publ., Cambridge, UK.
- [5] Blanes, S., Budd, C., 2004, "Explicit Adaptive Symplectic (Easy) Integrators: A Scaling Invariant Generalisation of the Levi-Civita and KS Regularisations," *Celestial Mechanics and Dynamical Astronomy*, Vol. 89, No. 4, pp. 383{405.
- [6] Boccaletti, D., Pucacco, G., 2001, *Theory of Orbits. 1: Integrable Systems and Non-perturbative Methods*, Springer-Verlag, Berlin Heidelberg, pp. 316 and .
- [7] Chang D. E. and Marsden, J. E., 2003, "Geometric derivation of the Delaunay variables and geometric phases," *Celest Mech & Dyn Astron* 86: 185{208.
- [8] Delaunay, Ch E., 1867, "Theorie du Mouvement de la Lune," *Memoires de l'Academie des Sciences de l'Institut Imperial de France*, Vol. 28, See Chap. 1, para. 5, pp. 9{11.
- [9] Deprit, A., 1967, "Free Rotation of a Rigid Body Studied in the Phase Space," *American J Physics* 35, 424{428.
- [10] Deprit, A. and Rom, A., 1970, *Celest Mech.* 2, 166.
- [11] Deprit, A., 1981, "A Note Concerning the TR-T Transformation," *Celestial Mechanics*, Vol. 23, pp. 299{305.
- [12] Deprit, A., 1993, "Complete Reduction of the Euler-Poinsot Problem," *Journal of the Astronautical Sciences*, Vol. 41, No. 4, pp. 603{628.
- [13] Fassò, F., 1996, "The Euler-Poinsot top: a non-commutatively integrable system without global action-angle coordinates," *J. Appl. Math. Phys. (ZAMP)* 47, 953{976.
- [14] Fassò, F., 2005, "Superintegrable Hamiltonian Systems: Geometry and Perturbations," *Acta Applicandae Mathematicae*, Vol. 87, pp. 93{121.
- [15] Ferrer, S., Lara, M., 2009, "Orbital Canonical Transformations. The Isochronal Family," *XII Jornadas Mecanica Celeste*, Laln, Spain 1-3 July 2009.
- [16] Fukushima, T., 2008, "Simple, Regular, and Efficient Numerical Integration of Rotational Motion," *The Astronomical Journal*, Vol. 135, pp. 2298{2322.
- [17] Fukushima, T., 2008, "Canonical and Universal Elements of the Rotational Motion of a Triaxial Rigid Body," *The Astronomical Journal*, Vol. 136, pp. 1728{1735.
- [18] Goldstein, H., Poole, C., and Safko, J., 2002, *Classical Mechanics*, 3rd edition, Pearson Addison-Wesley.
- [19] Grandati, Y., Berard, A., and Mohrbach, H., 2008, "Bohlin-Arnold-Vassiliev's duality and conserved quantities," *arXiv:0803.2610v2 [math-ph]*

- [20] Gur l, P ., 2005, Canonical Formalism for Modelling and Control of Rigid Body Dynamics, in New Trends in Astrodynamics and Applications, Ann. N.Y. Acad. Sci., Vol. 1065: 391{413.
- [21] Henon, M ., 1959, \L'am as isochrone I," Annales d'A strophysique, Vol. 22, pp. 126{39.
- [22] Hill, G .W ., 1913, \Motion of a system of m aterial points under the action of gravitation," Astronom ical Journal, Vol. 27, iss. 646{647, pp. 171{182.
- [23] Classical M echanics,.....
- [24] Kinoshita, H ., 1972, \First-Order Perturbations of the Two Finite Body Problem ," Publications of the Astronom ical Society of Japan, Vol. 24, pp. 423{457.
- [25] Levi-C ivita, T ., 1913, \Nouvo sistem a canonico di elem enti ellittici," Annali di M atem atica, serie III, Vol. XX , pp. 153{170.
- [26] Levi-C ivita, T ., \Sur la regularization du problem e des trois corps," Acta m athem atica, Vol. 42, 1918, pp. 99{144.
- [27] Lum , K -Y , Bloch, A M ., 1999, Generalized Serret-Andoyer Transform ation and Applications for the Controlled Rigid Body, Dynamics and Control, Vol. 9, 39{66.
- [28] Marsden, J.E ., and Ratiu, T . 1999, Introduction to M echanics and Sym m etry, TAM 17, 2nd Ed, Springer, New York.
- [29] Ortega, J. P. and Ratiu, T . 2004, M om entum M aps and Ham iltonian Reduction, Birkhauser, Boston.
- [30] Poincare, H ., 1893, Les m ethodes m ouvelles de la m ecanique celeste, Vol. 2, Gauthier-V illars et F ils, Paris, pp. 315{342.
- [31] Struckmeier, J., 2005, \Ham iltonian Dynamics on the sym plectic extended phase space for autonom ous and non-autonom ous system s," Journal of Physics A : M athem atical and General, Vol. 38, No. 6, pp. 1257{1278.
- [32] Sadov, Yu. A ., 1970, \The Action-A ngles Variables in the Euler-Poinsot Problem ," Prikladnaya M atem atika i M ekhanika, Vol. 34, pp. 962-964. [Journal of Applied M athem atics and M echanics (English translation), Vol. 34, No. 5, 1970, pp. 922{925]
- [33] Sadov, Yu. A ., 1970, \The Action-A ngle Variables in the Euler-Poinsot Problem ," Preprint No. 22 of the Keldysh Institute of Applied M athem atics of the A cadem y of Sciences of the U SSR , M oscow (in Russian).
- [34] Scheifele, G ., 1970, \Generalisation des elem ents de Delaunay en m ecanique celeste. Application au m ouvem ent d'un satellite arti ciel", Les Com ptes rendus de l'A cadem ie des sciences de Paris, Ser. A , Tom e 271, pp. 729{732.
- [35] Sussman, G .J., Wisdom , J., 2001, Structure and Interpretation of Classical M echanics, The M IT Press, Cam bridge, pp. 403 and ..

- [36] Whittaker, E.T., A treatise on the analytical dynamics of particles and rigid bodies, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1988, Reprint of the 1937 edition.
- [37] Yanguas, P., 2001, "Perturbations of the isochrone model," Nonlinearity, Vol. 14, pp. 1{34.