M asanori Iye¹

ABSTRACT

N ational A stronom ical O bservatory is an inter-university institute serving as the national center for ground based astronom y o ering observational facilities covering the optical, infrared, radio wavelength dom ain. NAOJ also has solarphysics and geo-lunar science groups collaborating with JAXA for space m issions and a theoretical group with computer simulation facilities. The outline of NAOJ, its various unique facilities, and some highlights of recent science achievem ents are reviewed.

Subject headings:

1. The Outline of NAOJ

The National Astronom ical Observatory of Japan (NAOJ), as the national center of astronom ical researches in Japan, deploys ve observatories and three VERA stations in Japan, Subaru Telescope in Hawaii, and ALMA Observatory in Chile as shown in Fig.1. NAOJ, belonging to the National Institutes for Natural Sciences (NINS) as one of the ve inter-university institutes, o ers its various top-level research facilities for researchers in the world.

M a jor facilities of NAOJ include 8.2m Subaru Telescope at M auna K ea, H aw aii, 45m R adio Telescope and R adio H eliograph at N obeyam a, 1.8m telescope at O kayam a A strophysical O bservatory, VERA interferom eter for G alactic radio astrom etry, solar facilities at M itaka and N orikura, super com puting facility and others. In addition to these ground based facilities, NAOJ scientists have been corem em bers for som e of the ISAS spacem issions, H inode, VSOP, and K aguya.

The total number of NAOJ sta in 2007 amounts to 258, including 33 professors, 48 associate professors, and 82 research associates. In addition, about 200 contract sta s are supporting the daily activities at each campus of NAOJ. A total of 72 graduate students and 34 post doctoral fellows are a liated to NAOJ.



Fig. 1. NAOJ's ve observatory cam puses and three VERA stations. Also shown are Subaru Telescope facilities in Hawaii and ALMA facility under construction in Chile.

¹O pticaland Infrared A stronom y D ivision, N ationalA stronom icalObservatory, M itaka, Tokyo, 181-8588 Japan

The annual total budget of NAOJ delivered from the M inistry of E ducation, Science, Culture and Sports, MEXT, for the scal year 2009 was 15.4 B including 3.3 B for personnelexpenses, 8.1 B for research expenses, and 3.5 B for facility developm ent expenses. In addition, 66 com – petitive research funds, am ounting 0.6 B¥ were acquired by applications of individual researchers in FY 2008.

All the NAOJ facilities are o ered for open use for scientists in Japan and some are o ered for world wide community. The selection process of successful proposals is based on the refereeing system for each facility and these processes are overseen by dedicated committees. In the scal year 2007, a total of 354 proposals were approved and a total of 1300 researchers, including 200 foreigners, used NAOJ facilities for researches.

O bservational and theoretical research achievem ents spanning the solar system, stars and clusters, interstellar m atter, galaxies, active galactic nuclei, and cosm ology were published in 376 refereed papers and 603 proceedings papers in English in FY 2007.

NAOJ, like all national universities and interuniversity institutes, was reformed to be a sem igovernmental agency back in 2004, according to the structural reformation scheme of the government. The science achievements of NAOJ are highly respected and the government continues its nancial supports, but this transformation in the academic system generated some turbulence. NAOJ regards, however, its increased exibility as a merit to plan for the better future taking the recommendations by external panel reviews.

NAOJ pays much e ort also to promote public relations. NAOJ opens its 4D 2U dome theater, inaugurated in 2007, as a new facility to show original numerical simulation videos in three dimensions using stereographic technology. NAOJ home page is one of the most popular site among the research institutes and universities in Japan. More than million accesses are observed every month, regularly.

2. Subaru Telescope

The 8.2m Subaru Telescope (Fig 2.), completed in 1999 atop the sum m it of 4200-m eter M auna K ea on the island of H aw ai'i, provides astronom ers of



Fig. 2. 8m Subaru Telescope has four foci and 8 open-use scienti c instrum ents.



Fig. 3. New laser guide star adaptive optics system will enhance the spatial resolution of Subaru Telescope by an order of magnitude.

Japan and around the world with opportunities to carry out top-level observations in the optical to m id-infrared (0.3-30 m) dom ain. O bservational targets range widely, e.g., objects in our Solar system, putative planets around nearby stars, galaxies and clusters of galaxies, even to the furthest objects at the edge of the observable universe. In order to address this wide range of research interests, Subaru o ers a suite of eight high perform ance cam eras and spectrographs in the optical and infrared dom ain.

Continuous R&D e orts to design and build new instrum ents are made possible by winning signi cant science budgets. A new adaptive optics system, AO 188, was constructed in 2006 to provide di raction limited imaging capability in the near infrared bands with Strehl ration as high as 0.6. This system, together with a new ly developed laser-guide star generating system (Fig.3.), which creates an articial reference star in the upper atm osphere to operate the A0188 anywhere in the sky, will greatly enhance the Subaru capability from 2009. New instruments, HiCIAO, the successor to the current stellar coronagraph C IAO, to be used with AO 188, Fiber Multi-O bject Spectrograph, FMOS, with 400 ber feed, and the HyperSuprim eC am, the successor of im ager Suprim e-Cam, providing 10 times larger eld coverage, will be on line shortly.

NAOJ set up Extrem ely Large Telescope (ELT) project o ce in 2005 and hopes to establish an international partner ship to realize the Thirty M eter Telescope (TMT) project as a reality at M auna K ea by around 2018.

Subaru Telescope has been yielding m any science results, discovery of the m ost distant galaxy at redshift 6.96 (Iye et al. 2006) as shown in Fig.4, led to the new insight on the cosm ic dawn of reionization of the universe. It is noteworthy that nine from the top-ten m ost distant galaxies have been discovered by the Subaru Telescope as of M arch 2009. Subaru also successfully m apped the distribution of dark m atter through weak lensing analysis.

Chem ical abundance studies of old stars in the G alaxy are other areas, where Subaru m ade significant science achievem ents. F ig 5 shows the m ost m etal de cient star A okiet al. 2006) discovered by Subaru H igh D ispersion Spectrograph, HDS.



Fig. 4. | Them ost distant galaxy, IOK-1, at 12.88 billion light years away.



Fig. 5. High dispersion spectra of the Sun (above) and the most metal de cient star ever found, HE1327-2326 (below). This star is likely to be the rst generations of stars in the universe.



Fig. 6. Peculiar streak of HII gas found in the Com a Cluster

A nother new surprise brought by Subaru's Suprim e-C am camera is the discovery of an unusual streak of ionized hydrogen gas associated with a galaxy 300 m illion light-years from Earth (Fig.6) (Yagi et al. 2007). The lament of gas is only 6 thousand light-years wide, yet extends 200,000 light-years, about the distance between the M ilky W ay Galaxy and its companion, the Large M agellanic C loud. Finding such an extrem ely narrow and long ionized gas cloud is som ething quite unexpected.



Fig. 7. Planet HR 8799b as in aged by Subaru in 2002.

Fig. 7 shows the image of the outermost planet HR 8799b as con med in the image taken by Subaru Telescope back in 2002. The data were re-examined stimulated by the discovery paper of three planets around HR 8799 in 2008 (M arois, et al. 2008). This m easurement is used to the determ ination of the orbit of this planet.

3. A tacam a Large M illim eter/subm illim eter A rray (A LM A)

ALMA is an international radio astronom y facility that is under construction on the Chilean plateau at an altitude of 5,000m above sea level. ALMA is a partnership between Japan, Europe and North America in cooperation with the Republic of Chile. NAOJ is leading the construction and operation of ALM A on behalf of the Japanese and Taiwanese science communities. ALMA consists of an array of sixty-eight 12-m antennas and twelve 7-m antennas extending over 18.5km span at maximum, allowing high spatial resolution equivalent to that of a telescope of the same size. NAOJ is building Atakam a Compact Array as part of the ALM A project (Fig.8). The resolution of ALM A will be 10 tim es as high as that of the world's most sophisticated optical telescopes, such as Subaru. With such a high resolution one can recognize a coin from a distance of 400km. ALMA aims to explore the dark and invisible universe by detecting m illim eter/subm illim eter waves between radio wave and far-infrared spectral regions in order to gure out the mysteries on the birth of stars and planets, as well as the synthesis of organic molecules in the universe. A LM A is scheduled to start its full operation from 2012.



Fig. 8. Four 12m anntenas for A tacam a C om – pact A rray (ACA) as delivered to C hile.

4. N obeyam a R adio O bservatory

The Nobeyama Radio Observatory (NRO) has two large radio telescopes, the Nobeyam a 45m telescope and the Nobeyam a Millim eter A rray (NMA). The 45m telescope has been the largest radio telescope operated at millim eter wavelengths since its commissioning in 1982. The advantage of the 45m telescope is its high sensitivity. NMA consists of six 10m movable parabola antennas and provides high angular resolution, 10 times better than that of the 45m telescope. They have benn widely used for various observations of starform ing regions in our galaxy to distant galaxies, how ever NMA ceased its science operation since FY 2008.

The discovery of the kinematical evidence of a massive black hole at the center of a galaxy NGC 4258 by VLBA in NRAO/USA as shown in Fig.9 (M iyoshiet al. 1995) was based on the previous discovery of very high velocity water vapor m asers by the 45m telescope.

ASTE (Atacam a Submillim eter Telescope Experim ent) is the rst 10-m subm illim eter telescope in the southern hem isphere located 4800 m above sea level at the Atacam a desert in Chile and operated by NRO/NAOJ and University of Tokyo in collaboration with the University of Chile etc. One of the important purposes of ASTE is exploring the southern sky at submillimeter wavelengths from 1 mm down to 0.3 mm (up to 950 GHz). ASTE equipped with 1.1 mm continuum cam era perform ed the intensive surveys of distant dusty forming galaxies during 2007 to 2008 and discovered a cluster of extrem e star-form ing galaxies (Fig.10) which is located 11.5 billion light-years away from the Earth and is associated with a large scale structure of galaxies (Tam ura et al. 2009).

5. M izu saw a VERA Observatory

VERA (VLBIExploration of Radio A strom etry) consists of four radio telescopes deployed over the Japanese archipelago to measure the parallax and proper motion of maser sources in our Galaxy using the Very Long Baseline Interferom etry (VLBI) technique. VERA uses a two-beam observing system, which makes it possible to observe two celestial objects simultaneously. This enables the phase uctuation caused by the atmospheric turbulence to be compensated for, and as



Fig. 9. Discovery of massive black hole at the center of NGC 4258, proven by the K eplerian motion of water maser sources.



Fig. 10. Submillimeter galaxies (red) detected by ASTE superposed on the distribution of infrared sources (green) and optical sources (blue).

a result, the positions of the radio sources can be determ ined very accurately. VERA is also able to determ ine the distance to objects with 10 m icroarc-second accuracy. This is a kind of triangulation m ethod based on the Earth's orbit.

VERA dem onstrated (cf. Fig.11) its high capability of precise astrom etry by breaking the world record of measuring the largest distance by means of trigonom etric parallax (Honma et al. 2007) as well as by providing the best distance measurement for Orion KL, one of the most important G alactic objects.



Fig. 11. Annual parallax of a waterm as source in the star forming region S269 was measured by VERA for the rst time to be 189 8 m icro arcsec.

6. O kayam a A strophysical O bservatory

The 1.88m telescope at 0 kayam a A strophysical 0 bservatory (OAO) (Yoshida 2005) was the largest optical infrared telescope and the heart of the modern astrophysics in Japan during 1960–1990s. The telescope has been used as a com - mon use facility and more than 200 researchers are visiting at OAO. High dispersion optical spectroscopy of stars is the main research activity (Sadakane et al. 2005). One of the recent achievem ents is discovery of ten extra-solar planets around G-giant stars (Sato et al. 2008). Fig. 12 shows an exam ple of the orbitalm odulation of radial velocity of such a star. There is a plan to construct a 3.8m telescope in the O kayam a cam - pus as a new facility of K yoto University.



Fig. 12. Observed radial velocities of HD 167042 (dots). The Keplerian orbital t is shown by the solid line.

7. TAM A 300 G ravitational W ave D etection Experim ent

The gravitational wave detection facility TAM A 300, with a Fabry-Perot interferom eter of 300m span to measure the time varying deformation of the space due to the gravitational wave, has been in contiguous operation since 1999. Sensitivity of the system has been much im proved by the development of the new seism ic attenuation system as is shown in Fig.13 (A rai et al. 2007).



Fig. 13. Tam a300 sensitivity with new seism ic attenuation system (SAS) compared with that of old SAS.

8. Kaguya-Lunar m ission

KAGUYA (SELENE) was launched on Septem – ber 14, 2007 by H-IIA to study the structure and the origin of the M oon. NAOJ is responsible for topography m easurem ent using a laser altimeter and gravity measurem ent using relay and VLBI subsatellites. NAOJ team reports lunar global shape and topography with two order of magnitude higher density and better vertical resolution than previously available. The result shows that the highest point on the M oon is on the southerm rim of the D irichlet-Jackson basin, and the lowest one is in the Antoniadi crater. The full-range topography spans about 19.81km, which is greater than previous estimation (A rakiet al. 2009). D etailed maps of polar regions were obtained for the

rst time (cfF ig.14). A dram atically improved lunar gravity model is also reported (N am iki et al. 2009). The model includes far side tracking data for the rst time. The results reveal that the far side impact basins do not resemble their nearside counterparts. The researchers interpret the results as indicating cooler, more rigid early conditions on the far side than previously thought. VLBI observations using VERA and international stations contribute signi cant improvement of orbital determ ination of satellites, which would improve lunar gravity model.



Fig. 14. Maps of Lunar northern polar region (left) and southern polar region (right) rst obtained by Kaquya mission.

9. Centers and D ivisions

A dvanced Technology Center(ATC) has machine shop, optical shop, space chamber shop (Fig. 15), design shop, high precision machining facilities and a coating facility. It was a center for developing Subaru Telescope instruments in late 90's. Currently, active projects run at ATC include band 4, 8, and 10 receiver developments for ALM A, R & D e orts for tera hertz detectors, gravitational wave detector and light weight mirror technology.



Fig. 15. Large clean room for testing the perform ances of telescope and instrum ents for space missions.

A stronom y Data Center (ADC) o ers computer systems for astronom ical data analysis for astronom ers and operates astronom ical data archives center. SM OKA provides access to the public science data obtained at Subaru Telescope, 188cm telescope at O kayam a A strophysical O bservatory, and 105cm Schm idt telescope at K iso O bærvatory (U niversity of Tokyo). Japanese V irtual O bærvatory project is also a part of ADC's activity.

Center for Com putational A strophysics (CfCA) has several supercom puters dedicated to astrophysical large-scale simulations. Currently, CfCA hosts NEC SX-9 vector supercom puter (2T ops) and Cray XT4 parallel supercom puter (28T ops), as well as GRAPE-6 and 7 specialpurpose com puters for gravitational N-body sim – ulations. These facilities are used by astrophysicists both within and outside Japan, for studies of supernova explosion, accretion disks, star form ation, planet form ation, galaxy form ation, cosm ology, and other areas of theoretical astrophysics. Fig. 16 shows the result of recent self-consistent simulation of a disk galaxy perform ed on the C ray system, in which gas cooling is followed down to 10K, resulting in the spiralarm swhich look quite similar to that of real spiral galaxies.

Public Relations Center (PRC) provide dedicated services for public outreach, publications, library, and ephemeris computation.



Fig. 16. A spiral galaxy as reproduced in a selfconsistent simulation incorporating physica processes.

There are four divisions for Optical and Infrared A stronom y, R adio A stronom y, Solar and P lasm a A strophysics, and T heoretical A stronom y, where m any of individual researchers are a liated.

10. O ther Space M issions

As for the H inode m ission, a new solar observatory in space, see (T suneta 2009). NAOJ astronom ers are collaborating with JAXA scientists for research and developm ent acitivities to realize 3.5m SPICA (SPace Infrared Telescope for C osm ology and A strophysics) m ission, a m idand far-infrared astronom y m ission with a cooled (4.5 K) telescope launched into Sun-Earth L2 orbit, V SOP -2, a space V LBI m ission following the V SOP (Haruka) to im age AGNs at 30 m icro arcsec resolution, and JASM INE (Japan A strom etry Satellite M ission for IN frared Exploration) to m easure position, distance and proper potions of stars in the bulge of the G alaxy.

Upon an urgent request from the editor, the

present paper was written by drafting the outline of the NAOJ and compiling the descriptions available in recent NAOJ web releases and those given in NAOJ 2008 annual report (NAOJ 2008) with some editing and revisions. The author acknowledges helpful quick comments delivered by M.fujimoto, R.K.awabe, J.M.akino, T.Sakurai, S.Sasaki, Y.Suematsu, K.T.atematsu, S.T.suneta, and M.Yoshida to improve the rst draft.

REFERENCES

- 2008, Annual Report of the NAOJ, 20
- Aoki,W .et al: 2006, ApJ, 638, 17.
- Arai, K. et al.: 2008, Class. Quantum Grav., 25, 114036.
- A raki,H. et al.: 2009, Science, 323, 897
- Fukagawa, M. et al, 2009, ApJL, 696, L1
- Honm a,M .et al: 2007, PASJ, 59, 889.
- Iye,M .et al.: 2006, N ature, 443, 186
- M arois, C. et al., 2008, Science, 322, 1348
- M iyoshi, M. et al.: 1995, Nature, 373, 127
- Nam iki, N. et al.: 2009, Science, 323, 900
- Sadakane, K., Hirata, R. and Tsuji, T.: 2005, PASJ, 57, 1.
- Sato, B. et al.: 2008, PASJ, 60, 1317.
- Tamura, et al. Nature, 495, 61.
- T suneta, S.: 2009, this volum e 11-18.
- YagiM .et al: 2007, ApJ, 660, 1209.
- Yoshida, M .: 2005, JKAS, 38, 117.

T h is 2-colum n preprint was prepared with the AAS ${\rm IAT}_EX$ m acros v5.2.