

Periodic and solitary wave solutions to the Fornberg-Whitham equation

Jiangbo Zhou*, Lixin Tian

*Nonlinear Scientific Research Center, Faculty of Science, Jiangsu University,
Zhenjiang, Jiangsu 212013, China*

Abstract

In this paper, new travelling wave solutions to the Fornberg-Whitham equation

$$u_t - u_{xxt} + u_x + uu_x = uu_{xx} + 3u_x u_{xx}$$

are investigated. They are characterized by two parameters. The expressions of the periodic and solitary wave solutions are obtained.

Key words: Fornberg-Whitham equation, solitary wave, periodic wave

1 Introduction

Recently, Ivanov [1] investigated the integrability of a class of nonlinear dispersive wave equations

$$u_t - u_{xxt} + \partial_x(\kappa u + \alpha u^2 + \beta u^3) = \nu u_x u_{xx} + \gamma u u_{xxx}, \quad (1.1)$$

where $\alpha, \beta, \gamma, \kappa, \nu$ are real constants.

The important cases of Eq.(1.1) are:

The hyperelastic-rod wave equation

$$u_t - u_{xxt} + 3uu_x = \gamma(2u_x u_{xx} + uu_{xxx}), \quad (1.2)$$

* Corresponding author.

Email address: zhoujiangbo@yahoo.cn (Jiangbo Zhou).

has been recently studied as a model, describing nonlinear dispersive waves in cylindrical compressible hyperelastic rods [2]-[7]. The physical parameters of various compressible materials put γ in the range from -29.4760 to 3.4174 [2,4].

The Camassa-Holm equation

$$u_t - u_{xxt} + 3uu_x = 2u_x u_{xx} + uu_{xxx}, \quad (1.3)$$

describes the unidirectional propagation of shallow water waves over a flat bottom [8,9]. It is completely integrable [1] and admits, in addition to smooth waves, a multitude of travelling wave solutions with singularities: peakons, cuspons, stumpons and composite waves [9]-[12]. The solitary waves of Eq.(1.2) are smooth if $\kappa > 0$ and peaked if $\kappa = 0$ [9,10]. Its solitary waves are stable solitons [13,14], retaining their shape and form after interactions [15]. It models wave breaking [16]-[18].

The Degasperis-Procesi equation

$$u_t - u_{xxt} + 4uu_x = 3u_x u_{xx} + uu_{xxx}, \quad (1.4)$$

models nonlinear shallow water dynamics. It is completely integrable [1] and has a variety of travelling wave solutions including solitary wave solutions, peakon solutions and shock waves solutions [20]-[27].

The Fornberg-Whitham equation

$$u_t - u_{xxt} + u_x + uu_x = uu_{xxx} + 3u_x u_{xx}, \quad (1.5)$$

appeared in the study qualitative behaviors of wave-breaking [28]. It admits a wave of greatest height, as a peaked limiting form of the travelling wave solution [29], $u(x, t) = A \exp(-\frac{1}{2} |x - \frac{4}{3}t|)$, where A is an arbitrary constant. It is not completely integrable [1].

The regularized long-wave or BBM equation

$$u_t - u_{xxt} + u_x + uu_x = 0, \quad (1.6)$$

and the modified BBM equation

$$u_t - u_{xxt} + u_x + 3u^2 u_x = 0, \quad (1.7)$$

have also been investigated by many authors [30]-[38].

Many efforts have been devoted to study Eq.(1.2)-(1.4),(1.6) and (1.7), however, little attention was paid to study Eq.(1.5). In [39], we constructed two types of bounded travelling wave solutions $u(\xi)$ ($\xi = x - ct$) to Eq.(1.5), which are defined on semifinal bounded domains and called kink-like and antikink-like wave solutions. In this paper, we continue to study the travelling wave solutions to Eq.(1.5). Following Vakhnenko and Parkes's strategy in [19], we obtain some periodic and solitary wave solutions $u(\xi)$ to Eq.(1.5) which are defined on $(-\infty, +\infty)$. The travelling wave solutions obtained in this paper are obviously different from those obtained in our previous work [39]. To the best of our knowledge, these solutions are new for Eq.(1.5). Our work may help people to know deeply the described physical process and possible applications of the Fornberg-Whitham equation.

The remainder of the paper is organized as follows. In Section 2, for completeness and readability, we repeat Appendix A in [19], which discuss the solutions to a first-order ordinary differential equation. In Section 3, we show that, for travelling wave solutions, Eq.(1.5) may be reduced to a first-order ordinary differential equation involving two arbitrary integration constants a and b . We show that there are four distinct periodic solutions corresponding to four different ranges of values of a and restricted ranges of values of b . A short conclusion is given in Section 4.

2 Solutions to a first-order ordinary differential equation

This section is due to Vakhnenko and Parkes (see Appendix A in [19]). For completeness and readability, we repeat it in the following.

Consider solutions to the following ordinary differential equation

$$(\varphi\varphi_\xi)^2 = \varepsilon^2 f(\varphi), \quad (2.1)$$

where

$$f(\varphi) = (\varphi - \varphi_1)(\varphi - \varphi_2)(\varphi_3 - \varphi)(\varphi_4 - \varphi), \quad (2.2)$$

and $\varphi_1, \varphi_2, \varphi_3, \varphi_4$ are chosen to be real constants with $\varphi_1 \leq \varphi_2 \leq \varphi \leq \varphi_3 \leq \varphi_4$.

Following [40] we introduce ζ defined by

$$\frac{d\xi}{d\zeta} = \frac{\varphi}{\varepsilon}, \quad (2.3)$$

so that Eq.(2.1) becomes

$$(\varphi_\zeta)^2 = f(\varphi). \quad (2.4)$$

Eq.(2.4) has two possible forms of solution. The first form is found using result 254.00 in [41]. Its parametric form is

$$\begin{cases} \varphi = \frac{\varphi_2 - \varphi_1 n \operatorname{sn}^2(w|m)}{1 - n \operatorname{sn}^2(w|m)}, \\ \xi = \frac{1}{\varepsilon p} (w \varphi_1 + (\varphi_2 - \varphi_1) \Pi(n; w|m)), \end{cases} \quad (2.5)$$

with w as the parameter, where

$$m = \frac{(\varphi_3 - \varphi_2)(\varphi_4 - \varphi_1)}{(\varphi_4 - \varphi_2)(\varphi_3 - \varphi_1)}, p = \frac{1}{2} \sqrt{(\varphi_4 - \varphi_2)(\varphi_3 - \varphi_1)}, w = p\zeta, \quad (2.6)$$

and

$$n = \frac{\varphi_3 - \varphi_2}{\varphi_3 - \varphi_1}. \quad (2.7)$$

In (2.5) $\operatorname{sn}(w|m)$ is a Jacobian elliptic function, where the notation is as used in Chapter 16 of [42]. $\Pi(n; w|m)$ is the elliptic integral of the third kind and the notation is as used in Section 17.2.15 of [42].

The solution to (2.1) is given in parametric form by (2.5) with w as the parameter. With respect to w , φ in (2.5) is periodic with period $2K(m)$, where $K(m)$ is the complete elliptic integral of the first kind. It follows from (2.5) that the wavelength λ of the solution to (2.1) is

$$\lambda = \frac{2}{\varepsilon p} |\varphi_1 K(m) + (\varphi_2 - \varphi_1) \Pi(n|m)|, \quad (2.8)$$

where $\Pi(n|m)$ is the complete elliptic integral of the third kind.

When $\varphi_3 = \varphi_4$, $m = 1$, (2.5) becomes

$$\begin{cases} \varphi = \frac{\varphi_2 - \varphi_1 n \tanh^2 w}{1 - n \tanh^2 w}, \\ \xi = \frac{1}{\varepsilon} \left(\frac{w \varphi_3}{p} - 2 \tanh^{-1}(\sqrt{n} \tanh w) \right). \end{cases} \quad (2.9)$$

The second form of solution of (2.5) is found using result 255.00 in [41]. Its

parametric form is

$$\begin{cases} \varphi = \frac{\varphi_3 - \varphi_4 n \operatorname{sn}^2(w|m)}{1 - n \operatorname{sn}^2(w|m)}, \\ \xi = \frac{1}{\varepsilon p} (w \varphi_4 - (\varphi_4 - \varphi_3) \Pi(n; w|m)), \end{cases} \quad (2.10)$$

where m, p, w are as in (2.6), and

$$n = \frac{\varphi_3 - \varphi_2}{\varphi_4 - \varphi_2}. \quad (2.11)$$

The solution to (2.1) is given in parametric form by (2.10) with w as the parameter. The wavelength λ of the solution to (2.1) is

$$\lambda = \frac{2}{\varepsilon p} |\varphi_4 K(m) - (\varphi_4 - \varphi_3) \Pi(n|m)|. \quad (2.12)$$

When $\varphi_1 = \varphi_2$, $m = 1$, (2.10) becomes

$$\begin{cases} \varphi = \frac{\varphi_3 - \varphi_4 n \tanh^2 w}{1 - n \tanh^2 w}, \\ \xi = \frac{1}{\varepsilon} \left(\frac{w \varphi_2}{p} + 2 \tanh^{-1}(\sqrt{n} \tanh w) \right). \end{cases} \quad (2.13)$$

3 Periodic and solitary wave solutions to Eq.(1.5)

Eq.(1.5) can also be written in the form

$$(u_t + uu_x)_{xx} = u_t + uu_x + u_x. \quad (3.1)$$

Let $u = \varphi(\xi) + c$ with $\xi = x - ct$ be a travelling wave solution to Eq.(3.1), then it follows that

$$(\varphi \varphi_\xi)_{\xi\xi} = \varphi \varphi_\xi + \varphi_\xi. \quad (3.2)$$

Integrating (3.2) twice with respect to ξ , we have

$$(\varphi \varphi_\xi)^2 = \frac{1}{4} (\varphi^4 + \frac{8}{3} \varphi^3 + a \varphi^2 + b), \quad (3.3)$$

where a and b are two arbitrary integration constants.

Eq.(3.3) is in the form of Eq.(2.1) with $\varepsilon = \frac{1}{2}$ and $f(\varphi) = (\varphi^4 + \frac{8}{3}\varphi^3 + a\varphi^2 + b)$. For convenience we define $g(\varphi)$ and $h(\varphi)$ by

$$f(\varphi) = \varphi^2 g(\varphi) + b, \text{ where } g(\varphi) = \varphi^2 + \frac{8}{3}\varphi + a, \quad (3.4)$$

$$f'(\varphi) = 2\varphi h(\varphi), \text{ where } h(\varphi) = 2\varphi^2 + 4\varphi + a, \quad (3.5)$$

and define φ_L , φ_R , b_L , and b_R by

$$\varphi_L = -\frac{1}{2}(2 + \sqrt{4 - 2a}), \varphi_R = -\frac{1}{2}(2 - \sqrt{4 - 2a}), \quad (3.6)$$

$$b_L = -\varphi_L^2 g(\varphi_L) = \frac{a^2}{4} - 2a + \frac{8}{3} + \frac{2}{3}(2 - a)\sqrt{4 - 2a}, \quad (3.7)$$

$$b_R = -\varphi_R^2 g(\varphi_R) = \frac{a^2}{4} - 2a + \frac{8}{3} - \frac{2}{3}(2 - a)\sqrt{4 - 2a}. \quad (3.8)$$

Obviously, φ_L , φ_R are the roots of $h(\varphi) = 0$.

In the following, suppose that $a < 2$ and $a \neq 0$ such that $f(\varphi)$ has three distinct stationary points: φ_L , φ_R , 0 and comprise two minimums separated by a maximum. Under this assumption, Eq.(3.3) has periodic and solitary wave solutions that have different analytical forms depending on the values of a and b as follows:

(1) $a < 0$

In this case $\varphi_L < 0 < \varphi_R$ and $f(\varphi_L) < f(\varphi_R)$. For each value $a < 0$ and $0 < b < b_R$ (a corresponding curve of $f(\varphi)$ is shown in Fig.1(a)), there are periodic inverted loop-like solutions to Eq.(3.3) given by (2.5) so that $0 < m < 1$, and with wavelength given by (2.8); see Fig.2(a) for an example.

The case $a < 0$ and $b = b_R$ (a corresponding curve of $f(\varphi)$ is shown in Fig.1(b)) corresponds to the limit $\varphi_3 = \varphi_4 = \varphi_R$ so that $m = 1$, and then the solution is an inverted loop-like solitary wave given by (2.9) with $\varphi_2 \leq \varphi < \varphi_R$ and

$$\varphi_1 = -\frac{1}{6}(2 + 3\sqrt{4 - 2a} + 2\sqrt{4 + 6\sqrt{4 - 2a}}), \quad (3.9)$$

$$\varphi_2 = -\frac{1}{6}(2 + 3\sqrt{4 - 2a} - 2\sqrt{4 + 6\sqrt{4 - 2a}}), \quad (3.10)$$

see Fig.3(a) for an example.




Fig. 1. The curve of $f(\varphi)$. (a) $a = -50$, $b = 233$; (b) $a = -50$, $b = 374.1346$; (c) $a = 1.5$, $b = -0.05$; (d) $a = 1.5$, $b = 0$; (e) $a = \frac{16}{9}$, $b = -0.1$; (f) $a = \frac{16}{9}$, $b = 0$; (g) $a = 1.9$, $b = -0.24$; (h) $a = 1.9$, $b = -0.2010$.

$$(2) \ 0 < a < \frac{16}{9}$$

In this case $\varphi_L < \varphi_R < 0$ and $f(\varphi_L) < f(0)$. For each value $0 < a < \frac{16}{9}$ and $b_R < b < 0$ (a corresponding curve of $f(\varphi)$ is shown in Fig.1(c)), there are periodic hump-like solutions to Eq.(3.3) given by (2.5) so that $0 < m < 1$, and with wavelength given by (2.8); see Fig.2(b) for an example.

The case $0 < a < \frac{16}{9}$ and $b = 0$ (a corresponding curve of $f(\varphi)$ is shown in Fig.1(d)) corresponds to the limit $\varphi_3 = \varphi_4 = 0$ so that $m = 1$, and then the solution can be given by (2.9) with φ_1 and φ_2 given by the roots of $g(\varphi) = 0$, namely

$$\varphi_1 = -\frac{4}{3} - \frac{1}{3}\sqrt{16 - 9a}, \varphi_2 = -\frac{4}{3} + \frac{1}{3}\sqrt{16 - 9a}. \quad (3.11)$$

In this case we obtain a weak solution, namely the periodic upward-cusp wave

$$\varphi = \varphi(\xi - 2j\xi_m), (2j - 1)\xi_m < \xi < (2j + 1)\xi_m, \ j = 0, \pm 1, \pm 2, \dots, \quad (3.12)$$

where

$$\varphi(\xi) = (\varphi_2 - \varphi_1 \tanh^2(\xi/4)) \cosh^2(\xi/4), \quad (3.13)$$

and

$$\xi_m = 4 \tanh^{-1} \sqrt{\frac{\varphi_2}{\varphi_1}}, \quad (3.14)$$

see Fig.3(b) for an example.




Fig. 2. Periodic solutions to Eq.(3.3) with $0 < m < 1$. (a) $a = -50$, $b = 233$ so $m = 0.7885$; (b) $a = 1.5$, $b = -0.05$ so $m = 0.6893$; (c) $a = \frac{16}{9}$, $b = -0.1$ so $m = 0.8254$; (d) $a = 1.9$, $b = -0.24$ so $m = 0.6121$.




Fig. 3. Solutions to Eq.(3.3) with $m = 1$. (a) $a = -50$, $b = 374.1346$; (b) $a = 1.5$, $b = 0$; (c) $a = \frac{16}{9}$, $b = 0$; (d) $a = 1.9$, $b = -0.2010$.

$$(3) a = \frac{16}{9}$$

In this case $\varphi_L < \varphi_R < 0$ and $f(\varphi_L) = f(0)$. For $a = \frac{16}{9}$ and each value $b_R < b < 0$ (a corresponding curve of $f(\varphi)$ is shown in Fig.1(e)), there are periodic hump-like solutions to Eq.(3.3) given by (2.10) so that $0 < m < 1$, and with wavelength given by (2.12); see Fig.2(c) for an example.

The case $a = \frac{16}{9}$ and $b = 0$ (a corresponding curve of $f(\varphi)$ is shown in Fig.1(f)) corresponds to the limit $\varphi_1 = \varphi_2 = \varphi_L = -\frac{4}{3}$ and $\varphi_3 = \varphi_4 = 0$ so that $m = 1$. In this case neither (2.5) nor (2.10) is appropriate. Instead we consider Eq.(3.3) with $f(\varphi) = \frac{1}{4}\varphi^2(\varphi + \frac{4}{3})^2$ and note that the bound solution has $-\frac{4}{3} < \varphi \leq 0$. On integrating Eq.(3.3) and setting $\varphi = 0$ at $\xi = 0$ we obtain a weak solution

$$\varphi = \frac{4}{3} \exp\left(-\frac{1}{2}|\xi|\right) - \frac{4}{3}, \quad (3.15)$$

i.e. a single peakon solution with amplitude $\frac{4}{3}$, see Fig.3(c).

$$(4) \quad \frac{16}{9} < a < 2$$

In this case $\varphi_L < \varphi_R < 0$ and $f(\varphi_L) > f(0)$. For each value $\frac{16}{9} < a < 2$ and $b_R < b < b_L$ (a corresponding curve of $f(\varphi)$ is shown in Fig.1(g)), there are periodic hump-like solutions to Eq.(3.3) given by (2.10) so that $0 < m < 1$, and with wavelength given by (2.12); see Fig.2(d) for an example.

The case $\frac{16}{9} < a < 2$ and $b = b_L$ (a corresponding curve of $f(\varphi)$ is shown in Fig.1(h)) corresponds to the limit $\varphi_1 = \varphi_2 = \varphi_L$ so that $m = 1$, and then the solution is a hump-like solitary wave given by (2.13) with $\varphi_L < \varphi \leq \varphi_3$ and

$$\varphi_3 = \frac{1}{6}(-2 + 3\sqrt{4 - 2a} - 2\sqrt{4 - 6\sqrt{4 - 2a}}), \quad (3.16)$$

$$\varphi_4 = \frac{1}{6}(-2 + 3\sqrt{4 - 2a} + 2\sqrt{4 - 6\sqrt{4 - 2a}}), \quad (3.17)$$

see Fig.3(d) for an example.

On the above, we have obtained expressions of parametric form for periodic and solitary wave solutions $\varphi(\xi)$ to Eq.(3.3). So in terms of $u = \varphi(\xi) + c$, we can get expressions for the periodic and solitary wave solutions $u(\xi)$ to Eq.(1.5).

4 Conclusion

In this paper, we have found expressions for new travelling wave solutions to the Fornberg-Whitham equation. These solutions depend, in effect, on two parameters a and m . For $m = 1$, there are inverted loop-like ($a < 0$), single peaked ($a = \frac{16}{9}$) and hump-like ($\frac{16}{9} < a < 2$) solitary wave solutions. For

$m = 1, 0 < a < \frac{16}{9}$ or $0 < m < 1, a < 2$ and $a \neq 0$, there are periodic hump-like wave solutions.

References

- [1] R. Ivanov, On the integrability of a class of nonlinear dispersive wave equations, J. Nonlinear Math. Phys. 1294 (2005) 462-468.
- [2] H. H. Dai, Model equations for nonlinear dispersive waves in a compressible Mooney-Rivlin rod, Acta Mech. 127 (1998) 193-207.
- [3] A. Constantin, W. A. Strauss, Stability of a class of solitary waves in compressible elastic rods, Phys. Lett. A 270 (2000) 140-148.
- [4] H. H. Dai, Y. Huo, Solitary shock waves and other travelling waves in a general compressible hyperelastic rod, Proc. R. Soc. A 456 (2000) 331-363.
- [5] Z. Y. Yin, On the cauchy problem for a nonlinearly dispersive wave equation, J. Nonlinear Math. Phys. 10 (2003) 10-15.
- [6] Y. Zhou, Stability of solitary waves for a rod equation, Chaos Solitons and Fractals 21 (2004) 977-981.
- [7] Y. Zhou, Blow-up of solutions to a nonlinear dispersive rod equation, Calc. Var. 25 (2005) 63-77.
- [8] R. S. Johnson, Camassa-Holm, Korteweg-de Vries and related models for water waves, J. Fluid. Mech. 457 (2002) 63-82.
- [9] R. Camassa, D. Holm, An integrable shallow water equation with peaked solitons, Phys. Rev. Lett. 71 (1993) 1661-1664.
- [10] Y. A. Li, P. J. Olver, P. Rosenau, Non-analytic solutions of nonlinear wave models, in nonlinear theory of generalised functions: Vienna 1997, Chapman & Hall/CRC Res. Notes Math. 401, Chapman & Hall/CRC, Boca Raton, FL, 1999, 129-145.
- [11] J. Lenells, Traveling wave solutions of the Camassa-Holm equation, J. Differ. Equ. 217 (2005) 393-430.
- [12] Y. A. Li, P. J. Olver, Well-posedness and blow-up solutions for an integrable nonlinearly dispersive model wave equation, J. Differ. Equ. 162 (2000) 27-63.
- [13] A. Constantin, W. A. Strauss, Stability of the Camassa-Holm solitons, J. Nonlinear Sci. 12 (2002) 415-422.
- [14] A. Constantin, W. A. Strauss, Stability of peakons, Commun. Pure Appl. Math. 53 (2000) 603-610.
- [15] R. S. Johnson, On the solutions of the Camassa-Holm equation, Proc. R. Soc. London A 459 (2003) 1687-1708.

- [16] A . Constantin, Existence of permanent and breaking waves for a shallow water equation: a geometric approach, *Ann. Inst. Fourier (Grenoble)* 50 (2000) 321-362.
- [17] A. Constantin, J. Escher, Wave breaking for nonlinear nonlocal shallow water equations, *Acta Math.* 181 (1998) 229-243.
- [18] A. Constantin, J. Escher, On the blow-up rate and the blow-up set of breaking waves for a shallow water equation, *Math. Z.* 233 (2000) 75-91.
- [19] V. O. Vakhnenko, E. J. Parkes, Periodic and solitary-wave solutions of the Degasperis-Procesi equation, *Chaos, Solitons and Fractals* 20 (2004) 1059-1073.
- [20] H. Lundmark, Formation and dynamics of shock waves in the Degasperis-Procesi equation, *J. Nonlinear Sci.* 17 (2007) 169-198.
- [21] J. Escher, Y. Liu, Z. Yin, Shock waves and blow-up phenomena for the periodic Degasperis-Procesi equation, *Ind. Univ. Math. J.* 56 (2007) 187-117.
- [22] G. M. Coclite, K. H. Karlsen, N. H. Risebro, Numerical schemes for computing discontinuous solutions of the Degasperis-Procesi equation, *IMA J. Numer. Anal.* 28 (2008) 80-105.
- [23] G. M. Coclite, K. H. Karlsen, N. H. Risebro, On the uniqueness of discontinuous solutions to the Degasperis-Procesi equation, *J. Diff. Equ.* 234 (2007) 142-160.
- [24] J. Lenells, Traveling wave solutions of the Degasperis-Procesi equation, *J. Math. Anal. Appl.* 306 (2005) 72-82.
- [25] Y. Matsuno, Multisoliton solutions of the Degasperis-Procesi equation and their peakon limit, *Inverse Problems* 21 (2005) 1553-1570.
- [26] H. Lundmark, J. Szmigielski, Multi-peakon solutions of the Degasperis-Procesi equation, *Inverse Problems* 19 (2003) 1241-1245.
- [27] Y. Matsuno, Cusp and loop soliton solutions of short-wave models, for the Camassa-Holm and Degasperis-Procesi equations, *Phys. Lett. A* 359 (2006) 451-457.
- [28] G. B. Whitham, Variational methods and applications to water wave, *Proc. R. Soc. Lond. Ser. A* 299 (1967) 6-25.
- [29] B. Fornberg, G. B. Whitham, A numerical and theoretical study of certain nonlinear wave phenomena, *Philos. Trans. R. Soc. Lond. Ser. A* 289 (1978) 373-404.
- [30] A. C. Bryan, A. E. G. Stuart, Solitons and the regularized long wave equation: A nonexistence theorem , *Chaos, Solitons and Fractals* 7 (1996) 1881-1886.
- [31] D. Bhardwaj, R. Shankar, A computational method for regularized long wave equation, *Comput. Math. Appl.* 40 (2000) 1397-1404.
- [32] S. I. Zaki, Solitary waves of the splitted RLW equation, *Comput. Phys. Commun.* 138 (2001) 80-91.

- [33] J. I. Ramos, Solitary waves of the EW and RLW equations, *Chaos, Solitons and Fractals* 34 (2007) 1498-1518.
- [34] J. I. Ramos, Solitary wave interactions of the GRLW equation, *Chaos, Solitons and Fractals* 33 (2007) 479-491.
- [35] J. Nickel, Elliptic solutions to a generalized BBM equation, *Phys. Lett. A* 364 (2007) 221-226.
- [36] A. H. A. Ali, A. A. Soliman, K. R. Raslan, Soliton solution for nonlinear partial differential equations by cosine-function method, *Phys. Lett. A* 368 (2007) 299-304.
- [37] D. H. Feng, J. B. Li, J. L. Lu, T. L. He, New explicit and exact solutions for a system of variant RLW equations, *Appl. Math. Comput.* 198 (2008) 715-720.
- [38] Z. W. Lin, Instability of nonlinear dispersive solitary waves, *J. Funct. Anal.* 255 (2008) 1191-1224.
- [39] J. B. Zhou, L. X. Tian, A type of bounded traveling wave solutions for the Fornberg-Whitham equation, *J. Math. Anal. Appl.* 346 (2008) 255-261.
- [40] E. J. Parkes, The stability of solutions of Vakhnenko's equation, *J. Phys. A Math. Gen.* 26 (1993) 6469-75.
- [41] P. F. Byrd, M. D. Friedman, *Handbook of elliptic integrals for engineers and scientists*. Berlin: Springer; 1971.
- [42] M. Abramowitz, I. A. Stegun. *Handbook of mathematical functions*. New York: Dover; 1972.