

# Multiscale expansions of difference equations in the small lattice spacing regime, and a vicinity and integrability test. I

Paolo Maria Santini<sup>1,\*</sup>

<sup>1</sup> Dipartimento di Fisica, Università di Roma "La Sapienza", and  
Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1  
Piazzale Aldo Moro 2, I-00185 Roma, Italy

\*email: paolo.santini@roma1.infn.it

February 22, 2024

## Abstract

We propose an algorithmic procedure i) to study the "distance" between an integrable PDE and any discretization of it, in the small lattice spacing regime, and, at the same time, ii) to test the (asymptotic) integrability properties of such discretization. This method should provide, in particular, useful and concrete informations on how good is any numerical scheme used to integrate a given integrable PDE. The procedure, illustrated on a fairly general 10-parameter family of discretizations of the nonlinear Schrödinger equation, consists of the following three steps: i) the construction of the continuous multiscale expansion of a generic solution of the discrete system at all orders in  $\hbar$ , following [1]; ii) the application, to such expansion, of the Degasperis–Procesi (DP) integrability test [2, 3], to test the asymptotic integrability properties of the discrete system and its "distance" from its continuous limit; iii) the use of the main output of the DP test to construct infinitely many approximate symmetries and constants of motion of the discrete system, through novel and simple formulas.

## 1 Introduction

Given a partial differential equation (PDE) and a partial difference equation (P E) discretizing it, it is interesting to know, when the lattice spacing is small, "how close" the two models are. In particular, if the PDE is

integrable, it is important to have a way to establish if such a discretization preserves integrability or, at least, how "close" is to an integrable system, detecting the order, in  $\epsilon$ , at which the discretization departs from integrability and, correspondingly, the time scale at which one should expect numerical evidence of nonintegrability and/or chaos. In addition, given a PDE and two PEs discretizing it, it is also interesting to know, when the lattice spacing is small, "how close" the two PEs are.

In this paper we propose to answer these basic questions in the following way. Concentrating on an integrable PDE and on a PE discretizing it,

1) we construct and study in detail the multiscale expansion at all orders of a generic solution of the PE under scrutiny, generated in the small

regime, following the procedure developed in [1]. At  $O(1)$ , the leading term  $u$  of such asymptotic expansion satisfies the integrable PDE; to keep the expansion asymptotic, we eliminate the secularities due to the linear part of the PE, arising at each order, introducing infinitely many slow (time) variables and establishing that the evolution of  $u$  with respect to such slow times is described by the infinite hierarchy of commuting flows of the integrable PDE, as in [1].

2) We make use of the asymptotic integrability test developed by Degasperis-Procesi (DP) in [2, 3] on such a multiscale expansion to test, at all orders, the "asymptotic" integrability properties of the PE; in particular, detecting the order in  $\epsilon$  (and, correspondingly, the time scale) at which the discretization departs from integrability. At this time scale, f.i., numerical simulations are expected to give some evidence of non-integrable and/or chaotic behaviour.

3) We finally show how to make use of the main output of the DP test to construct infinitely many "approximate" symmetries, at a required order in  $\epsilon$ , of the PE under scrutiny, using novel and simple formulas.

Recent studies on the performances, as numerical schemes for their continuous limits, of PEs possessing the same (continuous) Lie point symmetries as their continuous limits can be found in [4, 5]. Studies on the performances, as numerical schemes for their continuous limits, of integrable discretizations of integrable PDEs can be found, f. i., in [6] and [7]; in this case, the integrable discretization possesses infinitely many exact generalized symmetries and constants of motion in involution at any order in  $\epsilon$ , reducing to the generalized symmetries and constants of motion of the integrable PDE in the continuous limit. The PEs selected by our approach possess instead infinitely many approximate generalized symmetries and constants of motion in involution at the required order in  $\epsilon$  (see x3.1), reducing to the generalized

symmetries and constants of motion of the integrable PDE in the continuous limit.

The procedure we propose should allow one to have a control on the "distance" between the PDE and its continuous limit, as well as on the distance between two different discretizations of the same PDE. Indeed, suppose we construct an asymptotic expansion of the form  $u_\epsilon = u + O(\epsilon^\alpha)$ ;  $\alpha > 0$ , where

$u$  is a generic solution of the PDE and  $u$  is the corresponding solution of its continuous limit; if, at  $O(\epsilon^\alpha)$ ,  $\alpha > 0$ , the PDE passes the DP test, we infer that  $\|u_\epsilon - u\| = O(\epsilon^\alpha)$  at time scales of  $O(\epsilon^{-\beta})$ , where  $\| \cdot \|$  is the uniform norm wrt  $x$  and  $t$  (the norm used to test the asymptotic character of the generated multiscale expansion). In this way, since we control the distance between "generic solutions" of the PDE and of its continuous limit, we also control the distance between the PDE and its continuous limit. In addition, if the multiscale expansions of two different discretizations of the same PDE pass the DP test at  $O(\epsilon^\alpha)$ , we infer, from the triangular inequality, that  $\|u_\epsilon - v_\epsilon\| \leq \|u_\epsilon - u\| + \|u - v_\epsilon\| = O(\epsilon^\alpha)$  at time scales of  $O(\epsilon^{-\beta})$ , where  $u, v$  are solutions of the two different discretizations of the PDE corresponding to the same generic initial-boundary data; therefore we have a control also on the distance between the two different discretizations of the same PDE.

Some historical remarks are important, at this point, on the theory of multiscale expansions in connection with integrable systems, to put the results of this paper into a proper perspective. Multiscale expansions of a given PDE are very useful tools for investigating the properties of such a PDE and for identifying important model (universal) equations of physical phenomena. For instance, if the original nonlinear PDE has a dispersive linear part, a small amplitude monochromatic wave evolving according to it develops a slow space-time amplitude modulation described by the celebrated nonlinear Schrödinger (NLS) equation [8, 9, 10, 11, 12] (see also [13, 14, 15])

$$iu_t + u_{xx} + 2c|u|^2u = 0; \quad u = u(x;t) \in C; \quad (1)$$

integrable if  $c$  is a real constant [16]. Considering, instead, three monochromatic waves and imposing a suitable resonance condition on their wave numbers and dispersion relations, one generates another integrable universal model, the 3-wave resonant system [17]. In the above two examples, the expansion is constructed around "approximate" particular solutions of the original PDE (the monochromatic waves). It is also possible to expand around "exact" particular solutions of the original PDE; for instance, as

shown in [18], expanding around the exact solution  $u_0 = \exp(2ict)$  of (1), the first nontrivial term of the asymptotic expansion evolves according to another important model equation: the Korteweg-de Vries (KdV) equation [19], sharing with NLS the property of integrability [20]. Since multiscale expansions preserve integrability [18], i) if the original PDE is a "C-integrable" system (i.e., it is linearized by a "change of variables" [15, 21], like the Burgers equations [22]), the model equation generated by it is linear [15, 21]; ii) if the original PDE is an "S-integrable" system, or soliton equation (like the NLS equation), integrated in a more complicated way via a Riemann-Hilbert or  $\bar{\partial}$ -problem [23, 24, 25, 26], the model equation generated by it is also "S-integrable"; viceversa, iii) if the model equation generated by the expansion is not integrable, then the original equation is not integrable too. This criterion has been used in [15, 27, 28, 29] as a simple test of integrability. In addition, the universal character of the identified model equations (NLS, KdV or others) is also the reason why model equations possess very distinguished mathematical properties and, often, they are integrable [15, 27, 21].

Multiscale expansions can also be carried, in principle, to all orders and, as a consequence of eliminating the secular terms at each order, a sequence of slow time variables  $t_n = \epsilon^n t$  must be introduced and the dependence of the leading term of the expansion on such slow times is described by the hierarchy of commuting flows of the integrable model equation [1]. This multiscale expansion at all orders has been used in [2, 3] to build an efficient asymptotic integrability test for the original PDE (see x3 for more details on such test). An alternative asymptotic integrability test, based on the existence of approximate symmetries for the original PDE, can be found in [30]. The ideas and procedures developed in [1, 2, 3] have been recently used to build an integrability test also for PEs [31, 32, 33]; in this approach, one expands, as for the PDE case, around approximate or exact particular solutions of the PE under investigation, obtaining a continuous multiscale expansion at all orders, following [1], and applying on it the DP test. The main difference between the procedure followed in [31, 32, 33] and the results of this paper is the following. The standard multiscale approach used in [31, 32, 33], obtained expanding around approximate or exact particular solutions of the PE under investigation, cannot give informations on how close this PE and its continuous limit are, the main goal of the present paper. The common features of the procedure in [31, 32, 33] and of that used in this paper are that, in both cases, one constructs, from the given PE, continuous multiscale expansions carried to all orders, as in [1], and one applies to them

the DP integrability test. Therefore both procedures can be used to test the integrability and the asymptotic integrability of the original PDE.

Another integrability test for PDEs is the so-called "symmetry approach" [34], based on the existence of higher order symmetries and originally developed to test the integrability of PDEs [35, 36].

The results of this paper are illustrated on the basic prototype example of the NLS equation (1), starting from the following discretization of it:

$$i u_{n,t} + \frac{1}{2} (u_{n+1} + u_{n-1} - 2u_n) + F(u_{n-1}; u_n; u_{n+1}) = 0;$$

$$F(u_{n-1}; u_n; u_{n+1}) = 2a_1 u_n^2 + a_2 u_n^2 (u_{n+1} + u_{n-1}) + a_3 u_n^2 (u_{n+1} + u_{n-1}) + a_4 u_n (u_{n+1}^2 + u_{n-1}^2) + a_5 u_n (u_{n+1} u_{n-1} + u_{n+1} u_{n-1}) + a_6 u_n (u_{n+1}^2 + u_{n-1}^2) + 2a_7 u_n u_{n+1} u_{n-1} + a_8 (u_{n+1}^2 u_{n+1} + u_{n-1}^2 u_{n-1}) + a_9 (u_{n+1}^2 u_{n-1} + u_{n-1}^2 u_{n+1}) + a_{10} (u_{n+1}^2 u_{n-1} + u_{n-1}^2 u_{n+1});$$

where the constant coefficients  $a_j$ ;  $j = 1; \dots; 10$  are real, reducing to (1) in the natural continuous limit in which the lattice spacing  $\Delta x \rightarrow 0$  and  $n \rightarrow x \in \mathbb{R}$ ,  $u_n(t) \rightarrow u(x;t)$ , with

$$c = \sum_{j=1}^{10} a_j; \quad (3)$$

The 10-parameter family of equations (2) has been recently taken in [37] as the starting point of an analysis devoted to the identification of discretizations of NLS that possess, at the same time, a solitary wave and a breather solution reducing, respectively, to the one soliton and breather solutions of the NLS equation (1), in the continuous limit  $\Delta x \rightarrow 0$ . We remark that, rescaling the dependent variable, one can always introduce one normalization for the 10 coefficients; for instance, one can choose one of these coefficients, say  $a_j$ , to be sign( $a_j$ ) or, better for our purposes, one can normalize the sum (3) of the 10 coefficients to coincide with the prescribed coefficient  $c$  of the NLS equation (1).

The linear part of the discrete NLS (dNLS) (2) is the standard discretization of  $(i u_t + u_{xx})$ ; its nonlinear part is uniquely fixed by the following, physically sound, properties [37]. a) Equation (2) must possess the gauge symmetry of first kind (i.e., if  $u_n$  is a solution,  $u_n e^{i\theta}$  is a solution too, where  $\theta$  is an arbitrary real parameter), corresponding to the infinitesimal gauge

symmetry  $i_n$ . b) The nonlinearity is cubic; i.e., is the weakest nonlinearity compatible with the above gauge symmetry. c) Only first neighbours interactions are considered. d) Equation (2) is invariant under the symmetry transformation  $n \rightarrow n+1$  (space isotropy).

The dNLS (2) contains, in particular:

1) the integrable Ablowitz-Ladik (AL) equation [38]

$$i \psi_n \tau + \psi_n^2 (\psi_{n+1} + \psi_{n-1} - 2\psi_n) + a_2 j_n j_n^* (\psi_{n+1} + \psi_{n-1}) = 0; \quad (4)$$

for  $a_j = a_2 j^2$ ;  $j = 1; \dots; 10$ ;

2) the discretization

$$i \psi_n \tau + \psi_n^2 (\psi_{n+1} + \psi_{n-1} - 2\psi_n) + 2a_1 j_n j_n^* \psi_n = 0; \quad (5)$$

for  $a_j = a_1 j$ ;  $j = 1; \dots; 10$ , relevant in several applications [39, 40, 41, 42, 43], whose nonintegrability has been recently shown in [31, 32] using the DP test;

3) the discretization corresponding to

$$a_{10} = a_8; \quad a_1 = a_4 = a_5 = a_6 = a_7 = a_9 = 0; \quad (6)$$

with  $a_2; a_3; a_8$  arbitrary, possessing a solitary wave as well as a breather solution reducing, respectively, to the one soliton and breather solutions of the NLS equation in the limit  $\tau \rightarrow 0$  [37];

4) the discretization corresponding to

$$a_8 = a_3; \quad a_2 = 2a_3; \quad a_4 = 2a_6; \quad a_5 = a_7 = a_9 = a_{10} = 0; \quad (7)$$

where  $a_1; a_3; a_6$  are given in terms of physical quantities, describing coupled optical waveguides embedded in a material with Kerr nonlinearities [44];

5) the discretization corresponding to

$$a_4 = a_2; \quad a_1 = a_3 = a_6 = a_8 = a_{10} = 2; \quad a_5 = a_7 = a_9 = a_{10} = 0 \quad (8)$$

(a particular case of (7)), appearing in the modelling of the Fermi-Pasta-Ulam problem [45].

For special values of the coefficients  $a_j$ 's the dNLS equation (2) is Hamiltonian. For instance, equations (4), (5), (7) and (8) are Hamiltonian [44].

If  $0 < \tau < 1$ , the discrete scheme (2) approximates the NLS equation (1), (3) with an error of  $O(\tau^2)$ . To study more precisely how close equations

(2) and (1) are and, in particular, the integrability properties of (2), in this paper we follow the procedure indicated in the first part of this introduction, obtaining the following results.

1) Due to the structure of the vector field in (2), the generated  $\epsilon$ -expansion contains only even powers. At  $O(\epsilon^2)$ , the dNLS (2) passes the DP test if the 10 coefficients satisfy the elegant quadratic constraint

$$(a_1 - 3a_3 - 2a_4 - 6a_5 - 5a_6 + 3a_7 - 5a_8 - 13a_9 - a_{10}) \left( \sum_{j=1}^{10} a_j \right) = 0; \quad (9)$$

factorized into two linear constraints. If the first constraint  $\sum_{j=1}^{10} a_j = 0$  is satisfied, we are in the C-integrability framework and the dNLS (2) approximates the linear Schrodinger equation with an error of  $O(\epsilon^2)$ , for time scales of  $O(\epsilon^{-2})$ . If, instead, the second constraint is satisfied:

$$a_1 - 3a_3 - 2a_4 - 6a_5 - 5a_6 + 3a_7 - 5a_8 - 13a_9 - a_{10} = 0; \quad (10)$$

we are in the S-integrability framework and the dNLS (2) approximates the NLS equation (1),(3) with an error of  $O(\epsilon^2)$ , for time scales of  $O(\epsilon^{-2})$ .

We remark that, among the ten single dNLS equations obtained choosing only one of the ten coefficients different from zero in (2), only the AL equation (4) satisfies the constraint (9) and passes the test at  $O(\epsilon^2)$ .

2) At  $O(\epsilon^4)$  we have the following two scenarios. In the C-integrability framework, the dNLS (2) approximates, with an error of  $O(\epsilon^2)$ , the linear Schrodinger equation for time scales of  $O(\epsilon^{-4})$  if the following four linear constraints

$$\begin{aligned} \sum_{j=1}^{10} a_j &= 0; & a_1 + a_2 + a_6 + a_7 &= 0; & a_4 - a_5 + 2a_8 - 2a_9 &= 0; \\ a_2 + 2(a_3 + 3a_5 + 3a_6 - a_7 + a_8 + 7a_9) &= 0 \end{aligned} \quad (11)$$

are satisfied by the coefficients. Since one of the real  $a_j$ 's can always be fixed rescaling the dependent variable  $\psi$ , equations (11) characterize a 5-parameter family of discrete NLS equations (2) passing the test at such a high order.

In the S-integrability framework, the dNLS (2) approximates, with an error of  $O(\epsilon^2)$ , the NLS equation (1),(3) for time scales of  $O(\epsilon^{-4})$ , if the coefficients satisfy, together with the linear constraint (10), the five quadratic

constraints (53), (54)–(58). Since these five constraints do not contain the term  $(a_2)^2$ , they are trivially satisfied by the integrable AL equation (4), as it has to be. In general we do not expect a parametrization of such constraints in terms of elementary functions; however we have been able to construct the following two explicit examples of dNLS equations

$$\begin{aligned} i_{n,t} + \frac{1}{2} (j_{n+1} + j_{n-1} - 2j_n) + a_6 \left( 8j_n j_{n-1} + \frac{4}{3} j_n j_{n+1} (j_{n+1} + j_{n-1}) + \right. \\ \left. 4j_n^2 (j_{n+1} + j_{n-1}) - 4j_n (j_{n+1} j_{n-1} + j_{n+1} j_{n-1}) + j_n (j_{n+1}^2 + j_{n-1}^2) + \right. \\ \left. 2j_n j_{n+1} j_{n-1} \right) = 0; \end{aligned} \quad (12)$$

$$\begin{aligned} i_{n,t} + \frac{1}{2} (j_{n+1} + j_{n-1} - 2j_n) + a_9 \left( 48j_n j_{n-1} - 8j_n (j_{n+1} j_{n-1} + \right. \\ \left. j_{n-1} j_{n+1}) - 8j_n (j_{n+1} j_{n-1} + j_{n+1} j_{n-1}) + 10j_n (j_{n+1}^2 + j_{n-1}^2) + \right. \\ \left. 4j_n j_{n+1} j_{n-1} - 7(j_{n+1} j_{n+1} + j_{n-1} j_{n-1}) + \right. \\ \left. (j_{n+1}^2 j_{n-1} + j_{n-1}^2 j_{n+1}) + 6(j_{n+1} j_{n-1} + j_{n-1} j_{n+1}) \right) = 0; \end{aligned} \quad (13)$$

satisfying such complicated quadratic constraints, corresponding to particular cases in which the associated five quadrics degenerate into hyperplanes.

These two distinguished models, passing the test at such a high order through the above degeneration mechanism, are obviously good candidates to be S-integrable discretizations of NLS. A detailed study of their performances as numerical schemes for NLS, and of their possible integrability structure (Lax pair, etc..) is postponed to a subsequent paper.

To obtain the above results, it is essential to use well-known integrability properties of equation (1) (shared by all integrable systems; see, f.i. [46]–[49]) that we summarize here, for completeness.

The NLS equation belongs to a hierarchy of infinitely many commuting flows:

$$u_{t_n} = K_n(u); \quad n \geq 1 \quad (14)$$

ie., such that

$$[K_n(u), K_m(u)] = K_n^0(u) [K_m(u)] - K_m^0(u) [K_n(u)] = 0; \quad n, m \geq 1; \quad (15)$$

where

$$K_n^0(u) [f] = \lim_{\epsilon \rightarrow 0} \frac{K_n(u + \epsilon f) - K_n(u)}{\epsilon} \quad (16)$$

is the usual Frechet derivative of  $K_n(u)$  wrt  $u$  in the direction  $f$ . The commuting vector fields  $fK_n$ ,  $n \geq 1$  are arbitrary linear combinations, with constant coefficients, of the following basic symmetries  $f_n$ ,  $n \geq 1$ , generated by

the recursion relation

$$\begin{aligned} \hat{R}_{n+1} &= \hat{R}_n; \quad \hat{R}_0 = i\partial_x; \quad n \geq 1; \\ \hat{R}f &:= i f_x + 2cu_x^{-1}(uf + u_x f); \end{aligned} \quad (17)$$

where  $\hat{R}$  is the recursion operator of the NLS hierarchy [50]. The basic symmetries used in this paper are:

$$\begin{aligned} \hat{R}_0 &= i\partial_x; \quad \hat{R}_2 = i(u_{xx} + 2cu_x^2 u); \\ \hat{R}_4 &= i(u_{xxxx} + 2c(u^2 u_{xx} + 2u_x u_{xx}^2 + 4u_x^2 u_{xx} + 3u_x^2 u) + 6c^2 u_x^4 u); \\ \hat{R}_6 &= i(u_{xxxxx} + 2c(u^2 u_{xxx} + 6u_x u_{xxx}^2 + 4u u_x u_{xxx} + 9u u_x u_{xxx} + \\ &15u u_x u_{xxx} + 11u_x u_{xx}^2 + 10u_x^2 u_{xx}) + 10c^2(2u^2 u_x^2 u_{xx} + 2u u_x^2 u_{xx} + \\ &5u_x^2 u_{xx} + 5u_x^4 u_{xx} + u^3 u_x^2 + 6u_x u_x^2 u_x^2 + 7u_x u_x^2 u_x^2) + 20c^3 u_x u_x^2); \end{aligned} \quad (18)$$

and the NLS equation (1) corresponds to the flow  $u_{t_2} = K_2(u) = \hat{R}_2(u)$ .

Equivalently, the basic symmetries  $f_n, n \geq 1$  are elements of the kernel of the "linearized"  $n$ th flow operator  $\hat{M}_n; n \geq 1$ , defined by

$$\hat{M}_n f := f_{t_n} - K_n^0(u)[f]; \quad (19)$$

ie.:

$$\hat{M}_{n-m} = 0; \quad n, m \geq 1; \quad (20)$$

Due to (15), these linearized operators commute:

$$\hat{M}_n \hat{M}_m = \hat{M}_m \hat{M}_n; \quad n, m \geq 1; \quad (21)$$

The linearized operators used in this paper are:

$$\begin{aligned}
\hat{M}_2 f &:= f_{t_2} - i \frac{f_{xx}}{h} + 2c(u^2 f + 2j_x j f) ; \\
\hat{M}_4 f &:= f_{t_4} - \frac{i}{12} f_{xxxx} + 2c(u(6u_x f_x + 4u f_{xx}) + u(2u_x f_x + 2u_x f_x + \\
&uf_{xx}) + (6c j_x j u^2 + 3u_x^2 + 4uu_{xx})f + (9c j_x j^4 + 2j_x j^2 + 4uu_{xx} + \\
&2uu_{xx})f) ; \\
\hat{M}_6 f &:= f_{t_6} - \frac{i}{360} f_{xxxxxx} + 2c(10cu^3(u_x f_x + u f_{xx}) + 5(2u_x^2 f_{xx} + \\
&4uu_{xx} f_{xx} + u_x(5u_x f_{xx} + 5u_{xx} f_x + 4u_{xx} f_x + 3u f_{xxx}) + (5u_x u_{xx} + \\
&3uu_{xxx})f_x)) + u(70cu^2 u_x f_x + 11u_{xx} f_{xx} + 11f_{xx} u_{xx} + 9u_x f_{xxx} + \\
&4u_x f_{xxx} + 9u_{xx} f_x + 6u f_{xxx}) + u^2(5cu(6f_x u_x + 6f_x u_x + 5u f_{xx}) + \\
&f_{xxxx}) + f(30c^2 j_x j^4 u^2 + 10cu^2(3j_x j^2 + 5uu_{xx}) + 10cu^3 u_{xx} + \\
&5(2u_{xx}^2 + 3u_x u_{xxx}) + u(70cu u_x^2 + 6u_{xxxx})) + f(40c^2 j_x j^2 + 35cu^2 u_x^2 + \\
&11j_{xx} j^2 + 15cu^2(u_x^2 + 2uu_{xx}) + 9u_x u_{xxx} + 4u_x u_{xxx} + 6uu_{xxx} + \\
&2u(5cu(6j_x j^2 + 5uu_{xx}) + u_{xxxx})) : 
\end{aligned} \tag{22}$$

At last, if  $c = 0$ , equations (1) and (17) lead to the linear Schrodinger (LS) equation

$$iu_t + u_{xx} = 0 \tag{23}$$

and to its (trivial) symmetries ( $i^{n+1} \partial_x^n u$ ).

The paper is organized as follows. In §2 we construct the multiscale expansion, in the small regime, of a generic solution of (1), establishing, in particular, that the leading term of such expansion evolves wrt the infinitely many "even" time variables  $t_{2k} := 2^{(k-1)}t$ ;  $k \geq N_+$  according to the flows of the NLS hierarchy. In §3, after summarizing the DP test and after showing how to use the main output of this test to construct infinitely many approximate symmetries of the original PDE through novel and simple formulas, we apply the DP test to the PDE (2), isolating the constraints on the coefficients  $a_j^0$ ;  $j = 1, \dots, 10$  allowing one to pass the test at time scales of  $O(\epsilon^2)$  and of  $O(\epsilon^4)$ , in both scenarios of C- and S-integrability. In §4 we summarize the results of the paper and we discuss the research perspectives opened by this work. In the Appendix (§5) we display the long outputs of the DP test, obtained using the algebraic manipulation program of Mathematica.

## 2 Multiscale expansion in the small lattice spacing regime

If the lattice spacing is small:  $0 < \epsilon \ll 1$ , as consequence of the invariance of (2) under the transformation  $x_{n+1} = x_n + \epsilon$  and of the well-known formula

$$f_{n+1} = \sum_{k=0}^{\infty} \frac{\epsilon^k}{k!} \partial_x^k f; \quad (24)$$

only even  $x$ -derivatives appear at all (even) orders in  $\epsilon$ , implying that also the asymptotic expansion of  $f_n$  contains only even powers of  $\epsilon$ . Consequently, to eliminate the secularities appearing at all even orders in  $\epsilon$ , the coefficients of such expansion must depend on infinitely many "even" slow times [1]:

$$\tau = (t_2; t_4; t_6; \dots); \quad t_{2k} = \epsilon^{2(k-1)} t; \quad k \geq 1; \quad (25)$$

implying that

$$\partial_t = \partial_{t_2} + \epsilon^2 \partial_{t_4} + \epsilon^4 \partial_{t_6} + \dots \quad (26)$$

Therefore we are lead to the following ansatz for the asymptotic expansion of the "generic" solution of (2)

$$f_n(t) = \sum_{k=0}^{\infty} \epsilon^{2k} u^{(2k+1)}(x; \tau); \quad u^{(1)}(x; \tau) = u(x; \tau); \quad (27)$$

Plugging (24), (26) and (27) into equation (2) and equating to zero the coefficients of all powers in  $\epsilon$ , we obtain the following results.

At the leading  $O(1)$ , we obtain the NLS equation for the leading term  $u^{(1)} = u$  wrt the first time  $t_2 = t$ :

$$u_{t_2} = K_2(u); \quad K_2(u) = -\frac{1}{2} u = i(u_{xx} + 2c|u|^2 u); \quad c = \sum_{j=1}^{\infty} a_j \epsilon^{2j} \quad (28)$$

As usual in perturbation theory, at the next relevant order ( $O(\epsilon^2)$  in our case), the "linearization"  $\hat{M}_2 u^{(3)}$  of  $(u_{t_2} - K_2(u))$  appears, together with the linear term  $(u_{t_4} - (i-1)u_{xxxx})$ , coming from the linear part of (2), and with a nonlinear term  $G_5$ , coming from the nonlinear part of (2):

$$\hat{M}_2 u^{(3)} = u_{t_4} - \frac{i}{4!} u_{xxxx} + G_5; \quad (29)$$

where

$$\begin{aligned}
G_5 &= i(s_1 u^2 u_{xx} + s_2 j_1 j_1 u_{xx} + s_3 u j_x j_1 + s_4 u u_x^2); \\
s_1 &= a_3 + a_4 + a_5 + a_8 + a_9 + a_{10}; \\
s_2 &= a_2 + a_4 + a_5 + 2(a_6 + a_7 + a_8 + a_9 + a_{10}); \\
s_3 &= 2(a_4 - a_5 + 2a_8 - 2a_9); \quad s_4 = 2(a_6 - a_7 + a_8 + a_9 - a_{10});
\end{aligned} \tag{30}$$

Concentrating on the linear terms in round bracket, we observe that  $u_{t_4} = 2K_4 u$  and  $(i(2=4!)u_{xxxx})$  is the linear part of the symmetry  $(2=4!)_4(u) = 2K_4 u$ . Therefore, adding and subtracting the symmetry  $(2=4!)_4$ , equation (29) is conveniently rearranged in the following way, isolating the resonant terms in round bracket:

$$\hat{M}_2 u^{(3)} = u_{t_4} + \frac{2}{4!} (u) + g_5; \tag{31}$$

where

$$g_5 = i(c_1 j_1 j_1 u + c_2 u u_x^2 + c_3 u j_x j_1 + c_4 j_1 j_1 u_{xx} + c_5 u^2 u_{xx}) \tag{32}$$

and

$$\begin{aligned}
c_1 &= \frac{1}{2} c^2; \\
c_2 &= \frac{1}{2} (a_1 + a_2 + a_3 + a_4 + a_5 - 3a_6 + 5a_7 - 3a_8 - 3a_9 + 5a_{10}); \\
c_3 &= \frac{1}{2} (a_1 + a_2 + a_3 - 5a_4 + 7a_5 + a_6 + a_7 - 11a_8 + 13a_9 + a_{10}); \\
c_4 &= \frac{1}{3} [2a_1 - a_2 + 2a_3 - a_4 - a_5 - 4(a_6 + a_7 + a_8 + a_9 + a_{10})]; \\
c_5 &= \frac{1}{6} [a_1 + a_2 + a_6 + a_7 - 5(a_3 + a_4 + a_5 + a_8 + a_9 + a_{10})];
\end{aligned} \tag{33}$$

To eliminate the secularity in bracket, we are forced to choose

$$u_{t_4} = K_4(u) = \frac{2}{4!} (u); \tag{34}$$

so that (31) finally becomes the following secularity free equation for the first correction  $u^{(3)}$ :

$$\hat{M}_2 u^{(3)} = g_5; \tag{35}$$

This procedure iterates without essential differences at all orders. The term  $\hat{M}_2 u^{(3)}$  and  $(u_{t_4} - K_4(u))$  in (31) generate, at  $O(4)$ , the terms  $\hat{M}_2 u^{(5)}$  and  $\hat{M}_4 u^{(3)}$  respectively, while the new linear term  $(u_{t_6} - i(2=6!)u_{xxxxxx})$  is rearranged again into the secular factor  $(u_{t_6} - (2=6!)_6)$  that must be set to zero, to avoid secularities. Since, at  $O(2k)$ , we produce the linear term

$u_{t_{2k}} = i \frac{2}{(2k)!} \partial_x^{2k} u$ , one infers, in analogy with [1], that  $u$  evolves wrt to the higher times according to the even flows of the NLS hierarchy as follows:

$$u_{t_{2k}} = K_{2k}(u) := (-1)^{k+1} \frac{2}{(2k)!} \partial_x^{2k}(u); \quad k \geq N_+; \quad (36)$$

and one is left with the following triangular set of equations [2, 3]

$$\begin{aligned} O(2) : & \quad \hat{M}_2 u^{(3)} = g_5; \\ O(4) : & \quad \hat{M}_2 u^{(5)} + \hat{M}_4 u^{(3)} = g_7; \\ O(6) : & \quad \hat{M}_2 u^{(7)} + \hat{M}_4 u^{(5)} + \hat{M}_6 u^{(3)} = g_9; \\ & \quad \vdots \\ O(2k) : & \quad \hat{M}_2 u^{(2k+1)} + \hat{M}_4 u^{(2k-1)} + \dots + \hat{M}_{2k} u^{(3)} = g_{2k+3}; \\ & \quad \vdots \end{aligned} \quad (37)$$

where, f.i., the expression of  $g_7$  is presented in formula (62) of the Appendix. It remains to remark, following [2, 3], that the symmetries  $f_n g$  and the expressions in (37) bring naturally to the definition of the following vector spaces.

**Definitions** Let  $P_n$  be the vector space of all differential polynomials in the functions  $(\partial_x^j u^{(k)})$  and  $(\overline{\partial_x^j u^{(k)}})$  of order  $n$ , possessing the gauge symmetry of first kind, where:

$$\text{order}(\partial_x^j u^{(k)}) = \text{order}(\overline{\partial_x^j u^{(k)}}) = j + k \quad (38)$$

(so that  $\text{order}(\partial_x^j u) = j + 1$ , since  $u = u^{(1)}$ ), and let  $P_n(m)$  be the subspace of  $P_n$  of all differential polynomials in the functions  $(\partial_x^j u^{(k)})$  and  $(\overline{\partial_x^j u^{(k)}})$ , of order  $n$ , possessing the gauge symmetry of first kind and such that  $k \leq m$ . It is easy to see that, for instance,  $n \geq K_n \geq P_{n+1}(1)$ ,  $g_5 \in P_5(1)$  and  $g_7 \in P_7(3)$  (see (62)).

### 3 Applying the DP integrability test

Suppose we generate, from the model to be tested, a NLS-type multiscale expansion (as in our example); then we have the following scenarios. If such a model is S-integrable (C-integrable),

1) the leading term  $u$  of the asymptotic expansion evolves, with respect to

the slow times  $t_n$ , according to the NLS (LS) hierarchy [1];

2) there exist elements  $f_n^{(m)} \in P_{m+n}$  such that the following equations hold [2, 3]:

$$\hat{M}_n u^{(m)} = f_n^{(m)} \in P_{m+n}; \quad m, n \in N_+; \quad (39)$$

implying, due to (21), the compatibility conditions

$$\hat{M}_n f_m^{(j)} = \hat{M}_m f_n^{(j)}; \quad m, n, j \in N_+; \quad (40)$$

Therefore equations (40) are necessary conditions to be satisfied, in cascade, for the model under investigation to be  $S \rightarrow (C \rightarrow)$  integrable; they are also sufficient to guaranty the asymptotic character of the expansion. If equations (40) are satisfied only up to a certain order, the model under investigation is not integrable, being nevertheless "asymptotically integrable up to that order" [2, 3].

### 3.1 The DP test and approximate symmetries

Equations (39), (40), the basic formulae of the DP test, have been derived in [2, 3] as a consequence of the existence of a Lax pair for the starting integrable model. It follows that, if the conditions (39), (40) are satisfied up to a certain order, the equation under scrutiny admits an approximate Lax pair up to that order.

In this subsection we show how to derive the conditions (39), (40) from the existence of infinitely many symmetries of the starting integrable model. This derivation allows one to establish the important relations (to the best of our knowledge so far unknown) between the functions  $f_n^{(m)} \in P_{m+n}$  of the DP test and the symmetries of the starting model. We concentrate our attention on the case of difference equations, but our considerations have general validity.

Let  $u_{t_2} = K_2(u)$  be an integrable model, say, the AL equation (4), and let  $u_{t_{2m}} = K_{2m}(u)$ ;  $m > 2$ , be one of its infinitely many higher order commuting flows (symmetries), reducing, in the continuous limit, to the higher commuting flow  $u_{t_{2m}} = K_{2m}(u)$  of NLS.

On one hand, from equations (25), (27), we have that

$$u_{t_{2m}} = u_{t_{2m}} + \sum_{k=0}^{m-1} (u_{t_{2(m+k+1)}} + u_{t_{2m}}^{(3)}) + \sum_{k=0}^{m-1} \sum_{j=m}^{m+k} u_{t_{2j}}^{(2(m+k-j)+1)}; \quad (41)$$

where  $u_{t_{2m}} = K_{2m}(u)$  (from (36)) and  $u_{t_{2j}}^{(2(m+k-j)+1)} = K_{2j}^0[u^{(2(m+k-j)+1)}] + f_{2j}^{(2(m+k-j)+1)}$ , for some functions  $f_{2j}^{(2(m+k-j)+1)}$  to be specified. On the other hand:

$$K_{2m}(u) = K_{2m}(u) + \sum_{k=1}^{2m} K_{2m}^{(2k)}(u) = K(u) + \sum_{k=1}^{2m} K_{2m}^{(2k)}(u); \quad (42)$$

where  $K_{2m}^{(2k)} \in P_{2(m+k)+1}$ . Equating equations (41) and (42), we infer that  $f_n^{(m)} \in P_{m+n}$ ;  $m, n \in \mathbb{N}_+$  (the basic formula (39) of the DP test), and we also construct the asymptotic expansion of the generic higher order symmetry

$$K_{2m}(u) = K_{2m}(u) + \sum_{k=0}^{2m} K_{2(m+k)}^{(2k)}(u) + K_{2m}^0[u^{(3)}] + f_{2m}^{(3)} + \dots = \sum_{k=0}^{2m} K_{2(m+k)}^{(2k)}(u) + K_{2m}^0[u^{(3)}] + f_{2m}^{(3)} + \dots \quad (43)$$

in terms of the NLS higher order symmetries, of their Frechet derivatives in the direction of the corrections  $u^{(j)}$ ;  $j > 1$  of the leading term  $u$  of the expansion (27), and of the output functions  $f_n^{(m)} \in P_{m+n}$  of the DP test.

Therefore, if  $f_{2n}^{(2k+1)} \in P_{2(k+n)+1}$  exists, but  $f_{2n}^{(2k+3)} \in P_{2(k+n)+3}$  does not,  $n \in \mathbb{N}_+$ , it follows that:

- i) the solution  $u^{(2k+1)}$  of (39) is uniformly bounded and the expansion (27) is asymptotic up to the  $O(2^k)$ ; therefore the P.E. under scrutiny approximates well its continuous limit, with an error of  $O(2^k)$ , for time scales up to the  $O(2^k)$ .
- ii) The P.E. possesses infinitely many "approximate" generalized symmetries in the form (43) up to the  $O(2^k)$ ; therefore it is integrable up to that order. We remark that, due to the Hamiltonian theory of integrable systems [46]–[49], it is also possible to associate with the P.E. infinitely many "approximate" constants of motion in involution, a very useful information in a numerical check.

### 3.2 C- and S-integrability at $O(2)$

In our example, the first of equations (37) is already in the form (39), with  $g_5 = f_2^{(3)} \in P_5(1)$ . Assuming now that  $\hat{M}_4 u^{(3)} = f_4^{(3)}$ , we arrive at the consistency

$$\hat{M}_4 f_2^{(3)} = \hat{M}_2 f_4^{(3)}; \quad (44)$$

that must be viewed as an equation for the unknown  $f_4^{(3)}$ . Since  $g_5 = f_2^{(3)} \in P_5(1)$ , it follows that one must look for  $f_4^{(3)} \in P_7(1)$ . The calculation, plain but lengthy, has been performed using the algebraic manipulation program of Mathematica, and gives the following result.

Lemma 1. Equation (44) admits a unique solution  $f_4^{(3)} \in P_7(1)$  (presented in formula (64) of the Appendix) if the coefficients  $a_j$ 's appearing in (2) satisfy the following quadratic constraint

$$(a_1 - 3a_3 - 2a_4 - 6a_5 - 5a_6 + 3a_7 - 5a_8 - 13a_9 - a_{10}) \left( \sum_{j=1}^{10} a_j \right) = 0; \quad (45)$$

Once  $f_4^{(3)}$  is constructed,  $f_2^{(5)} \in P_7(3)$  ( $f_2^{(5)} = \hat{M}_2 u^{(5)}$ ) is found from the second of equations (37):

$$f_2^{(5)} = g_7 - f_4^{(3)} \quad (46)$$

and is presented in formula (67) of the Appendix.

We first notice the nice factorization of the quadratic constraint (45) into two linear constraints:

$$c = \sum_{j=1}^{10} a_j = 0; \quad (47)$$

$$a_1 - 3a_3 - 2a_4 - 6a_5 - 5a_6 + 3a_7 - 5a_8 - 13a_9 - a_{10} = 0; \quad (48)$$

Therefore we have the following two different scenarios.

1) If the first constraint (47) is satisfied by the coefficients  $a_j$ 's, the continuous limits of dNLS (2) is the linear Schrodinger equation. It follows that, in this case, equation (2) is "asymptotically C-integrable" at  $O(\epsilon^2)$  and one expects that, for generic initial data and at time scales of  $O(\epsilon^{-2})$ , the dynamics according to (2), (47) be well approximated by the dynamics according to the linear Schrodinger equation (23) with an error of  $O(\epsilon^2)$ .

In particular, the dNLS (2), (7) is "asymptotically C-integrable" at  $O(\epsilon^2)$  if:

$$a_1 + 4a_3 + 3a_6 = 0; \quad (49)$$

2) If, instead, the second constraint (48) is satisfied by the coefficients  $a_j$ 's, the dNLS equation (2) is "asymptotically S-integrable" at  $O(\epsilon^2)$  and one expects that, for generic initial data and at time scales of  $O(\epsilon^{-2})$ , the dynamics according to the dNLS equation (2), (48) approximates well the dynamics according to the NLS equation (1), (3) with an error of  $O(\epsilon^2)$ .

In particular, i) the dNLS (2),(6) is "asymptotically S-integrable" at  $O(\epsilon^2)$  i the following additional constraint is satisfied:

$$a_3 + 2a_8 = 0; \quad (50)$$

ii) the dNLS (2),(7) is "asymptotically S-integrable" at  $O(\epsilon^2)$  i the following additional constraint is satisfied:

$$a_1 - 8a_3 - 9a_6 = 0; \quad (51)$$

while the dNLS (2),(8) is not "asymptotically S-integrable" at  $O(\epsilon^2)$  (therefore it is not integrable).

In addition, since the dNLS equation (2) is the linear combination of ten different discretizations of NLS, it is immediate to check if some of these ten discretizations satisfy the constraint (48). Calling dNLS<sub>k</sub> the single discretization of NLS obtained choosing in (2)  $a_j = a_{k-jk}$ ;  $j = 1; \dots; 10$ , it is straightforward to see (since the coefficient  $a_2$  is the only one absent in (48)) that only the dNLS<sub>2</sub> equation (coinciding with the AL equation (4)) satisfies the constraint (48) (as it has to be, being an integrable system). All the other dNLS<sub>k</sub>;  $k \neq 2$  equations, including the dNLS<sub>1</sub> equation (5), do not satisfy the constraint (48); therefore they are not "asymptotically S-integrable" at  $O(\epsilon^2)$  (consequently, they are not integrable) and, for generic initial data and at time scales of  $O(\epsilon^{-2})$ , their dynamics are expected to be quite different from that of NLS (1),(3), presumably exhibiting numerical evidence of nonintegrability and/or chaos.

We finally infer that the discretizations (2),(6) and (2),(7) satisfying respectively the constraints (50) and (51), the AL equation and any other dNLS equation (2) satisfying the constraint (48) are all close to NLS (once the free coefficients of each model are normalized to satisfy (3)) and are all close together at time scales of  $O(\epsilon^{-2})$ , in the sense mentioned in the introduction.

It is interesting now to push the integrability test to the next order. Due to the above factorization of the constraint (45), the test bifurcates and, in the next two subsections, we explore both cases. Before doing that, we observe that, given  $f_2^{(3)} \in P_5(1)$  and assuming that the constraint (45) be satisfied, equation  $\hat{M}_6 f_2^{(3)} = \hat{M}_2 f_6^{(3)}$  admits a unique solution  $f_6^{(3)} = \hat{M}_6 u^{(3)} \in P_9(1)$ , presented in formula (69) of the Appendix, and no additional constraint appears in this derivation, as predicted by the DP test.

### 3.3 C - integrability at $O(\epsilon^4)$

Let us assume that the constraint (47) be satisfied. For the construction of  $f_4^{(5)} = \hat{M}_4 u^{(5)} \in P_9(3)$  from the equation

$$\hat{M}_4 f_2^{(5)} = \hat{M}_2 f_4^{(5)} \quad (52)$$

we have the following result.

**Lemma 2** Equation (52) admits a unique solution  $f_4^{(5)} \in P_9(3)$ , presented in formula (71) of the Appendix, if the coefficients  $a_j$ 's satisfy the four linear constraints (11), defining a 6-parameter family (but one of these parameters can always be rescaled away) of dNLS equations (2) "asymptotically C-integrable" at  $O(\epsilon^4)$ . Therefore one expects that, for generic initial data and at time scales of  $O(\epsilon^{-4})$ , the dynamics according to (2), (11) will approximate the dynamics according to the linear Schrodinger equation (23) with an error of  $O(\epsilon^2)$ .

For instance, the discretization (2), (7) satisfies the constraints (11) if  $a_6 = -a_3 = a_1$ .

The 6-parameter family of dNLS equations (2), (11) (or at least some particular case of it), being C-integrable at such a high order, is a natural candidate to be a C-integrable discrete system.

### 3.4 S - integrability at $O(\epsilon^4)$

Let us assume that the constraint (48) be satisfied. For the construction of a unique  $f_4^{(5)} = \hat{M}_4 u^{(5)} \in P_9(3)$  from equation (52), we have the following result.

**Lemma 3.** If the constraint (48) is satisfied, equation (52) admits a unique solution  $f_4^{(5)} \in P_9(3)$ , presented in formula (71) of the Appendix, if the following five quadratic constraints are satisfied:

$$Q_j = 0; \quad j = 1; \dots; 5; \quad (53)$$

where the  $Q_j$ 's are the following quadratic forms in the 9 variables

$a_2; \dots; a_{10}:$

$$Q_1 = 4a_{10}^2 + a_{10}a_2 + 2a_{10}a_3 - a_2a_3 + 2a_3^2 - a_{10}a_4 - 2a_2a_4 + a_3a_4 + 3a_{10}a_5 - 2a_2a_5 - 3a_3a_5 - 8a_4a_5 - 8a_5^2 + 18a_{10}a_6 + 6a_3a_6 - 6a_{10}a_7 + 4a_2a_7 + 6a_3a_7 + 4a_4a_7 + 20a_5a_7 + 24a_6a_7 - 8a_7^2 + 12a_{10}a_8 - 3a_2a_8 + 6a_3a_8 + 3a_4a_8 - 9a_5a_8 - 6a_6a_8 + 18a_7a_8 + 20a_{10}a_9 - 3a_2a_9 - 2a_3a_9 - 13a_4a_9 - 25a_5a_9 - 6a_6a_9 + 50a_7a_9 - 24a_8a_9 - 24a_9^2; \quad (54)$$

$$Q_2 = 14a_{10}^2 + 6a_{10}a_2 + 44a_{10}a_3 + 4a_2a_3 + 26a_3^2 + 36a_{10}a_4 + 5a_2a_4 + 40a_3a_4 + 17a_4^2 + 72a_{10}a_5 + 7a_2a_5 + 88a_3a_5 + 68a_4a_5 + 75a_5^2 + 64a_{10}a_6 + 10a_2a_6 + 72a_3a_6 + 60a_4a_6 + 128a_5a_6 + 60a_6^2 - 24a_{10}a_7 - 2a_2a_7 - 24a_3a_7 - 16a_4a_7 - 44a_5a_7 - 32a_6a_7 + 4a_7^2 + 64a_{10}a_8 + 8a_2a_8 + 60a_3a_8 + 54a_4a_8 + 106a_5a_8 + 100a_6a_8 - 20a_7a_8 + 42a_8^2 + 168a_{10}a_9 + 28a_2a_9 + 220a_3a_9 + 170a_4a_9 + 382a_5a_9 + 332a_6a_9 - 108a_7a_9 + 284a_8a_9 + 466a_9^2; \quad (55)$$

$$Q_3 = 20a_{10}^2 + 15a_{10}a_2 + 38a_{10}a_3 - 5a_2a_3 - 22a_3^2 + 39a_{10}a_4 - a_2a_4 - 23a_3a_4 - a_4^2 + 63a_{10}a_5 - 11a_2a_5 - 83a_3a_5 - 52a_4a_5 - 75a_5^2 + 70a_{10}a_6 - 14a_2a_6 - 78a_3a_6 - 48a_4a_6 - 148a_5a_6 - 84a_6^2 - 18a_{10}a_7 + 10a_2a_7 + 42a_3a_7 + 32a_4a_7 + 76a_5a_7 + 88a_6a_7 - 20a_7^2 + 88a_{10}a_8 - 7a_2a_8 - 54a_3a_8 - 15a_4a_8 - 119a_5a_8 - 134a_6a_8 + 82a_7a_8 - 36a_8^2 + 72a_{10}a_9 - 59a_2a_9 - 302a_3a_9 - 211a_4a_9 - 539a_5a_9 - 526a_6a_9 + 234a_7a_9 - 472a_8a_9 - 788a_9^2; \quad (56)$$

$$Q_4 = 32a_{10}^2 - 24a_{10}a_2 - 56a_{10}a_3 + 6a_2a_3 + 36a_3^2 - 70a_{10}a_4 - a_2a_4 + 30a_3a_4 + a_4^2 - 114a_{10}a_5 + a_2a_5 + 78a_3a_5 + 20a_4a_5 + 27a_5^2 - 120a_{10}a_6 + 14a_2a_6 + 88a_3a_6 + 48a_4a_6 + 84a_5a_6 + 84a_6^2 + 24a_{10}a_7 - 14a_2a_7 - 64a_3a_7 - 52a_4a_7 - 80a_5a_7 - 112a_6a_7 + 28a_7^2 - 164a_{10}a_8 + 2a_2a_8 + 48a_3a_8 + 12a_4a_8 + 16a_5a_8 + 124a_6a_8 - 116a_7a_8 + 36a_8^2 - 220a_{10}a_9 + 22a_2a_9 + 208a_3a_9 + 96a_4a_9 + 196a_5a_9 + 292a_6a_9 - 204a_7a_9 + 176a_8a_9 + 300a_9^2; \quad (57)$$

$$Q_5 = 4a_{10}^2 + 3a_{10}a_2 - 2a_{10}a_3 - a_2a_3 - 14a_3^2 + 3a_{10}a_4 - a_2a_4 - 19a_3a_4 - 5a_4^2 - 5a_{10}a_5 - 3a_2a_5 - 47a_3a_5 - 36a_4a_5 - 39a_5^2 - 2a_{10}a_6 - 6a_2a_6 - 38a_3a_6 - 32a_4a_6 - 68a_5a_6 - 36a_6^2 + 6a_{10}a_7 + 2a_2a_7 + 18a_3a_7 + 16a_4a_7 + 28a_5a_7 + 24a_6a_7 - 4a_7^2 + 8a_{10}a_8 - 3a_2a_8 - 30a_3a_8 - 19a_4a_8 - 59a_5a_8 - 62a_6a_8 + 26a_7a_8 - 20a_8^2 - 40a_{10}a_9 - 23a_2a_9 - 150a_3a_9 - 119a_4a_9 - 255a_5a_9 - 230a_6a_9 + 82a_7a_9 - 216a_8a_9 - 356a_9^2; \quad (58)$$

The homogeneous quadratic constraints (53), (54)–(58) for nine unknowns, characterizing the intersection of 5 quadrics in the real projective space of dimension 8, define, in principle, a 4-parameter family of solutions (but one of these parameters can always be rescaled away) whose parametrization does not appear to be expressible, in general, in terms of elementary functions. The corresponding dNLS equation (2) is asymptotically S-integrable at  $O(\epsilon^4)$  and should well approximate the NLS equation for times up to the  $O(\epsilon^4)$ .

We observe that, in all these quadratic constraints,  $a_2$  is the only coefficient appearing always multiplied by other coefficients (the term  $(a_2)^2$  is absent); therefore the choice

$$a_j = a_2 \cdot j; \quad j = 1; \dots; 10; \quad (59)$$

corresponding to the AL equation (4), satisfies all constraints, as it has to be. Other less trivial explicit solutions of (48), (53), (54)–(58) can also be constructed, corresponding to the case in which all quadrics degenerate into pairs of hyperplanes. Here we display the following two examples:

$$\begin{aligned} a_1 &= 4a_6; \quad a_2 = \frac{4a_6}{3}; \quad a_3 = 4a_6; \quad a_4 = 0; \\ a_5 &= 4a_6; \quad a_7 = a_6; \quad a_8 = a_9 = a_{10} = 0; \end{aligned} \quad (60)$$

$$\begin{aligned} a_1 &= 24a_9; \quad a_2 = a_3 = 0; \quad a_4 = a_5 = 8a_9; \\ a_6 &= 10a_9; \quad a_7 = 2a_9; \quad a_8 = 7a_9; \quad a_{10} = 6a_9; \end{aligned} \quad (61)$$

corresponding, respectively, to the dNLS equations (12) and (13) presented in the Introduction, "asymptotically S-integrable" at  $O(\epsilon^4)$ . Therefore one expects that, for generic initial data and at time scales of  $O(\epsilon^4)$ , the dynamics according to equations (12) and (13) are good approximations of the dynamics according to the NLS equation (1), with an error of  $O(\epsilon^2)$ , at time scales of  $O(\epsilon^4)$ . Of course, these distinguished equations, passing the test at such a high order, are also good candidates to be S-integrable difference equations.

We finally observe that there is no choice of parameters for which the dNLS equations (2), (6) and (2), (7) satisfy the above constraints; therefore these two models are not S-integrable at this order (they are not S-integrable at all) and do not approximate well NLS at time scales of  $O(\epsilon^4)$ .

## 4 Summary of the results and future perspectives

In this paper we have proposed an algorithmic procedure allowing one: i) to study the distance between an integrable PDE and any PDE discretizing it, in the small lattice spacing regime; ii) to test the (asymptotic) integrability properties of such a PDE, and iii) to construct infinitely many (approximate) symmetries and conserved quantities for it. This method should provide, in particular, useful and concrete informations on how good is a numerical scheme used to integrate a given integrable PDE.

The procedure we have proposed, illustrated on the basic prototype example of the nonlinear Schrödinger equation (1) and of its discretization (2), consists in the following three steps: i) the construction of the multiscale expansion of a generic solution of the dNLS (2) at all orders in  $\epsilon$ , following [1]; ii) the application, to such expansion, of the DP integrability test [2, 3]; iii) the use the main output of such test to construct infinitely many approximate symmetries of the dNLS equation (2), through novel formulas presented in this paper.

This approach allows one to study the distance between the integrable PDE and any PDE discretizing it. Suppose, for instance, that the asymptotic expansion we construct reads  $\psi = u + O(\epsilon^k)$ ;  $k > 0$ , where  $\psi$  is a generic solution of the dNLS (2) and  $u$  is the corresponding solution of (1), then if the DP test is passed at  $O(\epsilon^k)$ , we conclude that: i) the dynamics according to the NLS equation (1) is well approximated (with an error of  $O(\epsilon^k)$ ) by the dynamics according to its discretization (2), for time scales of  $O(\epsilon^{-k})$ ; ii) the dNLS equation is asymptotically integrable up to that order, constructing its infinitely many approximate symmetries and constants of motion in involution. On the contrary, if the DP test is not passed at that order, the dNLS equation is not integrable and one should expect, at the corresponding time scale, numerical evidence of nonintegrability and/or chaos.

We have carried the above procedure up to the  $O(\epsilon^4)$  and we have been able to isolate the constraints on the coefficients of the dNLS equation (2) allowing one to pass the test at that order, in both scenarios of S- and C-integrability.

Numerical experiments to test such theoretical findings are presently under investigation; preliminary results seem to confirm the theoretical predictions contained in this paper [51].

With the same methodology and goals, we are presently investigating families of discretizations of the Korteweg-de Vries and Burgers equations [52], other two basic integrable models of natural phenomena. Of course we also plan to investigate discretizations of integrable PDEs in which also the time variable is discretized.

## 5 Appendix

In this Appendix we display, for completeness, the long outputs of the DP test, obtained using the algebraic manipulation program of Mathematica.

The differential polynomial  $g_7$  in (37) reads:

$$\begin{aligned}
g_7 = & i_1 u_j f + i_2 j f u^{(3)} + i_3 u u^{(3)^2} + i_4 u^2 j f u^{(3)} + i_5 u j^{(3)} f + \\
& i_6 u j f u_x^2 + i_7 u_x^2 u^{(3)} + i_8 u j f j u_x f + i_9 j u_x f u^{(3)} + i_{10} u^3 u_x^2 + \\
& i_{11} u u_x u_x^{(3)} + i_{12} u u_x u_x^{(3)} + i_{13} u u_x u_x^{(3)} + i_{14} j f u_{xx} + i_{15} u u_{xx} u^{(3)} + \\
& i_{16} u u_{xx} u^{(3)} + i_{17} j u_x f u_{xx} + i_{18} u u_{xx}^2 + i_{19} u^2 j f u_{xx} + i_{20} u u_{xx} u^{(3)} + \\
& i_{21} u_x^2 u_{xx} + i_{22} u j u_{xx} f + i_{23} j f u_{xx} + i_{24} u^2 u_{xx}^{(3)} + i_{25} u u_x u_{xxx} + \\
& i_{26} u u_x u_{xxx} + i_{27} u u_x u_{xxx} + i_{28} j f u_{xxx} + i_{29} u^2 u_{xxx} ;
\end{aligned} \tag{62}$$

where

$$\begin{aligned}
i_1 = & \frac{1}{18} c^3; \quad i_2 = \frac{3}{2} c^2; \quad i_3 = 2c; \quad i_4 = c^2; \quad i_5 = 4c; \quad i_6 = 7i_{10}; \quad i_7 = \frac{1}{2} i_{11}; \\
i_8 = & 6i_{10}; \quad i_9 = \frac{1}{3} (a_1 + a_2 + a_3 - 5a_4 + 7a_5 + a_6 + a_7 - 11a_8 + 13a_9 + a_{10}); \\
i_{10} = & \frac{1}{36} c^2; \quad i_{11} = (a_1 + a_2 + a_3 + a_4 + a_5 - 3a_6 + 5a_7 - 3(a_8 + a_9) + 5a_{10}); \\
i_{12} = & i_{13}; \quad i_{13} = \frac{1}{3} (a_1 + a_2 + a_3 - 5a_4 + 7a_5 + a_6 + a_7 - 11a_8 + 13a_9 + a_{10}); \\
i_{14} = & 5i_{10}; \quad i_{15} = i_{16}; \\
i_{16} = & \frac{1}{3} (2a_1 - a_2 + 2a_3 - a_4 - a_5 - 4(a_6 + a_7 + a_8 + a_9 + a_{10})); \\
i_{17} = & \frac{1}{36} (5(a_1 + a_2 + a_3 + a_4 + a_5 + a_6 + a_7 + a_{10}) - 67a_8 + 77a_9); \\
i_{18} = & \frac{1}{18} (a_1 + a_2 + a_3 + a_4 + a_5 - 8(a_6 + a_7 + a_8 + a_9 + a_{10})); \\
i_{19} = & 2i_{10}; \quad i_{20} = \frac{1}{3} (a_1 + a_2 + a_6 + a_7 - 5(a_3 + a_4 + a_5 + a_8 + a_9 + a_{10})); \\
i_{21} = & \frac{1}{18} (a_1 + a_2 + a_3 + a_4 + a_5 + a_6 + a_7 - 17(a_8 + a_9) + 19a_{10}); \\
i_{22} = & \frac{1}{180} (11(a_1 + a_2 + a_3) - 79(a_4 + a_5) + 11(a_6 + a_7) - 169(a_8 + a_9 + a_{10})); \\
i_{23} = & \frac{1}{3} (2a_1 - a_2 + 2a_3 - a_4 - a_5 - 4(a_6 + a_7 + a_8 + a_9 + a_{10})); \quad i_{24} = \frac{1}{2} i_{20}; \\
i_{25} = & \frac{1}{12} (a_1 + a_2 + a_3 + a_4 + a_5 + 9a_7 - 7(a_6 + a_8 + a_9) + 9a_{10}); \\
i_{26} = & \frac{1}{60} (3(a_1 + a_2 + a_3 + a_{10}) - 17a_4 + 23a_5 + 3a_6 + 3a_7 - 37a_8 + 43a_9); \\
i_{27} = & \frac{1}{45} (a_1 + a_2 + a_3 - 14a_4 + 16a_5 + a_6 + a_7 - 29a_8 + 31a_9 + a_{10}); \\
i_{28} = & \frac{1}{60} (2a_1 - 3a_2 + 2a_3 - 3a_4 - 3a_5 - 8(a_6 + a_7 + a_8 + a_9 + a_{10})); \\
i_{29} = & \frac{1}{180} (a_1 + a_2 + a_6 + a_7 - 14(a_3 + a_4 + a_5 + a_8 + a_9 + a_{10}));
\end{aligned} \tag{63}$$

The solution  $f_4^{(3)} \in P_7(1)$  of  $\hat{M}_4 f_2^{(3)} = \hat{M}_2 f_4^{(3)}$ , where  $f_2^{(3)} = g_5$  is given in (32), (33), exists

unique and reads

$$f_4^{(3)} = i_1 u j j + i_2 u_{xx} j j^4 + i_3 u_{xx} u^2 j j^2 + i_4 u_x^2 j j^2 u + i_5 j j^2 j j^2 u + i_6 u_x^2 u^3 + i_7 u_{xxxx} j j^2 + i_8 u_{xxxx} u^2 + i_9 u_{xxx} u_x u + i_{10} u_{xxx} u_x u + i_{11} u_{xxx} u_x u + i_{12} u_{xx}^2 u + i_{13} j j_{xx} j^2 u + i_{14} u_{xx} j j_x j^2 + i_{15} u_{xx} u_x^2 ; \quad (64)$$

where

$$\begin{aligned} i_1 &= \frac{c_2^2}{3} (2c_2 - c_3 + c_4 + 3c_5); & i_2 &= \frac{c}{6} (4c_2 - 2c_3 + 6c_4 + 5c_5); \\ i_3 &= \frac{c}{12} (2c_2 - c_3 + 3c_4 + 10c_5); & i_4 &= \frac{c}{24} (40c_2 - 11c_3 + 25c_4 + 20c_5); \\ i_5 &= \frac{c}{12} (8c_2 + 5c_3 + 7c_4 + 16c_5); & i_6 &= \frac{c}{24} (4c_2 + c_3 + c_4 + 8c_5); \\ i_7 &= \frac{c_4}{6}; & i_8 &= \frac{c_5}{12}; & i_9 &= \frac{1}{12} (4c_2 + 3c_4); & i_{10} &= \frac{1}{12} (c_3 + 2c_5); \\ i_{11} &= \frac{1}{12} (2c_3 + c_4); & i_{12} &= \frac{1}{12} (3c_2 + 2c_4); & i_{13} &= \frac{1}{12} (c_3 + c_4 + 4c_5); \\ i_{14} &= \frac{1}{12} (2c_2 + 5c_3 + c_4); & i_{15} &= \frac{1}{12} (c_2 + c_3 + 3c_5); \end{aligned} \quad (65)$$

the following constraint is satisfied

$$2c_1 - c(2c_2 - c_3 + c_4 + 4c_5) = 0 \quad (66)$$

on the coefficients  $c_j$ 's defined in (33). This constraint is equivalent to (45).

$f_2^{(5)} = g_7 - f_4^{(3)}$  consequently reads, from (62) and (64):

$$\begin{aligned} f_2^{(5)} &= i_1 d_1 u j j + d_2 j j^4 u^{(3)} + d_3 u u^{(3)^2} + d_4 u^2 j j^2 u^{(3)} + d_5 u j j^{(3)} j^2 + \\ &d_6 u j j^2 u_x^2 + d_7 u_x^2 u^{(3)} + d_8 u j j^2 j j_x j^2 + d_9 j j_x j^2 u^{(3)} + d_{10} u^3 u_x^2 + d_{11} u u_x u_x^{(3)} + \\ &d_{12} u u_x u_x^{(3)} + d_{13} u u_x u_x^{(3)} + d_{14} j j^4 u_{xx} + d_{15} u u_{xx} u^{(3)} + d_{16} u u_{xx} u^{(3)} + \\ &d_{17} j j_x j^2 u_{xx} + d_{18} u u_{xx}^2 + d_{19} u^2 j j^2 u_{xx} + d_{20} u u_{xx} u^{(3)} + d_{21} u_x^2 u_{xx} + d_{22} u j j_{xx} j^2 + \\ &d_{23} j j^2 u_{xx}^{(3)} + d_{24} u^2 u_{xx}^{(3)} + d_{25} u u_x u_{xxx} + d_{26} u u_x u_{xxx} + d_{27} u u_x u_{xxx} + \\ &d_{28} j j^2 u_{xxxx} + d_{29} u^2 u_{xxxx} ; \end{aligned} \quad (67)$$

where

$$\begin{aligned}
d_1 &= \frac{c^2}{9} (5a_1 + 2a_2 - 4a_3 - a_4 - 13a_5 - 13a_6 + 11a_7 - 10a_8 - 34a_9 + 2a_{10}); \\
d_2 &= \frac{3}{2}c^2; \quad d_3 = 2c; \quad d_4 = -c^2; \quad d_5 = 4c; \\
d_6 &= \frac{c}{72} (95a_1 + 20a_2 + 35a_3 + 26a_4 - 106a_5 - 295a_6 + 185a_7 - 223a_8 - 487a_9 + \\
&125a_{10}); \quad d_7 = \frac{1}{2} (a_1 + a_2 + a_3 + a_4 + a_5 - 3a_6 + 5a_7 - 3(a_8 + a_9) + 5a_{10}); \\
d_8 &= \frac{c}{12} (11a_1 + 4a_2 - 5a_3 - 22a_4 - 2a_5 - 19a_6 + 13a_7 - 55a_8 - 15a_9 - 3a_{10}); \\
d_9 &= d_{13}; \quad d_{10} = \frac{c}{72} (11a_1 + 8a_2 - 13a_3 - 22a_4 - 10a_5 - 19a_6 + 29a_7 - 55a_8 \\
&31a_9 + 5a_{10}); \quad d_{11} = 2d_7; \quad d_{12} = d_{13}; \\
d_{13} &= \frac{1}{3} (a_1 + a_2 + a_3 - 5a_4 + 7a_5 + a_6 + a_7 - 11a_8 + 13a_9 + a_{10}); \\
d_{14} &= \frac{c}{18} (16a_1 - 2a_2 + a_3 - 5a_4 - 29a_5 - 44a_6 + 4a_7 - 35a_8 - 83a_9 - 11a_{10}); \\
d_{15} &= d_{16}; \quad d_{16} = \frac{1}{3} (2a_1 - a_2 + 2a_3 - a_4 - a_5 - 4(a_6 + a_7 + a_8 + a_9 + a_{10})); \\
d_{17} &= \frac{1}{36} (5a_1 + 2a_2 + 5a_3 - 28a_4 + 32a_5 - 13a_6 + 11a_7 - a_8 - 25a_9 + 11a_{10}); \\
d_{18} &= \frac{1}{72} (13a_1 + a_2 + 13a_3 + a_4 + a_5 - 11a_6 + 61a_7 - 11(a_8 + a_9) + 61a_{10}); \\
d_{19} &= \frac{c}{36} (11a_1 + 2a_2 - 19a_3 - 22a_4 - 34a_5 - 19a_6 + 5a_7 - 37a_8 - 61a_9 - 25a_{10}); \\
d_{20} &= \frac{1}{3} (a_1 + a_2 + a_6 + a_7 - 5(a_3 + a_4 + a_5 + a_8 + a_9 + a_{10})); \\
d_{21} &= \frac{1}{36} (2a_1 + 2a_2 - 7a_3 - 13a_4 - a_5 - 4a_6 + 8a_7 + 11a_8 + 35a_9 - 37a_{10}); \\
d_{22} &= \frac{1}{180} (14a_1 - a_2 - 46a_3 - a_4 + 59a_5 - 16a_6 - 16a_7 + 44a_8 + 164a_9 + \\
&104a_{10}); \\
d_{23} &= \frac{1}{3} (2a_1 - a_2 + 2a_3 - a_4 - a_5 - 4(a_6 + a_7 + a_8 + a_9 + a_{10})); \quad d_{24} = \frac{1}{2}d_{20}; \\
d_{25} &= \frac{1}{4} (a_1 + a_3 - a_6 - a_7 - a_8 - a_9 - a_{10}); \\
d_{26} &= \frac{1}{180} (11a_1 - 4a_2 + 11a_3 - 4a_4 - 4a_5 - 19(a_6 + a_7 + a_8 + a_9 + a_{10})); \\
d_{27} &= \frac{1}{30} (a_1 + a_2 - 4a_3 + a_4 - 9a_5 + a_6 + a_7 + 6a_8 - 14a_9 - 4a_{10}); \\
d_{28} &= \frac{1}{180} (14a_1 - a_2 + 14a_3 - a_4 - a_5 - 16(a_6 + a_7 + a_8 + a_9 + a_{10})); \\
d_{29} &= \frac{c}{120};
\end{aligned} \tag{68}$$

The unique solution  $f_6^{(3)} = \hat{M}_6 u^{(3)} \in P_9(1)$  of equation  $\hat{M}_6 f_2^{(3)} = \hat{M}_2 f_6^{(3)}$  reads:

$$\begin{aligned}
f_6^{(3)} &= i_1 j_1 f_1 u + i_2 u_{xx} j_1 f_1 + i_3 u_{xx} j_1 f_1 u^2 + i_4 u_x^2 j_1 f_1 u + i_5 j_1 x j_1 f_1 u + \\
&i_6 u_x^2 u^3 j_1 f_1 + i_7 u_{xxxx} j_1 f_1 + i_8 u_{xxx} u_x j_1 f_1 u + i_9 u_{xxx} u_x j_1 f_1 u + i_{10} u_{xx}^2 j_1 f_1 u + \\
&i_{11} j_1 x x j_1 f_1 u + i_{12} u_{xxx} u_x j_1 f_1 u + i_{13} u_{xxx} u_x u^3 + i_{14} u_{xx} u_x^2 u^2 + \\
&i_{15} u_{xx} j_1 x j_1 f_1 + i_{16} u_{xx} u_x^2 u^2 + i_{17} u_x^3 u_x u + i_{18} j_1 x f_1 u + i_{19} (u_{xx})^2 u^3 + \\
&i_{20} u_{xx} j_1 x f_1 u^2 + i_{21} u_{xx} u_x^2 j_1 f_1 + i_{22} u_{xxxxxx} j_1 f_1 + i_{23} u_{xxxxxx} u^2 + \\
&i_{24} u_{xxxxxx} j_1 f_1 + i_{25} u_{xxxxxx} u u_x + i_{26} u_{xxxxxx} j_1 x f_1 + i_{27} u_{xxxxxx} u_{xx} u + \\
&i_{28} u_{xxxxxx} u_{xx} u + i_{29} u_{xxxxxx} u_x^2 + i_{30} u_{xxxxxx} u_{xx} u + i_{31} u_{xxx}^2 u + i_{32} u_{xxx} u_{xx} u_x + \\
&i_{33} u_{xxx} u_{xx} u_x + i_{34} j_1 x x x f_1 u + i_{35} j_1 x x f_1 u_{xx} + i_{36} u_{xxxxxx} u_x u + \\
&i_{37} u_{xx} u_{xxx} u_x + i_{38} u_{xxxxxx} j_1 f_1 u^2;
\end{aligned} \tag{69}$$

where

$$\begin{aligned}
1 &= \frac{c^3}{48} (6c_2 - 3c_3 + 3c_4 + 8c_5); & 2 &= \frac{c^2}{36} (10c_2 - 5c_3 + 10c_4 + 12c_5); \\
3 &= \frac{c^2}{36} (4c_2 - 2c_3 + 4c_4 + 9c_5); & 4 &= \frac{c^2}{18} (13c_2 - 5c_3 + 9c_4 + 10c_5); \\
5 &= \frac{c^2}{36} (18c_2 - 3c_3 + 14c_4 + 28c_5); & 6 &= \frac{c^2}{36} (5c_2 - c_3 + 3c_4 + 8c_5); \\
7 &= \frac{c}{180} (6c_2 - 3c_3 + 15c_4 + 7c_5); & 8 &= \frac{c}{360} (108c_2 - 29c_3 + 126c_4 + 56c_5); \\
9 &= \frac{c}{360} (36c_2 + 7c_3 + 66c_4 + 52c_5); & 10 &= \frac{c}{720} (152c_2 - 41c_3 + 169c_4 + 84c_5); \\
11 &= \frac{c}{360} (44c_2 - 7c_3 + 79c_4 + 118c_5); & 12 &= \frac{c}{360} (16c_2 + 7c_3 + 26c_4 + 62c_5); \\
13 &= \frac{c}{360} (8c_2 + c_3 + 6c_4 + 26c_5); & 14 &= \frac{c}{72} (33c_2 - 7c_3 + 21c_4 + 14c_5); \\
15 &= \frac{c}{360} (278c_2 + 11c_3 + 276c_4 + 236c_5); & 16 &= \frac{c}{360} (39c_2 + 8c_3 + 33c_4 + 68c_5); \\
17 &= \frac{c}{72} (22c_2 - c_3 + 13c_4 + 16c_5); & 18 &= \frac{c}{720} (158c_2 + 31c_3 + 101c_4 + 176c_5); \\
19 &= \frac{c}{720} (12c_2 - c_3 + 9c_4 + 44c_5); & 20 &= \frac{c}{120} (16c_2 + 7c_3 + 12c_4 + 42c_5); \\
21 &= \frac{c}{360} (114c_2 - 12c_3 + 103c_4 + 158c_5); & 22 &= \frac{c_4}{120}; & 23 &= \frac{c_5}{360}; \\
24 &= \frac{1}{240} (2c_3 + 3c_4); & 25 &= \frac{1}{360} (c_3 + 4c_5); & 26 &= \frac{1}{720} (18c_2 + 21c_3 + 25c_4); \\
27 &= \frac{1}{360} (15c_2 + 13c_4); & 28 &= \frac{1}{720} (9c_3 + 11c_4 + 12c_5); \\
29 &= \frac{1}{360} (c_2 + 2c_3 + 10c_5); & 30 &= \frac{1}{360} (2c_3 + c_4 + 11c_5); \\
31 &= \frac{1}{144} (4c_2 + 3c_4); & 32 &= \frac{1}{720} (50c_2 + 35c_3 + 34c_4); \\
33 &= \frac{1}{360} (11c_2 + 17c_3 + 10c_4 + 15c_5); & 34 &= \frac{1}{720} (11c_3 + 4c_4 + 18c_5); \\
35 &= \frac{1}{720} (20c_2 + 25c_3 + 11c_4 + 20c_5); \\
36 &= \frac{1}{240} (4c_2 + 5c_4); & 37 &= \frac{1}{720} (8c_2 + 31c_3 + 4c_4 + 50c_5); \\
38 &= \frac{c}{360} (2c_2 - c_3 + 5c_4 + 14c_5);
\end{aligned} \tag{70}$$

and no additional constraint on the coefficients  $c_j$ 's appears.

The unique solution  $f_4^{(5)} = \hat{M}_4 u^{(5)} \in P_9(3)$  of equation  $\hat{M}_4 f_2^{(5)} = \hat{M}_2 f_4^{(5)}$  reads

$$\begin{aligned}
f_4^{(5)} = & i_1 u j j^2 + i_2 u_{xx} j j^2 + i_3 u_{xx} u^2 j j^2 + i_4 u_x^2 u j j^2 + i_5 j j^2 u j j^2 + \\
& i_6 u_x^2 j j^2 u^3 + i_7 u_{xxxx} j j^2 + i_8 u_{xxx} u_x j j^2 u + i_9 u_{xxx} u_x j j^2 u + i_{10} u_{xx}^2 j j^2 u + \\
& i_{11} j j^2 u j j^2 u + i_{12} u_{xxx} u_x j j^2 u + i_{13} u_{xxx} u_x u^3 + i_{14} u_{xx} u_x^2 u^2 + i_{15} u_{xx} j j^2 u j j^2 + \\
& i_{16} u_{xx} u_x^2 u^2 + i_{17} u_x^3 u_x u + i_{18} j j^2 u^4 + i_{19} u_{xx}^2 u^3 + i_{20} u_{xx} j j^2 u^2 + i_{21} u_{xx} u_x^2 j j^2 + \\
& i_{22} u_{xxxxxx} j j^2 + i_{23} u_{xxxxxx} u^2 + i_{24} u_{xxxxxx} u_x u + i_{25} u_{xxxxxx} u_x u + i_{26} u_{xxxxxx} j j^2 + \\
& i_{27} u_{xxxxxx} u_{xx} u + i_{28} u_{xxxxxx} u_{xx} u + i_{29} u_{xxxxxx} u_x^2 + i_{30} u_{xxxxxx} u_{xx} u + i_{31} u_{xx}^2 u + \\
& i_{32} u_{xxx} u_{xx} u_x + i_{33} u_{xxx} u_{xx} u_x + i_{34} j j^2 u_{xxx} + i_{35} u_{xx} j j^2 u_{xx} + i_{36} u_{xxxxxx} u_x u + \\
& i_{37} u_{xx} u_{xxxx} u_x + i_{38} u^2 j j^2 u_{xxxx} + i_{39}^{(3)} u_{xxx} j j^2 + i_{40}^{(3)} u_{xxx} u_x u + i_{41}^{(3)} u_{xxx} u_x u + \\
& i_{42}^{(3)} u_{xx} u_{xx} u + i_{43}^{(3)} u_{xx} j j^2 + i_{44}^{(3)} u_{xx} u u_{xx} + i_{45}^{(3)} u_x u_{xxx} u + i_{46}^{(3)} u_{xxx} u + i_{47}^{(3)} u_{xx} u_x + \\
& i_{48}^{(3)} u_x u_x u_{xx} + i_{49}^{(3)} u_x u u_{xxx} + i_{50}^{(3)} u_{xxx} u + i_{51}^{(3)} u_{xxx} u_x + i_{52}^{(3)} j j^2 u_{xxx} + \\
& i_{53}^{(3)} u_x u_{xxx} + i_{54}^{(3)} u u_{xxxx} + i_{55}^{(3)} u_{xxxx} u^2 + i_{56}^{(3)} u_{xxx} u_x u + i_{57}^{(3)} u_{xx} u_{xx} u + \\
& i_{58}^{(3)} u_{xx} u_x^2 + i_{59}^{(3)} u_{xxx} u + i_{60}^{(3)} u_{xx} u_x + i_{61}^{(3)} u_{xxxx} u + i_{62}^{(3)} u_{xxx} u_x + \\
& i_{63}^{(3)} u_{xx}^2 + i_{64}^{(3)} j j^2 + i_{65}^{(3)} u_x j j^2 u + i_{66}^{(3)} u_x j j^2 u + \\
& i_{67}^{(3)} u_{xx} j j^2 u + i_{68}^{(3)} u_x^2 u^2 + i_{69}^{(3)} j j^2 u j j^2 + i_{70}^{(3)} u_x^2 u^2 + i_{71}^{(3)} u_{xx} j j^2 u + \\
& i_{72}^{(3)} u_{xx} j j^2 u^2 + i_{73}^{(3)} u_x j j^2 u + i_{74}^{(3)} u_x u^3 + i_{75}^{(3)} u_{xx} j j^2 u + i_{76}^{(3)} u_x^2 j j^2 + \\
& i_{77}^{(3)} j j^2 u^2 + i_{78}^{(3)} u_{xx} u^3 + i_{79}^{(3)} j j^2 + i_{80}^{(3)} j j^2 u^2 + i_{81}^{(3)} u_{xx} u^{(3)} u + \\
& i_{82}^{(3)} u^{(3)} u + i_{83}^{(3)} u^{(3)} u_x + i_{84}^{(3)} u_{xx}^{(3)} + i_{85}^{(3)} u_{xx} u^{(3)} u + i_{86}^{(3)} j j^2 u + \\
& i_{87}^{(3)} u^{(3)} u_x + i_{88}^{(3)} u_{xx}^{(3)} u + i_{89}^{(3)} u_x^{(3)} u_x + i_{90}^{(3)} j j^2 u_{xx} + i_{91}^{(3)} (u^{(3)})^2 j j^2 u + \\
& i_{92}^{(3)} j j^2 j j^2 u + i_{93} (u^{(3)})^2 u^3 ;
\end{aligned}$$

(71)

where:

$$\begin{aligned}
 1 &= \frac{c}{576} 336d_1 + 4(6c_2 - 3c_3 + 2c_4 + 10c_5)d_2 + 6c_5(2c_2 - c_3 + 2c_4 + 4c_5)d_3 + \\
 &4(6c_2 - 3c_3 + 3c_4 + 2c_5)d_4 + (4c_4^2 + 2c_2c_5 - c_3c_5 + c_4c_5 - 2c_5^2)d_5 + \\
 &c(2(14c_2 - 5c_3 + 9c_4 + 28c_5)d_{11} - 2(10c_2 - 3c_3 + 7c_4 + 8c_5)d_{12} - 4(6c_2 - \\
 &3c_3 + c_4 + 16c_5)d_{13} + 16d_{14} + 2(18c_2 - 7c_3 + 11c_4 + 30c_5)d_{15} - 2(6c_2 - \\
 &3c_3 + 3c_4 + 14c_5)d_{16} - 48d_{19} - 4(6c_2 - 3c_3 + 3c_4 + 8c_5)d_{20} + 4(4c_2 + c_3 + \\
 &2c_4 + 15c_5)d_{23} + 4(6c_2 + 5c_3 + c_4 - 6c_5)d_{24} + 32d_6 - 16d_8 + 4(2c_2 - c_3 + \\
 &c_4 + 4c_5)d_9) + 8c^2(-2d_{17} + 4d_{18} + 2d_{22} + 7d_{25} - 7d_{26} - 5d_{28} + 16d_{29}) ; \\
 2 &= \frac{1}{144} 96d_1 + 10c_4d_2 + 4c_4c_5d_3 - 6c_5d_4 - 2c_4^2d_5 + c(2(4c_2 - c_3 + 6c_4 + \\
 &8c_5)d_{11} - 2(2c_2 + 5c_4)d_{12} - 2(6c_2 - 3c_3 + c_4 + 14c_5)d_{13} + 104d_{14} + (18c_2 - \\
 &7c_3 + 11c_4 + 24c_5)d_{15} + 3(2c_2 + c_3 - c_4 - 4c_5)d_{16} - 24d_{19} + (34c_2 - 19c_3 + \\
 &31c_4 + 80c_5)d_{23} + 2(6c_2 + 5c_3 - 5c_4 - 10c_5)d_{24} + 16d_6 - 8d_8 + 2(2c_2 - \\
 &c_3 + c_4 + 4c_5)d_9) + 4c^2(-2d_{17} + 4d_{18} + 2d_{22} + 13d_{25} - 13d_{26} + d_{28} + 4d_{29}) ; \\
 3 &= \frac{1}{48} (12d_1 + 2c_5d_2 + 2c_4d_4 + c(2c_5d_{11} - 2c_5d_{12} + 8d_{14} + 24d_{19} + (2c_2 - \\
 &c_3 + c_4 + 4c_5)d_{20} + 2(2c_2 - c_3 + c_4 + 5c_5)d_{23} + 5(2c_2 - c_3 + c_4)d_{24}) + \\
 &8c^2(d_{25} - d_{26} + d_{28} - 2d_{29})); \\
 4 &= \frac{1}{144} 216d_1 + 2(3c_2 + 6c_4 - 2c_5)d_2 + 2(2c_3c_4 + 2c_2c_5 + c_3c_5 + c_4c_5 - \\
 &2c_5^2)d_3 - 24c_5d_4 - (6c_4^2 - 3c_3c_5 + c_4c_5 + 2c_5^2)d_5 + c(24d_{10} + (52c_2 - 19c_3 + \\
 &9c_4 + 54c_5)d_{11} + 2(7c_2 + 3c_3 - 8c_4 + 9c_5)d_{12} + 2(10c_2 + 4c_3 - 5c_4 - \\
 &22c_5)d_{13} + 28d_{14} + 2(33c_2 - 4c_3 + 5c_4 + 16c_5)d_{15} + 2(22c_2 + 12c_3 - 18c_4 - \\
 &19c_5)d_{16} - 102d_{19} + (4c_2 - 10c_3 - 6c_4 + 7c_5)d_{20} + (154c_2 - 58c_3 + 121c_4 + \\
 &199c_5)d_{23} + 3(22c_2 + 18c_3 - 3c_4 - 22c_5)d_{24} + 132d_6 + (10c_2 - 5c_3 + 5c_4 + \\
 &4c_5)d_7 - 16d_8 + 2(6c_2 - 3c_3 + 3c_4 + 13c_5)d_9) + 2c^2(42d_{17} + 118d_{18} - 8d_{21} + \\
 &17d_{22} + 54d_{25} - 37d_{26} + 7d_{27} + 193d_{28} + 160d_{29}) ; \\
 5 &= \frac{1}{48} 48d_1 + 2(3c_3 - 2c_4 + 4c_5)d_2 + 2c_4c_5d_3 + 2(c_3 - 6c_4 + 4c_5)d_4 + \\
 &2(c_4^2 - c_5^2)d_5 + c(8d_{10} + 4(c_2 + 4c_5)d_{11} + (10c_2 - 7c_3 + 11c_4 + 8c_5)d_{12} + \\
 &(6c_2 - 3c_3 + 3c_4 + 4c_5)d_{13} + 24d_{14} - 4(3c_2 - c_3 + c_4 + c_5)d_{15} + 4(4c_2 - \\
 &2c_3 + 3c_4 + 3c_5)d_{16} + 16d_{19} + 4c_5d_{20} + 2(28c_2 - 15c_3 + 6c_4 + 42c_5)d_{23} + \\
 &4(4c_2 + c_3 + 5c_4 - 17c_5)d_{24} + 8d_6 + 32d_8 + (2c_2 + c_3 - c_4 - 4c_5)d_9) + \\
 &4c^2(2d_{17} + 4d_{18} - 6d_{22} + 3d_{25} + d_{26} - 4d_{27} - 13d_{28} - 4d_{29}) ; \\
 6 &= \frac{1}{48} 12d_1 + 2c_5^2d_3 + 2(c_2 - c_4)d_4 - c_4c_5d_5 + c(20d_{10} + 2(2c_2 - \\
 &c_3 + 4c_5)d_{12} - 4d_{14} + 2(c_2 + 2c_5)d_{15} - 2(c_4 + c_5)d_{16} + 8d_{19} + (8c_2 - \\
 &2c_3 + 7c_4 + 8c_5)d_{23} + 2(10c_2 - 5c_3 + 11c_4 + 5c_5)d_{24} + 4d_6 + 4d_8) + \\
 &2c^2(2d_{17} + 12d_{18} - 2d_{22} + d_{25} + 3d_{26} - 4d_{27} + 9d_{28} + 12d_{29}) ; \\
 7 &= \frac{1}{24} 4d_{14} + 2c_4d_{23} - c_5d_{24} + 2c(d_{25} - d_{26} + 9d_{28} - 2d_{29}) ; \\
 8 &= \frac{1}{24} c_4d_{11} - c_5d_{13} + 8d_{14} + 2c_2d_{15} + (6c_2 + 7c_4)d_{23} - 2c_5d_{24} + \\
 &8d_6 + 2c(-2d_{17} + 8d_{18} + 12d_{25} - 2d_{26} - d_{27} + 27d_{28} - 4d_{29}) ; \\
 9 &= \frac{1}{24} c_4d_{12} + 4d_{14} + 2c_5d_{15} - 2c_4d_{16} + (3c_3 + 4c_4)d_{23} - c_3d_{24} + 4d_8 + \\
 &2c(4d_{18} - 2d_{22} + d_{25} + 9d_{26} - d_{27} + 18d_{28}) ; \\
 10 &= \frac{1}{24} 2c_4d_{11} - 2c_5d_{13} + 8d_{14} + 3c_4d_{27} - c_5d_{16} + 2c_2d_{23} + c_4d_{23} + 6d_6 + \\
 &2c(11d_{18} - d_{22} + 2d_{25} - 2d_{27} + 6d_{28}) ;
 \end{aligned} \tag{72}$$

$$\begin{aligned}
11 &= \frac{1}{24} (2c_5 d_{11} - 2c_4 d_{13} + 4d_{14} + c_5 d_{15} + c_4 d_{16} + 12d_{19} + c_4 d_{20} + (4c_3 + c_4 + \\
&6c_5) d_{23} + (c_4 + 2c_5) d_{24} + 2d_8 + 4c(d_{18} + 4d_{22} + d_{25} - d_{26} + 13d_{28} - 2d_{29}) ; \\
12 &= \frac{1}{24} (c_5 d_{11} + c_4 d_{13} + 6d_{19} + c_3 d_{23} + 10c_5 d_{23} + c_3 d_{24} - 4c_4 d_{24} + 2d_8 + \\
&2c(d_{25} + d_{26} + 7d_{27} + 8d_{28} + 8d_{29}) ; \\
13 &= \frac{1}{24} (4d_{10} + c_5 d_{12} + 2d_{19} + 2c_2 d_{24} + 2c_4 d_{24} + 2c(d_{25} + d_{26} + d_{27} + 8d_{29}) ; \\
14 &= \frac{1}{48} (4(3c_2 + c_4) d_{11} + 8d_{14} + 2(2c_2 - c_4) d_{15} - 2c_5 d_{16} + (12c_2 + 7c_4) d_{23} + \\
&2c_5 d_{24} + 24d_6 - 6c_5 d_7 + 2c(-4d_{17} + 20d_{18} - 6d_{21} - 2d_{22} + 37d_{25} - 2d_{26} + \\
&37d_{28} + 4d_{29}) ; \\
15 &= \frac{1}{24} ((5c_3 + c_4) d_{11} + (2c_2 + c_4) d_{12} + (-3c_3 + 2c_5) d_{13} + 8d_{14} + (c_3 + \\
&4c_5) d_{15} + (c_3 - 4c_4) d_{16} + (8c_2 + 11c_3 + 6c_4 - 4c_5) d_{23} + 4(c_3 + c_4 \\
&2c_5) d_{24} + 8d_6 - 4c_4 d_7 + 20d_8 + c_4 d_9 + 4c(4d_{17} + 8d_{18} + 11d_{25} + 5d_{26} - 7d_{27} + \\
&34d_{28} - 8d_{29}) ; \\
16 &= \frac{1}{48} (12d_{10} - 2c_5 d_{11} + 2(2c_3 + c_4 + 3c_5) d_{12} - 2(2c_2 + c_4) d_{13} + \\
&4d_{14} + 2c_2 d_{16} + 6d_{19} - 3c_5 d_{20} - (2c_3 + 7c_5) d_{23} + (4c_2 - 2c_3 + c_4 + \\
&4c_5) d_{24} + 8d_8 + 2c_5 d_9 + 2c(2d_{17} + 6d_{18} + 2d_{21} + d_{22} - 3d_{25} + 26d_{26} - 11d_{27} \\
&14d_{28} + 4d_{29}) ; \\
17 &= \frac{1}{24} ((c_2 + 3c_3) d_{11} - 2c_3 d_{13} + 2(c_2 + c_3 - c_5) d_{23} + 2c_4 d_{24} + 4d_6 + \\
&(c_3 - 2c_4) d_7 + 6d_8 + c_2 d_9 + 4c(d_{17} + 2d_{21} + 7d_{25} - 4d_{27} + 6d_{28}) ; \\
18 &= \frac{1}{24} (6d_{10} - 2c_5 d_{11} + (c_2 + 3c_3 + 6c_5) d_{12} - (3c_2 - c_3 + 2c_4) d_{13} + \\
&6d_{19} - 3c_5 d_{20} - (2c_3 + 7c_5) d_{23} + (2c_2 - 6c_3 + c_4) d_{24} + 2d_6 + c_2 d_7 + \\
&8d_8 + c_3 d_9 + 2c_5 d_9 + 2c(4d_{17} + 6d_{18} + 2d_{21} - 3d_{22} - 3d_{25} + 20d_{26} - 5d_{27} \\
&14d_{28} - 20d_{29}) ; \\
19 &= \frac{1}{24} (2d_{10} + 2d_{19} + c_5 d_{20} + (2c_2 + c_4) d_{24} + 2c(d_{18} + d_{22} + 6d_{29}) ; \\
20 &= \frac{1}{24} (24d_{10} + (c_3 + 2c_5) d_{12} + 2c_2 d_{13} + c_4 d_{13} + 12d_{19} + (c_3 + \\
&3c_5) d_{20} - 3c_5 d_{23} - (14c_2 - 3c_3 + 7c_4) d_{24} + 4d_8 + c_5 d_9 + 2c(2d_{17} \\
&6d_{18} + 2d_{21} + 7d_{22} - 3d_{25} + 6d_{26} + 9d_{27} - 36d_{29}) ; \\
21 &= \frac{1}{24} ((c_3 + 6c_5) d_{11} + (c_3 - 3c_4) d_{13} + 18d_{19} + c_2 d_{20} + (c_2 + 6c_3 + 10c_5) d_{23} \\
&4(c_3 + c_4) d_{24} + 4d_6 + c_4 d_7 + 4d_8 + 4c(4d_{21} + d_{22} + 5d_{25} - 2d_{26} + 3d_{27} + 20d_{28} \\
&16d_{29}) ; \\
22 &= \frac{d_{28}}{6} ; \quad 23 = \frac{d_{28}}{12} ; \quad 24 = \frac{1}{12} (2d_{26} + d_{28}) ; \quad 25 = \frac{1}{12} (d_{27} + 2d_{29}) ; \\
26 &= \frac{1}{12} (2d_{17} + d_{25} + 3d_{26} + d_{28}) ; \quad 27 = \frac{1}{12} (4d_{18} + 3d_{25} + 2d_{28}) ; \\
28 &= \frac{1}{12} (2d_{22} + d_{26} + d_{28}) ; \quad 29 = \frac{1}{12} (d_{21} + d_{27} + 3d_{29}) ; \\
30 &= \frac{1}{12} (d_{22} + d_{27} + 4d_{29}) ; \quad 31 = \frac{1}{12} (3d_{18} + 2d_{25}) ; \\
32 &= \frac{1}{12} (5d_{17} + 2d_{18} + d_{25} + 2d_{26}) ; \quad 33 = \frac{1}{12} (d_{17} + 4d_{21} + 3d_{22} + d_{25} + d_{26}) ; \\
34 &= \frac{1}{12} (d_{22} + d_{26} + 2d_{27}) ; \quad 35 = \frac{1}{12} (d_{17} + d_{18} + 3d_{21} + 2d_{22}) ; \\
36 &= \frac{1}{12} (2d_{25} + 3d_{28}) ; \quad 37 = \frac{1}{12} (d_{17} + 2d_{21} + d_{22} + 5d_{27}) ; \\
38 &= \frac{1}{24} (2d_{19} + c_5 d_{23} + c_4 d_{24} + 4c(d_{28} + 3d_{29})) ;
\end{aligned}$$

$$\begin{aligned}
1 &= \frac{d_{23}}{6}; \quad 2 = \frac{1}{12}(2d_{11} + 3d_{23}); \quad 3 = \frac{1}{12}(2d_{12} + d_{23}); \\
4 &= \frac{1}{12}(3d_{11} + 2d_{15} + 2d_{23}); \quad 5 = \frac{1}{12}(d_{11} + 3d_{12} + d_{23} + 2d_9); \\
6 &= \frac{1}{12}(d_{12} + 2d_{20} + d_{23}); \quad 7 = \frac{1}{12}(2d_{11} + 3d_{15}); \\
8 &= \frac{1}{12}(d_{11} + 2d_{12} + d_{15} + 3d_9); \quad 9 = \frac{1}{12}(d_{11} + d_{12} + 3d_{20} + d_9); \\
10 &= \frac{1}{12}(d_{12} + d_{20}); \quad 11 = \frac{d_{15}}{6}; \quad 12 = \frac{1}{12}(d_{15} + 2d_9); \\
13 &= \frac{1}{12}(d_{15} + 2d_{20} + d_9); \quad 14 = \frac{1}{12}(d_{20} + d_9); \quad 15 = \frac{d_{20}}{12}; \\
16 &= \frac{d_{24}}{12}; \quad 17 = \frac{1}{12}(d_{13} + 2d_{24}); \quad 18 = \frac{1}{12}(d_{13} + d_{16} + 4d_{24}); \\
19 &= \frac{1}{12}(d_{13} + 3d_{24} + d_7); \quad 20 = \frac{1}{12}(2d_{13} + d_{16}); \quad 21 = \frac{1}{12}(5d_{13} + d_{16} + 2d_7); \\
22 &= \frac{d_{16}}{6}; \quad 23 = \frac{1}{12}(3d_{16} + 4d_7); \quad 24 = \frac{1}{12}(2d_{16} + 3d_7); \\
25 &= \frac{1}{12}(2d_2 + c(d_{11} - d_{12} + 9d_{23} - 2d_{24})); \\
26 &= \frac{1}{12}(6d_2 + 2c_4d_3 - c_5d_5 + c(11d_{11} - d_{12} - d_{13} + 2d_{15} - 2d_{20} + 10d_{23} \\
&\quad 8d_{24})); \\
27 &= \frac{1}{12}(2d_2 + 2c_5d_3 - c_4d_5 + c(d_{11} + 7d_{12} + d_{13} + 2d_{15} - 2d_{16} + 8d_{23})); \\
28 &= \frac{1}{24}(8d_2 + 6c_4d_3 - c_5d_5 + 2c(2d_{11} - 2d_{13} + 11d_{15} - d_{16} - d_{20} + 2d_{23})); \\
29 &= \frac{1}{24}(6d_2 + 2(c_2 + c_3 - 2c_5)d_3 - (c_3 - c_4)d_5 + 2c(4d_{11} - 2d_{13} + 4d_{15} \\
&\quad 2d_{20} + 2d_{23} + d_7 + d_9)); \\
30 &= \frac{1}{24}(8d_2 + 2(c_3 + 4c_5)d_3 + (c_3 - 4c_4)d_5 + 4c(d_{11} + d_{12} + d_{13} + 4d_{15} \\
&\quad 2d_{16} + 2d_{20} + 6d_{23} - 4d_{24} + 4d_9)); \\
31 &= \frac{1}{24}(2d_2 + c_2d_5 + 2c(2d_{12} + 2d_{20} + 2d_{24} + d_7 + d_9)); \\
32 &= \frac{1}{24}(4d_2 + 2c_5d_3 + c_4d_5 + 2c(d_{15} + d_{16} + 7d_{20} + 2d_{23} + 2d_{24})); \\
33 &= \frac{1}{2}(d_4 + 2c(d_{23} + 3d_{24})); \\
34 &= \frac{1}{12}(3d_4 + c(d_{11} + d_{12} + 7d_{13} + 4d_{23} + 4d_{24})); \\
35 &= \frac{1}{12}(d_4 + c(d_{11} + d_{12} + d_{13} + 4d_{24})); \\
36 &= \frac{1}{24}(12d_4 + c_4d_5 + 2c(2d_{11} - 2d_{13} + 3d_{15} + 7d_{16} - d_{20} + 6d_{23} - 2d_{24})); \\
37 &= \frac{1}{24}(18d_4 + c_2d_5 + 4c(3d_{11} - 2d_{12} + d_{13} + d_{16} + 7d_{23} - 8d_{24} + 4d_7)); \\
38 &= \frac{1}{24}(18d_4 + c_3d_5 + 4c(d_7 + d_9 - 3d_{11} + 4d_{12} + d_{13} + 2d_{16} - 3d_{23} - 10d_{24})); \\
39 &= \frac{1}{24}(2d_4 + c_5d_5 + 2c(d_{15} + d_{16} + d_{20} + 2d_{24})); \\
40 &= \frac{c}{144}(80d_2 + 2(6c_2 - 3c_3 + c_4 + 12c_5)d_3 + 16d_4 + (10c_2 - 5c_3 + 5c_4 + \\
&\quad 8c_5)d_5 + 8c(3d_{11} - d_{12} - 2d_{13} + 3d_{15} - d_{16} - 2d_{20} + 2d_{23} - 2d_{24})); \\
41 &= \frac{c}{48}(8d_2 + 24d_4 + (2c_2 - c_3 + c_4 + 4c_5)d_5 + 8c(d_{11} - d_{12} + d_{23} - 2d_{24})); \\
1 &= \frac{d_3}{3}; \quad 2 = \frac{d_3}{4}; \quad 3 = \frac{d_3}{6}; \quad 4 = \frac{d_3}{12}; \quad 5 = 10 = \frac{d_5}{6}; \quad 6 = \frac{d_5}{12}; \quad 7 = \frac{d_5}{4}; \\
8 &= 9 = \frac{d_5}{12}; \quad 11 = \frac{c}{12}(7d_3 + d_5); \quad 12 = \frac{c}{6}(d_3 + 4d_5); \quad 13 = \frac{c}{12}(d_3 + d_5);
\end{aligned}$$

i the following three sets of constraints are satisfied:

$$\begin{aligned}
d_{15} - d_{16} &= d_5 - 2d_3 = c_4d_3 - 2cd_{23} = -d_{12} + d_{13} - d_{15} + d_{20} + d_{23} - 2d_{24} = \\
&\quad d_9 + d_{11} + d_{13} - 2d_7 = (2c_2 - c_3 + c_4)d_3 - 2(d_{11} - d_{13} + d_{16} - d_{20} + 2d_{24})c = \quad (75) \\
&\quad (c_2 - c_3 + c_4 + c_5)d_3 + 2(d_{13} - d_{16} - d_{24} - d_7)c = 0;
\end{aligned}$$

$$(2c_2 - c_3 + c_4 + 2c_5)d_3 - 2d_4 + 4d_{24}c = -4d_2 + (6c_2 - 3c_3 + 3c_4 + 8c_5)d_3 + 8d_{24}c = 0; \quad (76)$$

$$\begin{aligned}
& 4d_{10} + 2c_5d_{13} - 2c_5d_{16} + 2d_{19} + c_5d_{20} + c_5d_{23} + (2c_2 - c_4 - 4c_5)d_{24} + \\
& 2c(2d_{18} - d_{22} - d_{25} + 2d_{26} - d_{27} - 2d_{28} + 4d_{29}) = 0; \\
& 4d_{14} - 2c_4d_{11} + 2(c_2 - c_4 + c_5)d_{16} + (4c_2 + 3c_4)d_{23} + 2c_5d_{24} + 2c(-2d_{17} + \\
& 4d_{18} + 2d_{22} + d_{25} - d_{26} + 13d_{28} + 4d_{29}) = 0; \\
& 2d_8 - (c_4 - 2c_5)d_7 - 2(c_4 - c_5)d_{11} + (2c_2 + c_3 + 2c_4 + c_5)d_{13} - (c_3 + 3c_4)d_{16} + \\
& 6d_{19} + (c_3 + 2c_4 + c_5)d_{20} + (5c_2 - c_3 + 3c_4)d_{23} - (4c_2 + c_3 + 6c_4)d_{24} + \\
& 2c(2d_{17} - 2d_{18} - 4d_{21} - 7d_{22} - d_{25} + 2d_{26} + 11d_{27} + 10d_{28} - 20d_{29}) = 0; \\
& 4d_6 + (2c_2 + 2c_3 - c_4 - 4c_5)d_7 - 12d_{10} + (c_3 + 2c_5)d_{11} - (c_2 - 4c_4 + 9c_5)d_{13} \\
& 4d_{14} + (3c_2 - 5c_5)d_{16} + 12d_{19} - (2c_2 + c_3 + 2c_4 - 4c_5)d_{20} + (c_2 + 4c_3 + c_4 \\
& c_5)d_{23} + (6c_2 - 2c_3 + 3c_4 - 12c_5)d_{24} + (-8d_{17} + 20d_{18} + 4d_{21} - 14d_{22} + 10d_{25} + \\
& 4d_{26} + 10d_{27} + 44d_{28} - 16d_{29}) = 0; \\
& 24d_1 + 6(c_5^2 - c_4c_5)d_3 + 2(6c_2 - c_4)d_4 + (2(2c_2 + 2c_3 - c_4 + 4c_5)d_7 - 4d_8 \\
& 52d_{10} + 2(-4c_2 + c_3 - c_4 + 2c_5)d_{11} + 2(c_2 + c_3 - 6c_4 + 6c_5)d_{13} - 16d_{14} + 2(5c_2 \\
& c_3 + c_4 - c_5)d_{16} + 62d_{19} + (-4c_2 + 2c_3 - 12c_4 + 5c_5)d_{20} + (2c_2 + 12c_3 + 4c_4 + \\
& 13c_5)d_{23} + (6c_2 - 12c_3 + 5c_4)d_{24})c + 2c^2(-16d_{17} + 50d_{18} + 4d_{21} - 13d_{22} + d_{25} + \\
& 20d_{26} + 3d_{27} + 52d_{28} + 4d_{29}) = 0;
\end{aligned} \tag{77}$$

The first set of constraints (75) is automatically satisfied by the parametrizations (33) and (68), while the second set of two constraints (76) is satisfied by the parametrizations (33) and (68) and by the  $O(\epsilon^2)$  constraint (45). The remaining five constraints (77) are equivalent to the five quadratic constraints (53), (54)–(58) in the S-integrability scenario in which (48) holds, and to the linear constraints (11) in the C-integrability scenario in which  $c = 0$  (the last constraint is automatically satisfied by the condition  $c = 0$  and, in the remaining constraints, the quadrics degenerate into the hyperplanes described by equations (11)).

Acknowledgements. We acknowledge interesting discussions with U. Aglietti.

## References

- [1] A. Degasperis, S. V. Manakov and P. M. Santini, "Multi-scale perturbation beyond the nonlinear Schrödinger equation. I", *Physica D* 100, 187–211 (1997).
- [2] A. Degasperis and M. Procesi, "Asymptotic integrability" in *Symmetry and perturbation theory*, SPT 98 23–37, edited by A. Degasperis and G. Gaeta, World Scientific, Singapore (1999).

- [3] A. Degasperis, "Multiscale expansion and integrability of dispersive wave equations", lectures given at the Euro Summer School "What is integrability?", Isaac Newton Institute, Cambridge, U.K., 13-24 August (2001); in *Integrability* edited by A. Mikhailov, Lecture Notes in Physics 767, Springer, Berlin-Heidelberg (2009).
- [4] P. Winternitz, "Symmetries of discrete systems. Discrete integrable systems", 185{243, Lecture Notes in Phys. 644, eds. B. Grammaticos, Y. Kosm ann-Schwarzbach and T. Tam izhm ani, Springer, Berlin, 2004.
- [5] F. Valiquette and P. Winternitz, "Discretization of partial differential equations preserving their physical symmetries", arXiv:math-ph/0507061.
- [6] T. R. Taha and M. J. Ablowitz, "Analytical and numerical aspects of certain nonlinear evolution equations II. Numerical, nonlinear Schrödinger equation", *J. Comput. Phys.* 55, 192 (1984).
- [7] B. M. Herbst and M. J. Ablowitz, "Numerically induced chaos in the nonlinear Schrödinger equation", *Phys. Rev. Lett.* 62, 2065-2068 (1989).
- [8] P. L. Kelley, *Phys. Rev. Lett.* (1965) 15 1005.
- [9] V. E. Zakharov, *Soviet Phys. JETP* (1968) 994-998.
- [10] D. J. Benney and A. C. Newell, *J. Math. and Phys. (now Stud. Appl. Math.)* 46 (1967) 133-139.
- [11] A. Hasegawa and T. Tappert, *Appl. Phys. Lett.* 23 (1972) 142.
- [12] H. Hasimoto and H. Ono, *J. Phys. Soc. Japan* 33 (1972) 805.
- [13] T. Taniuti, *Suppl. Proc. Th. Phys.* 55, 1 (1974).
- [14] Y. Kodama and T. Taniuti, *J. Phys. Soc. Japan*, 45, 298 (1978).
- [15] F. Calogero and W. Eckhaus, "Nonlinear evolution equations, rescalings, model PDEs and their integrability: I" *Inverse Problems* 3, 229-262 (1987).
- [16] V. E. Zakharov and A. S. Shabat, *Soviet Phys. JETP* 34 (1972) 62.

- [17] V.E. Zakharov and S.V. Manakov, "Resonance interaction of wave packets", *Soviet Physics JETP* 42 (1975).
- [18] V.E. Zakharov and E.A. Kuznetsov, *Physica D* 18, 455 (1986).
- [19] D.J. Korteweg and F. de Vries, "On the Change of Form of Long Waves Advancing in a Rectangular Canal, and on a New Type of Long Stationary Waves." *Philos. Mag.* 39, 422-443, 1895.
- [20] C.S. Gardner, C.S. Greene, M.D. Kruskal, and R.M. Miura, "Method for Solving the Korteweg-de Vries Equation." *Phys. Rev. Lett.* 19, 1095-1097, 1967.
- [21] F. Calogero, "Why are certain nonlinear PDEs both widely applicable and integrable?" in *What is integrability?* 1-62, edited by V.E. Zakharov, Springer, Berlin-Heidelberg (1991).
- [22] E. Hopf, *Commun. Pure Appl. Math.* 3, 201 (1950). J.D. Cole, *Quant. Appl. Math.* 9, 225 (1951).
- [23] V.E. Zakharov, S.V. Manakov, S.P. Novikov and L.P. Pitaevsky, *Theory of Solitons. The Inverse Problem Method*, Plenum Press (1984).
- [24] F. Calogero and A. Degasperis, "Spectral Transform and solitons: tools to solve and investigate nonlinear evolution equations. Volume one", North Holland, Amsterdam (1982).
- [25] M.J. Ablowitz and P.C. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University Press, Cambridge, 1991.
- [26] B. Konopelchenko, *Solitons in Multidimensions*, World Scientific, Singapore (1993).
- [27] F. Calogero and W. Eckhaus, "Nonlinear evolution equations, rescalings, model PDEs and their integrability: II", *Inverse Problems* 4, 11-13 (1988).
- [28] F. Calogero, A. Degasperis and X.D. Ji, "Nonlinear Schrodinger-type equations from multiscale reduction of PDEs. I. Systematic derivation", *J. Math. Phys.* 41, 6399-6443 (2000).

- [29] F. Calogero, A. Degasperis and X.D. Ji, "Nonlinear Schrodinger-type equations from multiscale reduction of PDEs. II. Necessary conditions of integrability for real PDEs", *J. Math. Phys.* 42, 2635-2652 (2001).
- [30] Y. Kodama and A.V. Mikhailov, "Obstacles to asymptotic integrability", in *Algebraic aspects of integrable systems: in memory of Irene Dorfmán*, 173-204, edited by A.S. Fokas and I.M. Gelfand, Birkhauser, Boston (1996).
- [31] D. Levi, M. Petrera and C. Scimitema, "On the integrability of the discrete nonlinear Schrodinger equation", *E.P.L.* 84, 10003 (2008).
- [32] C. Scimitema, "Multiscale techniques for nonlinear difference equations", PhD Thesis, Dept. of Physics, University of Roma3, Roma, Italy (2009).
- [33] C. Scimitema, "Multiscale reduction of discrete Korteweg-de Vries equations", *J. Phys. A: Math. Theor.* (special issue for the Conference SDE 8).
- [34] R.I. Yamilov, "Symmetries and integrability criteria for differential difference equations". *J. Phys. A: Math. Gen.*, R541-R623 (2006).
- [35] V.V. Sokolov and A.B. Shabat, "Classification of integrable evolution equations", *Sov. Sci. Rev. section C* 4, 221-80 (1984).
- [36] A.V. Mikhailov, A.B. Shabat and R.I. Yamilov, "The symmetry approach to the classification of nonlinear equations. Complete lists of integrable systems", *Russian Math. Surveys* 42/4, 1-63 (1987).
- [37] D. Pelinovski, "Translationally invariant nonlinear Schrodinger lattices", *Nonlinearity* 19, 2695-2716 (2006).
- [38] M.J. Ablowitz and J.F. Ladik, *J. Math. Phys.* 17, 1011 (1979).
- [39] A.S. Davydov, *J. Theor. Biol.* 38 559 (1973).
- [40] W.P. Su, J.R. Schrieffer and A.J. Heeger, *Phys. Rev. Lett.* 42 698 (1979).
- [41] J.C. Eilbeck, P.S. Lomdhal, A.C. Scott, *Physica D* 16, 318 (1985).

- [42] D . Hennig and G . T sironis, *Physics Reports* 307, 333 (1999).
- [43] F . K . Abdullaev, B . B . Baizakov, S . A . D arm anyan, V . V . K onotop and M . Salerno, *Phys. Rev. A* 64, 043606 (2001).
- [44] M . O ster, M . Johansson and A . Eriksson, \Enhanced mobility of strongly localized modes in waveguide arrays by inversion of stability", *Phys. Rev. E* 67, 056606 (2003).
- [45] C . Claude, Y . S . K ishar, O . K luth and K . H . Spatschek, \Moving localized modes in nonlinear lattices", *Phys. Rev. B* 47, 14228-14232 (1993).
- [46] F . M agri, *J. Math. Phys.* 19, 1156 (1978).
- [47] I . G elfand and I . D orfm an, *Funct. Anal. Appl.* 13 (1979); 14 (1980).
- [48] A . S . Fokas and B . Fuchssteiner, *Lett. Nuovo C im ento* 28 299 (1980); *Physica* 4D 47 (1981).
- [49] P . M . Santini and A . S . Fokas: \Recursion operators and biham iltonian structures in m ultidim ensions.I"; *Com m M ath Phys.* 115, 375-419 (1988).
- [50] M . J . Ablowitz, D . K aup, A . C . Newell and H . Segur, *Stud. Appl. Math.* 53, 249 (1974).
- [51] U . A glietti and P . M . Santini, \Multiscale expansions of difference equations in the sm all lattice spacing regime. Integrability test and num erical con m ations", preprint (in preparation).
- [52] U . A glietti, P . M . Santini and C . Scim ita, preprint (in preparation).