

4C 02.27: a quasar with episodic activity?

M . Jam rozy^{1?}, D .J. Saikia^{2;3} and C .K onar⁴

¹ Obserwatorium Astronomiczne, Uniwersytet Jagielloński, ul. Orła 171, PL-30244 Kraków, Poland

² National Centre for Radio Astrophysics, TIFR, Ganeshkhind, Post Bag 3, Pune 411 007, India

³ Australia Telescope National Facility, CSIRO, PO Box 76, Epping, NSW 1710, Australia

⁴ Inter-University Centre for Astronomy and Astrophysics, Ganeshkhind, Post Bag 4, Pune 411 007, India

Accepted. Received

ABSTRACT

Striking examples of episodic activity in active galactic nuclei are the double-double radio galaxies (DDRGs) with two pairs of oppositely-directed radio lobes from two different cycles of activity. Although there are over about a dozen good examples of DDRGs, so far no case of one associated with a quasar has been reported. We present Giant Metrewave Radio Telescope observations of a candidate double-double radio quasar (DDRQ), J0935+0204 (4C 02.27), and suggest that radio jets in this source may also have been intrinsically asymmetric, contributing to the large observed asymmetries in the flux density and location of both pairs of radio lobes.

Key words: galaxies:active { galaxies:nuclei { galaxies:individual:4C 02.27 { radio continuum :galaxies

1 INTRODUCTION

A striking example of episodic nuclear activity is when a new pair of radio lobes is seen closer to the nucleus before the 'old' and more distant radio lobes have faded (e.g. Subrahmanyan, Saripalli & Hunstead 1996; Lara et al. 1999). Such sources have been christened as 'double-double' radio galaxies (DDRGs) by Schoenmakers et al. (2000a). More recently Brocksopp et al. (2007) have also reported the discovery of a triple-double radio galaxy with three distinct cycles of activity. Although more than approximately a dozen or so of such DDRGs are known in the literature (Saikia, Konar & Kulkarni 2006, and references therein), so far no case of one associated with a quasar has been reported. This could be partly due to difficulties in distinguishing a knot of emission in a jet from a hotspot, since quasars tend to have more prominent jets than galaxies due to effects of relativistic beaming. However, if radio galaxies and quasars are intrinsically similar, one should find evidence of emission from an earlier cycle of activity in quasars as well. This may be in the form of distinct pairs of lobes as in the DDRGs, or may be seen as diffuse relic emission beyond the extent of the younger double lobed radio source, as seen for example in the radio galaxy 4C 29.30 (Jam rozy et al. 2007).

Since the lobes from an earlier cycle of activity are likely to have a steep radio spectrum, low-frequency observations with telescopes such as the Giant Metrewave Radio

Telescope (GMRT) should help identify such features. However, although the DDRG J0041+3224 was discovered from GMRT and Very Large Array (VLA) observations (Saikia et al. 2006), deep GMRT observations of a few hundred sources at 153, 244, 610 and 1260 MHz have not yielded clear examples of sources with episodic activity (Sirothia et al. 2009). We have been systematically making observations, searching the literature as well as in ages from surveys made with the VLA such as NVSS (NRAO VLA Sky Survey; Condon et al. 1998) and FIRST (Faint Images of the Radio Sky at Twenty-cm; Becker, White & Helfand 1995) for candidate DDRGs and DDRQs or quasars with signs of episodic radio activity. Identifying a DDRG/DDRQ is not automated and each image is examined by eye. In this paper, we present observations with the GMRT at 619 MHz of possible relic emission associated with the quasar, 4C 02.27 (J0935+0204), and discuss the possibility that this might be a quasar with signs of episodic activity.

2 4C 02.27

The radio source 4C 02.27 (J0935+0204) is associated with a quasar at a redshift of 0.649117 ± 0.000210 as listed in the NASA Extragalactic Database (NED) from measurements made as part of the the SDSS (Sloan Digital Sky Survey; Abazajian et al. 2004; Schneider et al. 2005). The corresponding luminosity distance is 3880.7 Mpc and 1 arcsec corresponds to 6.918 kpc in a Universe with $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.27$, $\Omega_\Lambda = 0.73$ (Spergel et al. 2003).

? E-mail: jam.rozy@oa.uj.edu.pl (M J); djs@ncra.tifr.res.in (D JS); chiran.jb.konar@gmail.com (CK)




Figure 1. Left panel: VLA image of 4C 02.27 at 1413 MHz with an angular resolution of 1.7 arcsec reproduced from Hintzen et al. (1983), with the components NE_{inn}, SW_{inn} and SW_{out} marked. Right panel: GMRT image of 4C 02.27 at 619 MHz with an angular resolution of 5.42 × 4.68 arcsec² along a PA of 49° showing the outer north-eastern component, NE_{out}. The cross marks the position of the quasar.

The published radio images show a double-lobed radio source which is highly asymmetric in both flux density and location of the outer components, and has two prominent hotspots at the outer edges (labelled as SW_{out} and NE_{inn} in Fig. 1, left panel) with an overall angular separation of 47 arcsec (325 kpc). The separations of the peaks of emission in NE_{inn} and SW_{out} from the nucleus are 6.3 arcsec (43.6 kpc) and 40.6 arcsec (281 kpc) respectively (e.g. Hintzen, Uvestad & Owen 1983; Swarup, Sinha & Hildner 1984; Price et al. 1993). The corresponding arm-length ratio is 6, making it one of the very asymmetric sources in terms of the location of the components. The flux density ratio of the two components at 1400 MHz is 3.5, with SW_{out}, which is farther from the nucleus being also brighter. In addition, there is a weak component (SW_{inn}) with a flux density of 9 mJy at 1400 MHz located 3.5 arcsec (24.2 kpc) south-west of the nucleus along the axis of the source. There is no evidence of a distinct jet-like structure as defined by Bridle & Perley (1984). From VLA C-array observations at 5 GHz, Saikia et al. (1984) find the south-western component to be 15.4 per cent polarised compared with 5.7 per cent for the north-eastern one and 1.1 per cent for the core component.

In an optical study of the host galaxies of intermediate-redshift quasars, Ronnback et al. (1996) report an arm-like structure resembling a tidal tail in 4C 02.27 and show that the luminosity profile of the host galaxy follows an $r^{1.4}$ law. Their effective radius is consistent with earlier measurements by Romaniushin & Hintzen (1989) although Ronnback et al. find the host galaxy magnitude to be 0.4 mag brighter.

In the next Section we present GMRT observations of the source at 619 MHz as well as the NVSS and FIRST images of the source. In Section 3, we discuss the possibility

that 4C 02.27 (J0935+0204) could be a DDRQ exhibiting signs of episodic activity.

3 OUTER NORTH-EASTERN LOBE

In this section we present the results of the GMRT observations as well as the NVSS and FIRST images. The GMRT and VLA images show the presence of a diffuse lobe beyond the north-eastern hotspot.

3.1 GMRT observations and results

The source was observed with the GMRT on 2007 September 01 at 619 MHz for approximately 330 minutes on source. The observations were made in the standard manner, with each observation of the target-source interspersed with observations of the phase calibrator, 0943-083. 3C 286 and 3C 147 were both observed for flux density and bandpass calibration. The flux densities are on the Baars et al. (1977) scale. The data collected were calibrated and reduced in the standard way using the NRAO AIPS software package. Several rounds of self calibration were done to improve the quality of the images. The GMRT image has an angular resolution of 5.42 × 4.68 arcsec² along a position angle (PA) of 49° and an rms noise of 0.22 mJy beam⁻¹.

The GMRT image (Figure 1, right panel) shows clearly a diffuse lobe of emission to the north-east well beyond the eastern hotspot, which we will refer to as NE_{out}. The total flux density of this feature is 41 mJy, while the total flux density of the source is 1578 mJy. The total flux density of the south-western lobe (SW_{out}), is 1138 mJy, while that of the




Figure 2. 1400-MHz VLA images of 4C 02.27. NVSS and FIRST contour maps of the entire source overlaid on the optical field from the Digital Sky Survey (DSS). The contours are spaced by factors of 2 and the first contours are 1.35 and 0.6 mJy beam⁻¹ respectively. The sizes of the beams are indicated by ellipses in the bottom left corner of the image. The cross marks the position of the radio core.

central feature which contains the core and the inner north-eastern (NE_{inn}) and south-western (SW_{inn}) components is 393 mJy.

3.2 NVSS and FIRST images

In Figure 2, we present the NVSS and FIRST images of the source at 1400 MHz with angular resolutions of 45 and 6 arcsec respectively superimposed on the DSS image. The FIRST image is very similar to that of the GMRT one showing the diffuse outer lobe of emission to the north east (NE_{out}) in addition to the other features. The flux density of NE_{out} and SW_{out} are 15 mJy and 488 mJy, while that of the central feature consisting of the core and NE_{inn} and SW_{inn} is 230 mJy.

The spectral indices, (S/ν) , of the outer features, NE_{out} and SW_{out}, are 1.2 and 1.0 respectively between 610 and 1400 MHz, while the total flux densities yield a spectral index of 0.9. The two-point spectral index of the inner component consisting of the core, NE_{inn} and SW_{inn} is 0.7. The deconvolved size of the NVSS image, which is 52 × 12 arcsec² along a position angle of 52°, is consistent with the structure of the source. The total flux density of 787 mJy in the NVSS image is very similar to value of 800 mJy at 1413 MHz estimated by Hintzen et al. (1983). We have also examined the VLA Low-Frequency Sky Survey (VLSS; Cohen et al. 2007) image but this shows no new feature.

4 EPISODIC NATURE OF 4C 02.27

The GMRT and the VLA FIRST images clearly show two pairs of components, the inner and the outer doubles with angular sizes of 10 and 68 arcsec respectively, which corresponds to projected linear sizes of 70 and 470 kpc respectively. The inner and outer lobes appear distinct and do not appear to be multiple hotspots in the lobes (cf. Lonsdale & Barthel 1986). The eastern component of the inner double clearly shows an edge-brightened structure similar to the hotspots of classical FR II sources. On the other hand, the western component of the inner double is compact and weak and its detailed structure is not well determined although the high-resolution image published by Price et al. (1993) hints that it too might have an edge-brightened structure. The flux density ratio of these two inner components is 1.7 at 1400 MHz while the peak brightness ratio is 6 (Hintzen et al. 1983). The inner double is reminiscent of the inner structure of the radio galaxy 3C 219 which has been interpreted to be due a restarted jet activity (Bridle, Perley & Henriksen 1986; Clarke et al. 1992), and has been included in samples of double-double radio galaxies (Schoenmakers et al. 2000a; Saikia et al. 2006; Saripalli & Mack 2007). Let us examine whether the asymmetry in location and brightness of the components of the inner double in 4C 02.27 can be understood in terms of the relativistic beaming scenario. Here the arm-length ratio, R is given by $(1 + \cos \theta) = (1 - \cos \theta)$ and the brightness ratio is $(R)^{2+\alpha}$, where $v = \beta c$ is the component velocity and θ is the angle of inclination of the jet axis to the line of sight. For a velocity of 0.4c and $\theta = 40^\circ$, which is a reasonable value for a lobe-dominated quasar, and $\alpha = 1$, $R = 1.9$ and the brightness ratio is 6.7. These values are close to the observed values of 1.8 and 6 estimated from the image of Hintzen et al. (1983), suggesting consistency of the arm-length and peak brightness ratio with the relativistic beaming scenario with the NE_{inn} lobe approaching us. High-resolution depolarisation and rotation measure estimates would help to examine this possibility. The precise values of the brightness ratio at different locations will also be affected by the microphysics of the dissipation of energy by shocks along the flow of the jet. However, the flux density ratio is much larger with the NE_{inn} component being stronger by 1.7, suggesting intrinsic asymmetries as well over the lifetime of the inner double.

The situation is even more complex for the outer double. From the GMRT 619-MHz image the arm-length ratio of the outer lobes is 1.3, while the peak brightness ratio is 60 and the total flux density ratio 30. The values estimated from the FIRST image are similar and are in the opposite sense to that of the inner double, with the SW_{out} lobe now being farther and also brighter. Considering the outer lobes alone, it would be difficult to reconcile the arm-length and the large brightness ratios in the simple version of the relativistic beaming scenario. Environmental asymmetries coupled with the effects of relativistic motion can produce a wide range of arm-length and brightness asymmetries (e.g. Jeyakumar et al. 2005). For a lobe approaching us and propagating through a dense medium, the oppositely directed lobes could be somewhat symmetrically located with large asymmetries in brightness or flux density ratios. In this case, the SW_{out} lobe would be approaching us, which would be inconsistent with the inner double. On the other hand,

if the NE_{out} lobe is approaching us, one would expect it to be brighter due to the effects of relativistic motion as well as a denser environment, inferred from its closer distance to the radio nucleus. These suggest that the radio jets may have been intrinsically asymmetric during this cycle of activity. The possibility of intrinsically asymmetric jets has been suggested earlier, for example, for the highly asymmetric double-lobed radio source B0500+630 (Saikia et al. 1996), weak-cored one-sided sources (Saikia et al. 1989, 1990), the one-sided radio emission in 3C273 which has been imaged with a dynamic range of $10^4:1$ (Davis, Muxlow & Conway 1985; Conway & Davis 1994), the inner jets of the radio galaxies M87 (Kovalev et al. 2007) and NGC 6251 (Jones 1986; Jones & Wehrle 2002), and the optical counter jet in the radio galaxy 3C66B (Fraix-Burnet 1997). On the theoretical side, Icke (1983) has suggested a hydrodynamical model for gas flows in the nuclear region that can affect the de Laval nozzle causing one-sided jets, while Wang, Sulkanen & Lovelace (1992) have explored models where the ratio of jet luminosities depends directly on the degree of asymmetry of the magnetic field in the nuclear region.

To explain the extreme asymmetries one might also consider the possibility that the orientation of the jet might change during the different cycles of activity; in the extreme cases the approaching jet during one cycle may be the receding one in the next or vice versa. However, the alignment of all the four components in 4C02.27 suggests that this is not a likely scenario for this source.

Another interesting aspect on the episodic nature of 4C02.27 is the presence of a hotspot in the SW_{out} lobe. This is not unique to 4C02.27. While the outer doubles are often diffuse as for example in J1453+3308 (Schoenmakers et al. 2000a; Konar et al. 2006), hotspots are also sometimes seen in the outer lobes as for example in the northern lobe of B1834+620 (Schoenmakers et al. 2000b). This can be used to estimate the time scale of interruption of energy supply. For typical sizes of hotspots of a few to 10 kpc in large radio sources (cf. Jeyakumar & Saikia 2000), the hotspots are expected to fade in about 10^4 – 10^5 yr (e.g. Kaiser, Schoenmakers & Rottgering 2000). This is a small fraction of the time it takes for the material in the jets of large radio sources to reach the hotspots from the radio core. Therefore it is reasonable to assume that the hotspot fades soon after the last jet material passes through them. The presence of a hotspot in the SW_{out} lobe implies that it still receives jet material. The travel time of the jet material from the core to the hotspot, t_j , is given by D_{hs}/v_{jet} , where D_{hs} is the physical distance of the hotspot from the core and v_{jet} is the velocity of the jet. In our case, t_j is 2.9×10^6 yr for an inclination angle, $i = 40^\circ$, and a jet velocity of $0.5c$. However, this estimate will be affected by light travel time effects due to the orientation of the source axis. For example, if the SW_{out} lobe is on the receding side, it will take longer for the information to reach us compared with it being on the approaching side. The observed time difference, t_{obs} , between the ejection of the last jet material and its arrival at the hotspot is 1.8 and 3.9 M yr depending on the orientation of the source. If the time scale of interruption of the jet is larger than t_{obs} , the hotspot in the SW_{out} lobe and the inner structure cannot be observed simultaneously. Therefore, the interruption of jet activity must be less than t_{obs} . Also, within this time period the inner double forms,

Table 1. Large radio quasars

Source	Alt. name	Red-shift (M Hz)	LAS (arcsec)	LLS (kpc)	Refs
(1)	(2)	(3)	(4)	(5)	(6)
J0439 2422		0.8400	128	979	1
J0631 5405		0.2036	312	1034	2
J0750+ 6541		0.7470	222	1627	3
J0810 6800		0.2311	390	1425	2
J1027 2312		0.3090	198	893	1
J1130 1320		0.6337	297	2033	4
J1353+ 2631		0.3100	190	859	5
J1427+ 2632		0.3660	240	1212	5
J1432+ 1548		1.0050	168	1353	6
J1504+ 6856		0.3180	204	939	3
J1723+ 3417	4C 34.47	0.2060	244	816	7,8
J2042+ 7508	4C 74.26	0.1040	610	1151	9

1: Ishwara-Chandra & Saikia (1999); 2: Saripalli et al. (2005); 3: Lara et al. (2001); 4: Bhatnagar, Gopal-Krishna & Wisotzki (1998); 5: Rogora, Padrielli & de Ruiter (1986); 6: Singal, Konar & Saikia (2004); 7: Jagers et al. (1982); 8: Hooimeyer et al. (1992); 9: Riley et al. (1988)

so that the time scale for interruption is less than t_{obs} . It is also of interest to note that the time scale of interruption for this lobe with a hotspot is much smaller than for say J1453+3308 which has diffuse outer lobes. The dynamical and spectral ages of the diffuse outer lobes of J1453+3308 are 215 and 50 M yr, while that of the inner double is only 2 M yr, suggesting a much longer time scale of interruption (Kaiser et al. 2000; Konar et al. 2006).

5 CONCLUDING REMARKS

The quasar 4C02.27 with an overall linear size of 470 kpc appears to exhibit signs of episodic activity and we classify it as a DDRQ. The source also exhibits evidence of an intrinsic asymmetry of the oppositely-directed jets. Although most DDRGs appear to be associated with large radio galaxies, often with sizes larger than approximately 1 Mpc, signs of episodic activity are seen in smaller sources as well (Schoenmakers et al. 2000a,b; Saikia et al. 2006 and references therein). Assuming an inclination angle of 40° to the line of sight, the intrinsic size of 4C02.27 would be 730 kpc, comparable to some of the known or candidate DDRGs (Jamrozy et al. 2009). Considering the tendency for DDRGs to often have large sizes, we examined the structures of large quasars with sizes larger than 800 kpc (Table 1), including the quasar type object J0750+6541 (Lara et al. 2001). These do not show evidence of episodic activity, which along with our search of the literature suggests that such objects are not common. A deep low-frequency search amongst 374 sources, most of which were small, did not show any clear example of a DDRG/DDRQ (Sirothia et al. 2009). Considering only giant radio galaxies, there are 4 DDRGs in the well-defined sample of 49 sources (Schoenmakers 1999), suggesting that 10 per cent of these large sources may show signs of episodic activity. Considering the known giant radio galaxies selected from different samples and sometimes with incomplete structural information (e.g. Ishwara-Chandra &

Saikia 1999; Lara et al. 2001; Machalski, Jamroz & Zola 2001; Saripalli et al. 2005), also suggests a similar percentage. Although the numbers are very small at present, a similar fraction would be consistent with the unified scheme for radio galaxies and quasars (e.g. Barthel 1989).

ACKNOWLEDGMENTS

We thank an anonymous reviewer and the editor for their comments which have significantly improved the paper, and the staff of GMRT for their help with the observations. MJ acknowledges the MNRAS funds for scientific research during the years 2009–2012 under contract No 3812/B/H03/2009/36. The GMRT is a national facility operated by the NCRA, TIFR. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under co-operative agreement by Associated Universities Inc. This research has made use of the NASA/IPAC extragalactic database (NED) which is operated by the Jet Propulsion Laboratory, Caltech, under contract with the National Aeronautics and Space Administration. We acknowledge use of the Digitized Sky Surveys which were produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The images of these surveys are based on photographic data obtained using the Oschin Schmidt Telescope on Palomar Mountain and the UK Schmidt Telescope.

REFERENCES

- Abazajian K., et al., 2004, *AJ*, 128, 502
 Baars J.W.M., Genzel R., Pauliny-Toth I.I.K., Witzel A., 1977, *A & A*, 61, 99
 Barthel P.D., 1989, *ApJ*, 336, 606
 Becker R.H., White R.L., Helfand D.J., 1995, *ApJ*, 450, 559
 Bhatnagar S., Gopal-Krishna, Wisotzki L., 2004, *MNRAS*, 299, 25L
 Bridle A.H., Perley R.A., 1984, *ARA & A*, 22, 319
 Bridle A.H., Perley R.A., Henriksen R.N., 1986, *AJ*, 92, 534
 Brockopp C., Kaiser C.R., Schoenmakers A.P., de Bruyn A.G., 2007, *MNRAS*, 382, 1019
 Clarke D.A., Bridle A.H., Burns J.O., Perley, R.A., Norman M.L., 1992, *ApJ*, 385, 173
 Cohen A.S., Lane W.M., Cotton W.D., Kassim N.E., Lazio T.J.W., Perley R.A., Condon J.J., Erickson, W.C., 2007, *AJ*, 134, 1245
 Condon J.J., Cotton W.D., Greisen E.W., Yin Q.F., Perley R.A., Taylor G.B., Broderick J.J., 1998, *AJ*, 115, 1693
 Conway R.G., Davis R.J., 1994, *A & A*, 284, 724
 Davis R.J., Muxlow T.W.B., Conway R.G., 1985, *Nature*, 318, 343
 Fraix-Burnet D., 1997, *MNRAS*, 284, 911
 Hintzen P., Ulvestad J., Owen F., 1983, *AJ*, 88, 709
 Hooley J.R.A., Barthel P.D., Schilizzi R.T., Miley G.K., 1992, *A & A*, 261, 1
 Icke V., 1983, *ApJ*, 265, 648
 Ishwara-Chandra C.H., Saikia D.J., 1999, *MNRAS*, 309, 100
 Jagers W.J., Miley G.K., van Bruggel W.J.M., Schilizzi R.T., Conway R.G., 1982, *A & A*, 105, 278
 Jamroz M., Konar C., Saikia D.J., Stawarz L., Mack K.-H., Siemiginowska A., 2007, *MNRAS*, 378, 581
 Jamroz M., Konar C., Saikia D.J., Machalski J., 2009, in Saikia D.J., Green D.A., Gupta Y., Venturi T., eds, *ASP Conf. Ser. Vol. 407, The Low-Frequency Radio Universe*, Astron. Soc. Pac., San Francisco, p. 137
 Jayakumar S., Saikia, D.J., 2000, *MNRAS*, 311, 397
 Jayakumar S., Wita P.J., Saikia, D.J., Hooda, J.S., 2005, *A & A*, 432, 823
 Jones D.L., 1986, *ApJ*, 309, L5
 Jones D.L., Wehrle A.E., 2002, *ApJ*, 580, 114
 Kaiser C.R., Schoenmakers A.P., Rottgering H.J.A., 2000, *MNRAS*, 315, 381
 Konar, C., Saikia, D.J., Jamroz, M., Machalski, J., 2006, *MNRAS*, 372, 693
 Kovalev Y.Y., Lister M.L., Homan D.C., Kellermann K.I., 2007, *ApJ*, 668, L27
 Lara L., Marquez I., Cotton W.D., Feretti L., Giovannini G., Marcaide J.M., Venturi T., 1999, *A & A*, 348, 699
 Lara L., Marquez I., Cotton W.D., Feretti L., Giovannini G., Marcaide J.M., Venturi T., 2001, *A & A*, 378, 826
 Lonsdale C.J., Barthel P.D., 1986, *AJ*, 92, 12
 Machalski J., Jamroz M., Zola S., 2001, *A & A*, 371, 445
 Price R., Gower A.C., Hutchings J.B., Talon S., Duncan D., Ross G., 1993, *ApJS*, 86, 365
 Riley J.M., Warner P.J., Rawlings S., Saunders R., Pooley G.G., Eales S.A., 1988, *MNRAS*, 236, 13P
 Rogora A., Padrielli L., de Ruiter H.R., 1986, *A & A*, 64, 557
 Romanihin W., Hintzen P., 1989, *ApJ*, 341, 41
 Ronback J., van Groningen E., Wanders I., Omdahl E., 1996, *MNRAS*, 283, 282
 Saikia D.J., Shastri P., Kapahi V.K., Sinha R.P., Swarup G., 1984, *JApA*, 5, 429
 Saikia D.J., Junor W., Muxlow T.W.B., Tzioumis A.K., 1989, *Nature*, 339, 286
 Saikia D.J., Junor W., Comwell T.J., Muxlow T.W.B., Shastri P., 1990, *MNRAS*, 245, 408
 Saikia D.J., Thomasson P., Jackson N., Salter C.J., Junor W., 1996, *MNRAS*, 282, 837
 Saikia D.J., Konar C., Kulkarni V.K., 2006, *MNRAS*, 366, 1391
 Saripalli L., Mack K.-H., 2007, *MNRAS*, 376, 1385
 Saripalli L., Hunstead R.W., Subrahmanyan R., Boyce E., 2005, *AJ*, 130, 896
 Schoenmakers A.P., 1999, *PhD Thesis*, Universiteit Utrecht, The Netherlands
 Schoenmakers A.P., de Bruyn A.G., Rottgering H.J.A., van der Laan H., Kaiser C.R., 2000a, *MNRAS*, 315, 371
 Schoenmakers A.P., de Bruyn A.G., Rottgering H.J.A., van der Laan H., 2000b, *MNRAS*, 315, 395
 Schneider D.P., et al., 2005, *AJ*, 130, 367
 Singal A.K., Konar C., Saikia D.J., 2004, *MNRAS*, 347, 79L
 Sirothia S.K., Saikia D.J., Ishwara-Chandra C.H., Kantharia N.G., 2009, *MNRAS*, 392, 1403
 Spergel D.N. et al., 2003, *ApJS*, 148, 175
 Subrahmanyan R., Saripalli L., Hunstead R.W., 1996, *MNRAS*, 279, 257
 Swarup G., Sinha R.P., Hildrup K., 1984, *MNRAS*, 208, 813
 Wang J.C.L., Sulkanen M.E., Lovelace R.V.E., 1992, *ApJ*, 390, 46