Spectro-Polarimetric Observations of Solar Magnetic Fields and the SOHO/MDI Calibration Issue

M.L. Demidov¹ · H. Balthasar²

Received: 11 May 2009 / Accepted: 18 August 2009 / Published online:

Abstract Comparisons of solar magnetic-field measurements made in different spectral lines are very important, especially in those lines in which observations have a long history or (and) specific diagnostic significance. The spectral lines Fe_I 523.3 nm and Fe_I 525.0 nm belong to this class. Therefore, this study is devoted to a comprehensive analysis using new high-precision Stokesmeter full-disk observations. The disk-averaged magnetic-field strength ratio R = B(523.3)/B(525.0) equals 1.97 ± 0.02 . The center-to-limb variation (CLV) is $R = 1.74 - 2.43\mu + 3.43\mu^2$, where μ is the cosine of the center-to-limb angle. For the disk center, we find R = 2.74, and for near-limb areas with $\mu = 0.3$, R equals 1.32. There is only a small dependence of R on the spatial resolution. Our results are rather close to those published three decades ago, but differ significantly from recent magnetographic observations. An application of our results to the important SOHO/MDI magnetic data calibration issue is discussed. We conclude that the revision of the SOHO/MDI data, based only on the comparison of magnetic field measurements in the line pair Fe_I 523.3 nm and Fe_I 525.0 nm (increasing by a factor of 1.7 or 1.6 on average according to recent publications) is not obvious and new investigations are urgently needed.

Keywords: Magnetic fields, Photosphere

1. Introduction

Many tasks of solar and solar-terrestrial physics urgently require precise, quantitative information about the distribution of magnetic fields across the solar disk. However, the extremely complicated spatial structure of magnetohydrodynamic parameters in the solar atmosphere leads to differences in magnetic field data from different spectral lines. This is the reason why comprehensive analyses of magnetic-field measurements made in different spectral lines are important for

Russia

email: demid@iszf.irk.ru

¹ Institute of Solar-Terrestrial Physics, Siberian Branch,

Russian Academy of Sciences, 664033 Irkutsk, P.O. Box 291.

² Astrophysikalisches Institut Potsdam (AIP), An der Sternwarte 16, 14482 Potsdam, Germany

diagnostics of solar magnetism and for the analysis of different data sets. In the first case, it is better to use observations performed at the same instrument, to avoid possible influences of different instrumental, method, time-dependent, and other effects. In the second case, the aim is to construct composite long-term data sets combining different observatories to study, for example, the temporal variations of solar magnetic parameters on different time scales.

A very significant part of solar magnetic-field observations has been performed (and are performed up to now) in the spectral line Fe I 525.0 nm. Therefore, comparisons of observations in other spectral lines with observations in this particular line are extremely important. The atomic parameters of Fe I 525.0 nm (large Landé factor, g = 3, small lower level excitation potential, $\chi_l = 0.12 \text{ eV}$) stimulated vigorous discussions about the reliability of measurements in this line, which are not finished yet. Different authors (some aspects of this issue are given in the discussion in Demidov *et al.*, 2008 and in the recent paper of Ulrich *et al.*, 2009) offer different correction factors to reduce raw measurements in Fe I 525.0 nm to the "true" ones.

Because of its low temperature sensitivity, one of the lines used as a "standard" for such correction is the rather strong line Fe_I 523.3 nm with $g_{\rm eff} = 1.3$ and $\chi_l = 2.93$ eV (Harvey and Livingston, 1969; Howard and Stenflo, 1972; Frazier and Stenflo, 1972; Frazier and Stenflo, 1978; Ulrich, 1992; Ulrich *et al.*, 2002; Tran *et al.*, 2005; Ulrich *et al.*, 2009). However, the combination of this line and Fe_I 525.0 nm is far from being perfect for diagnostics of solar magnetic fields, because they have different thermal dependencies, and Fe_I 523.3 nm is not intrinsically narrow (see Solanki, 1993). With much more success, it could be used as an example for the thermal line ratio. As shown by Socas-Navarro *et al.* (2008), reliable information about the magnetic-field parameters in the quiet regions (which cover most of the solar surface) can be achieved only using a combination of several lines with particular characteristics.

Nevertheless, observations in the discussed lines are still important in the context of the cross-calibration of different data sets, especially for the data recalibration problem of the Michelson Doppler Imager (MDI) onboard the Solar and Heliospheric Observatory (SOHO), see http://soi.stanford.edu/magnetic and Solar News (issues No. 19, 2007 and No. 25, 2008). The SOHO/MDI observations of magnetic fields are performed in the spectral line Ni I 676.8 nm with $g_{\rm eff}$ = 1.43. The main reasons for two subsequent serious revisions of the magnetic field data from MDI, which are widely used in the scientific community, are based on the comparison of observations in Fe_I 523.3 nm and Fe_I 525.0 nm made at the Mount Wilson Observatory (MWO) and published by Tran et al. (2005) and Ulrich et al. (2009). According to the first paper, based on the results of Ulrich multiplied by the factor $\delta^{-1} = B(523.3)/B(525.0) = 4.5 - 2.5 \sin^2 \theta$ or, according to the second paper, by the factor $\delta^{-1} = 4.15 - 2.82 \sin^2 \theta$, where θ is the centerto-limb angle. As a consequence, to fit SOHO/MDI data to MWO observations in Fe_I 525.0 nm corrected in this way, they must be multiplied by a factor that is a function of θ , see Figure 6 and Table 3 of Tran *et al.* (2005) and whose mean value over the disk is 1.7. According to Ulrich *et al.* (2009), the factor is 1/0.619= 1.615 for $\theta \leq 30^{\circ}$.

The fact, that the new MWO results strongly contradict the previous ones of Howard and Stenflo (1972), who found that the center-to-limb variation (CLV) of the coefficient is $\delta^{-1} = 0.48 + 1.33\mu$, with $\mu = \cos\theta$, requires independent measurements. For disk center, the factor is only 1.8 instead of 4.12 according to Ulrich *et al.* (2009). Their factor between B(523.3) and B(525.0) can be up to 5.5 (see their Figure 3), which can hardly be explained by "the fact that the shifted Zeeman components at $\lambda 525.0$ nm are shifted beyond the sampling passband of the MWO Babcock magnetograph" (Ulrich *et al.*, 2009).

The above mentioned circumstances make an additional exploration of solar magnetic field observations in Fe_I 523.3 nm and Fe_I 525.0 nm an urgent task. The entire spectral-line profile should be considered. This is the main aim of our study.

2. Observations and Analysis

All previous observations concerning the relationship between magnetic-field data in Fe I 523.3 nm and Fe I 525.0 nm were obtained with photoelectrical magnetographs. Drawbacks of such data are evident, because direct information about the distribution across the spectral line profile of the Stokes parameters (Stokes V/I_c in the case of longitudinal field) is missing – the only reliable indicator of magnetic field in the spatial range of line formation. Some assumptions and calibrations are necessary to obtain magnetic-field strengths from intensity variations in the exit slits of the magnetograph, which might be different for different spectral lines in the case of simultaneous observations.

Spectropolarimetric observations, as used in the present study, are much more reliable and informative. To study the problem of the comparison of solar magnetic-field measurements in combination of these lines, several series of observations in different regimes have been performed with the Stokes-meter of the Solar Telescope for Operative Predictions (STOP) at the Sayan Solar Observatory (SSO). Basic information about this instrument and methodical issues can be found in Demidov et al. (2002) and Demidov et al. (2008). STOP is equipped with a linear, 29 mm wide Toshiba TCD CCD detector with 3424 pixels (height 200 μ m, width 8 μ m), which allows us to obtain high-precision measurements of the Stokes parameters I and V/I_c in several spectral lines simultaneously. The solar surface is scanned following a 2D-raster, and the spatial resolution depends on defocussing the solar image. The standard angular resolution used in the regular observing programs (magnetograms of the large-scale magnetic fields, LSMF) is 100'' and the scanning step is 91''. To cover both lines with a single exposure, we used the fourth order of the spectrograph, where more than 2.7 nm are available. Part of the observations obtained with this dispersion was done with the standard spatial resolution (Figures 7 and 8), but most with focused solar image and standard scanning step, what corresponds to a resolution of $\approx 10''$.

The cycle time (one phase of the electro-optical modulator $-KD^*P$ crystal in our case) was 30 ms, the signal integration time in every point of the scanning process across the Sun was four seconds with and four seconds without the halfwave plate. The $\lambda/2$ plate is periodically inserted into the light beam in front of the coelostat to monitor the zero-level position.

An example of Stokes I and Stokes V/I_c profiles across the whole CCD detector is shown in Figure 1. The data correspond to the point next to the solar disk center, observed on 3 February, 2009 with a resolution of about 10". It is easy to see how much information is provided by such observations.

To derive quantitative information about magnetic and thermodynamic parameters from Stokes profiles such as those shown in Figure 1 is a complicated task. We use the approach (see Demidov et al., 2002), which numerically imitates the work of solar magnetographs in the center-of-gravity mode (see Solanki, 1993), to obtain the magnetic flux density (in the following we will use the term "magnetic field strength" for it throughout the paper) from the recorded data. The code calculates the position of the spectral line in two states $(\pm \lambda/4)$ of the polarization analyzer. Parameters for the "exit slits" are selected corresponding to the different lines. This position corresponds to the middle between the "slits", where the intensities are equal. If λ_1 is the wavelength of the spectral line in one analyser state and λ_2 in the other one, the value $\Delta \lambda = (\lambda_1 - \lambda_2)/2$ is considered as the Zeeman shift associated with the magnetic-field strength B by the known equation $\Delta \lambda = 4.67 \times 10^{-5} g \lambda^2 B$, where B is in Gauss and λ in cm. For the case shown in Figure 1, the magnetic strength calculated this way is $-26 \,\mathrm{G}$ for Fe_I 525.0 nm and -64 G for Fe_I 523.3 nm. Parameters for the "exit slits" are: the width W is equal to 9.32 pm, the separation Δ (distance between line center and the "slits" centers) is equal to 6.21 pm for the first line and W = 24.84 pm, $\Delta = 15.53$ pm for the second line.

The data used for Figure 1 offer the possibility to estimate the errors of our measurements. To do this the noise level of the Stokes V/I_c signal in the continuum can be used. The best choice for it is the part of the spectrum on the right of Fe I 524.4 nm. Calculations made for the ≈ 150 pm wide band show that the rms value in this region is 1.2×10^{-4} . Taking into account that $V/I_c = 0.01$ corresponds to ≈ 26 G for Fe I 525.0 nm and ≈ 64 G for Fe I 523.3 nm, we conclude that the formal error of the measurements is ≈ 0.3 G for the first line and ≈ 0.8 G for the second.

Figure 2 shows the Stokes I and Stokes V/I_c profiles for this pair of lines from the same observations as in Figure 1. Figure 3 compares the Stokes V/I_c profiles of both lines. Despite of the large difference in the g-factors, the amplitudes of the Stokes V/I_c -profiles are almost the same and the peak positions of the Stokes V/I_c profiles are close to the steepest parts of the corresponding Stokes I profiles. As a consequence of the fact that Fe I 523.3 nm is deeper and wider than Fe I 525.0 nm, the separation of the Stokes V/I_c extrema for the first line is almost two times larger than for the second one, although the splitting factor of the latter is larger. Therefore, in the case of weak fields, the separation of the Stokes V/I_c profiles peaks cannot be a measure of the magnetic field strength when lines with different widths are used. Indeed, if the magnetic-field strength is far from the Zeeman saturation regime, the locations of the Stokes V/I_c profile peaks are insensitive to the field and determined from the derivative of the Stokes I profile.



Figure 1. Spectra of Stokes parameters I (top panel) and V/I_c (bottom panel) along the whole wavelength band including Fe_I 523.3 nm and Fe_I 525.0 nm, covered by the detector. These observations were obtained close to the solar disk center with a magnetic-field strength of -26 G measured in Fe_I 525.0 nm.



Figure 2. Examples of Stokes I (in relative units and normalized to fit the scale of graph) and Stokes V/I_c profiles for Fe_I 523.3 nm (top panel) and for Fe_I 525.0 nm (bottom panel). The same observations as in Figure 1 were used.



Figure 3. Example of Stokes V/I_c profiles of Fe₁ 523.3 nm and Fe₁ 525.0 nm. The same observations as in Figure 1 and Figure 2 were used.

Already Figure 3 allows for the disk center a preliminary estimate of the ratio R = B(523.3)/B(525.0). In agreement with Demidov *et al.* (2008), we prefer the notation R for the ratio or the regression coefficient. The ratio of the Stokes V/I_c amplitudes (average of the amplitudes of the blue and red Stokes V/I_c peaks) of the two lines is V(523.3)/V(525.0) = 0.0097/0.0106 = 0.915. Normalizing to the corresponding Landé factors, we obtain $R = 0.915 \times (3.0/1.3) = 0.915 \times 2.3 = 2.1$. This value is rather close to the estimation R = B(523.3)/B(525.0) = (-64 G/-26 G) = 2.4, obtained from calculations of the magnetic-field strengths using wide parts of the line wings.

One point is not enough for a reliable estimation of the B(523.3)/B(525.0)ratio. The scatter plot, calculated for three full-disk magnetograms observed on 1, 2, and 3 February, 2009, is shown in Figure 4. Parameters for the "exit slits" are the same as mentioned above. We see a rather high correlation coefficient $\rho = 0.93$, and a systematic difference with regression coefficient $R = 1.97 \pm 0.02$. To calculate R, the reduced major axis method (Davis, 1986) is used. A formula to estimate the error (which is determined from the scattering of the points, and which is less when the correlation coefficient is higher) is taken from this book too. According to our measurements, the average factor that should be used to adjust measurements in Fe I 525.0 nm to those in Fe I 523.3 nm, is 1.97 ± 0.02 .

The next logical step is to check for possible spatial variations of the ratio R = B(523.3)/B(525.0) across the solar disk. For this purpose, we divide the solar disk into polar and equatorial sectors as in Demidov *et al.* (2008). We derive a polynomial fit $R = 1.74 - 2.43 \,\mu + 3.43 \,\mu^2$, shown in Figure 5. There are no significant differences in the CLV of R between polar and equatorial sectors of



Figure 4. Correlation and regression analysis of solar magnetic field observations in Fe I 523.3 nm and Fe I 525.0 nm. Data for three days of observations, 1, 2 and 3 February, 2009, are used. N is the number of points, ρ is the correlation coefficient, and R is the coefficient of linear regression (the slope of the line through the scatter plot).

the disk. At disk center, R is 2.74, and closer to the limb, at $\mu = 0.3$, R is 1.32. The dashed line is a quadratic polynomial fit through the observed points.

A significant decrease of the ratio B(523.3)/B(525.0) with increasing heliocentric distance is illustrated in Figure 6, where V/I_c profiles of the analyzed lines are shown for the point next to the south pole with $\mu = 0.56$ and with magnetic field strengths B(525.0) = 11.6 G and B(523.3) = 19.8 G. The ratio of these strengths, $R \approx 1.5$, is rather close to R = 1.45 from the analytical formula in Figure 5 at $\mu = 0.56$.

As mentioned above, observations in Fe I 523.3 nm and Fe I 525.0 nm with the STOP traditional spatial resolution of 100" were performed besides the observations with the rather high spatial resolution in order to look for a possible dependence of R = B(523.3)/B(525.0) on the spatial resolution. Ulrich (1992) found a significant increase of R with decreasing spatial resolution: R = 4.5 at disk center at a resolution of 20" instead of R = 3.9 at a resolution of 5".

Examples of full-disk magnetograms with a resolution of 100", obtained on 1 February, 2009 are shown in Figure 7. Despite of the very weak magnetic field strengths everywhere across the disk (minimum of solar activity, there were no sunspots and active regions), both maps show identical spatial structures, and they are well correlated. This is proved by the regression and correlation analysis presented in Figure 8. Calculations of the CLV of R = B(523.3)/B(525.0) for these data yield: $R = 2.24 + 0.03\mu + 0.56\mu^2$. We see that R is 2.83 at disk center, slightly more than the value of R = 2.74 obtained with high-spatial resolution. Despite the decrease of the resolution by a factor of ten, we encounter only a



Figure 5. Center-to-limb variation of the ratio R = B(523.3)/B(525.0) for the polar (N-S) and equatorial (E-W) sectors of the solar disk.



Figure 6. Example of Stokes V/I_c profiles of Fe I 523.3 nm and Fe I 525.0 nm for a point near the south pole with $\mu = 0.56$ and with a magnetic-field strength B = 11.6 G from Fe I 525.0 nm.



Figure 7. Full-disk magnetograms observed with a spatial resolution of 100'' in Fe₁ 523.3 nm (left panel) and Fe₁ 525.0 nm (right panel) on 1 February, 2009.



Figure 8. Correlation and regression analysis of solar magnetic field observations in Fe I 523.3 nm and Fe I 525.0 nm for observations with a spatial resolution of 100'', made on 1 February, 2009. Notations are the same as in Figure 4.

small $\approx 3\%$ increase of *R*. The difference is much larger for large heliocentric distances, but most probably it is caused by a weakening of the field strengths and by an increasing influence of the noise.

3. Discussion and Conclusion

The cornerstone of the study by Ulrich *et al.* (2009) is the difference of magnetic field strengths measured in the different parts of the profile of Fe I 523.3 nm, *i.e.*

the sampling of the line. They wrote: "contrary to the assumptions of Frazier and Stenflo (1972), Stenflo (1973) and Frazier and Stenflo (1978) we find that the field indicated by λ 5233 Å depends on the spectral sampling" and that "we [...] recommend the use of a sampling point as close to the line core as is practical". In the work of Ulrich (1992), the parameters of the spectral configurations with fiber-optic image reformattors were: centered on Fe I 525.0 nm – separation $\Delta = \pm 3.9$ pm; centered on Fe I 523.3 nm: – $\Delta = \pm 4.5$ pm.

In the recent work, Ulrich *et al.* (2009) obtained new MWO observations with the same sampling for Fe I 525.0 nm and with a new set of separations of $\pm 0.9, \pm 2.9, \pm 8.4, \pm 10.2$, and ± 17.7 pm for Fe I 523.3 nm. Computations of magnetic fields from various spectral sampling pairs and at ± 8.4 pm have shown different values of correlations coefficients ρ and slopes in the corresponding scatter plots, depending on the value of the separation and the range of θ (see Figures 1 to 3 and Table I in Ulrich *et al.*, 2009). The correlation coefficients for measurements at ± 8.4 pm with the other ones were: $\rho = 0.50$ at ± 0.9 pm, $\rho = 0.90$ at ± 2.9 pm, $\rho = 0.90$ at ± 10.2 pm and $\rho = 0.50$ at ± 17.7 pm. Only the correlation coefficients for the pairs nearest to ± 8.4 pm are reliable. In two other cases they are too small for a solid statistical analysis. The examples of slopes obtained by Ulrich *et al.* (2009) for the central area of the disk (range of sin θ from 0.0 to 0.4) are: 1.665 ± 0.067 at ± 0.9 pm, 1.795 ± 0.027 at ± 2.9 pm, 1.162 ± 0.010 at ± 10.2 pm and 2.694 ± 0.094 at ± 17.7 pm.

The fact that magnetic fields measured in points near the line center $(\pm 2.9 \text{ pm})$ are weaker (by a factor of 0.73 after application of additional assumptions and calculations, see Figure 12 of Ulrich *et al.*, 2009) than those measured in a point near the middle of the line wings $(\pm 8.4 \text{ pm})$ is the basic argument for the decrease of the correction coefficient R, following from direct comparison of data sets in Fe I 523.3 nm (sampling $\pm 8.4 \text{ pm}$) and Fe I 525.0 nm. For the disk center, the value R = 4.12 is obtained instead of 5.5.

We are convinced that only the Stokes V/I_c profile over the whole spectral line can characterize the magnetic field properties in the range of the line formation. But for the comparison with MWO results, it is important to find relationships between strengths in the different parts of Fe_I 523.3 nm in our data. For this purpose, calculations with different "exit slit" parameters, close to those recently used at MWO, were performed. The results for 3 February, 2009 with 10" spatial resolution are given in the following.

The values of "slit" widths and separations are expressed in the integer numbers of CCD pixels. In observations, the width of one pixel is equal to 0.777 pm. Following Ulrich *et al.* (2009), all data were analyzed relatively to the measurements in the part of the wings with a separation of $\Delta = 8.5$ pm and with a "slit" width of W = 9.32 pm. The comparison of data with these "slit" parameters with other ones are given in Table 1. For the points at ± 3.1 pm, our result differs from that of Ulrich *et al.* (2009). According to our data, magnetic fields measured closer to the line center are almost the same (only 1.07 times weaker) as measured at ± 8.5 pm. That means, if we correct measurements at ± 8.5 pm taking into account the regression coefficient $B(\pm 8.5 \text{ pm})/B(\pm 3.1 \text{ pm}) = 1.07$, we get a small difference to measurements in Fe I 525.0 nm. Indeed, Ulrich *et al.* (2009) found $B(523.3 \pm 8.4 \text{ pm})/B(525.0 \pm 3.9 \text{ pm}) = 5.5$ for the disk center.

| $\Delta\lambda \; [\mathrm{pm}]$ | Slit width W [pm] | ρ | $R = B(\pm 8.5 \text{pm})/\text{B}(\pm \Delta \lambda)$ |
|----------------------------------|-------------------|-------|---|
| | | | |
| ± 15.53 | 24.82 | 0.991 | 1.271 ± 0.009 |
| | | | |
| ± 0.8 | 0.80 | 0.870 | 1.018 ± 0.028 |
| ± 3.1 | 1.55 | 0.977 | 1.073 ± 0.013 |
| ± 10.1 | 9.32 | 0.998 | 1.027 ± 0.003 |
| ± 17.8 | 9.32 | 0.935 | 1.528 ± 0.030 |

M.L.Demidov, H.Balthasar

Applying a special correction factor, they obtained the value of 4.12. In our case we find $5.5 \times (1/1.07) = 5.1$.

All results described above have a direct impact on the problem of the calibration of SOHO/MDI magnetic-field data. Some correction coefficients suggested by MWO scientists were already mentioned before. The correction factors for the MDI data are much less if we use the values derived from our observations instead of those obtained by Tran *et al.* (2005) and by Ulrich *et al.* (2009). For disk center, we obtain 1.61 /(4.15/2.74) = 1.06 instead of 1.61 according to Ulrich *et al.* (2009).

Our independent estimations of the SOHO/MDI correction factors are obtained using our previous results from the comparison of SOHO/MDI data with SSO observations (important: both analyzed data sets correspond to spatial resolution 100''), published by Demidov *et al.* (2008). From Figure 1(e) of that paper we get an average value of 2.75 for the B(SOHO/MDI)/B(SSO) ratio. In order to correct the SOHO/MDI data (when they coincide with the corrected data in Fe_I 525.0 nm), we have to multiply them with the coefficient 1.97/2.75= 0.72, if we take R = B(523.3)/B(525.0) = 1.97 from our data with 10" spatial resolution, and 2.55/2.75 = 0.93, if we take R = 2.55 from data (see Figure 8) with 100'' spatial resolution. Our correction coefficient is smaller by a factor of about two compared to the recent MWO results (Tran et al., 2005, Ulrich et al., 2009), and instead of increasing the magnetic field strengths from SOHO/MDI data we have to decrease them. We would like to note here, that, according to the study of Demidov et al. (2008), observations at Mount Wilson and Sayan observatories made in Fe_I 525.0 nm are in almost perfect agreement with each other. Therefore, the problem lies in the differences between the observations in the strong line Fe I 523.3 nm.

It seems that this result contradicts to the result of Berger and Lites (2003), who obtained a correction factor of 1.56, based on a comparison of SOHO/MDI and *Advanced Stokes Polarimeter* (ASP) data. But observations with the ASP were made in a different spectral line (Fe I 630.25 nm), and using different lines can lead to quite different results (see Gopasyuk *et al.*, 1972 and Demidov *et al.*, 2008). Further, Berger and Lites (2003) analyzed only observations made in active regions, instead of full-disk quiet-Sun observations in our case. Besides

that, the observations with ASP and SOHO/MDI were made with much higher spatial resolution.

In connection with this issue it is important to note that comparisons of SOHO/MDI with Kitt Peak spectro-polarimeter data (Jones and Ceja, 2001, Wenzler, Solanki, and Krivova, 2005) caused a decrease of the SOHO/MDI measurements by a factor of ≈ 1.4 to adjust them to the Kitt Peak data.

Summarizing, we arrive at the conclusion that the issue of calibrating SOHO/ MDI data is rather far from being solved, and new investigations and observations are urgently needed.

Acknowledgements We appreciate the comments of the unknown reviewer which lead to significant improvements of our paper. The results presented in this study were obtained partly under support by the DFG (German Science Foundation) grant BA 1875/5-1. The MDI data from board the SoHO spacecraft used in this study are producted by the Stanford-Lockheed Institute for Space Research and the Solar Oscillations Investigation (SOI) in the W.W. Hansen Experimental Physics Laboratory of Stanford University and the Solar and Astrophysics Laboratory of the Lockheed-Martin Advanced Technology Center, USA. We took the magnetograms from the web-site http://soi.stanford.edu/data/ supported by the SOI and MDI team. This study includes data from the synoptic program at the 150-Foot Solar Tower of the Mt. Wilson Observatory. The Mt. Wilson 150-Foot Solar Tower is operated by UCLA, with funding from NASA, ONR and NSF, under agreement with the Mt. Wilson Institute.

References

Berger, T.E, Lites, B.W.: 2003, Solar Phys. 213, 213.

- Davis, J.C.: 1986, Statistics and Data Analysis in Geology, John Wiley, New York.
- Demidov, M.L., Zhigalov, V.V., Peshcherov, V.S., Grigoryev, V.M.: 2002, Solar Phys. 209, 217.
- Demidov, M.L., Golubeva, E.M., Balthasar, H., Staude, J., Grigoryev, V.M.: 2008, Solar Phys. 250, 279.
- Frazier, E.N., Stenflo, J.O.: 1972, Solar Phys. 27, 330.
- Frazier, E.N., Stenflo, J.O.: 1978, Astron. Astrophys. 70, 789.
- Gopasyuk, S.I., Kotov, V.A., Severny, A.B, Tsap, T.T.: 1973, Solar Phys. 31, 307.
- Harvey, J., Livingston, W.: 1969, Solar Phys. 10, 283.
- Howard, R., Harvey, J.W.: 1970, Solar Phys. 12, 23.
- Jones. H.P., Ceja, J.A.: 2001, In Sigwarth, M. (ed.), Advanced Solar Polarimetry Theory, Observation, and Instrumentation, CS-236, Astron. Soc. Pac., San Francisco, 87.
- Howard, R., Stenflo, J.O.: 1972, Solar Phys. 22, 402.
- Socas-Navarro, H., Borrero, J.M., Asensio Ramos, A., Collados, M., Domínquez Cerdeña, I.,Khomenko, E.V., Martínez González, M.J., Martínez Pillet, V., Ruiz Cobo, B., Sánchez Almeida, J.: 2008, Astrophys. J. 674, 596.
- Solanki, S.K.: 1993, Space Sci. Rev. 63, 1.
- Stenflo, J.O.: 1973, Solar Phys. 32, 41.
- Tran, T., Bertello, L., Ulrich, R.K., Evans, S.: 2005, Astrophys. J. Suppl. 156, 295.
- Ulrich, R.K.: 1992, In M.S. Giampapa, M.S., Bookbinder, J.M. (eds.), Proc. 7th Cambridge Workshop, Cool Stars, Stellar Systems, and the Sun, CS-26, Astron. Soc. Pac., San Francisco, 265.
- Ulrich, R.K., Evans, S., Boyden, J.E., Webster, L.: 2002, Astrophys. J. Suppl. 139, 259.
- Ulrich, R.K., Bertello, L., Boyden, J.E., Webster, L.: 2009, *Solar Phys.* 255, 53.
- Wang, H, Sheeley, N.R., Jr.: 1995, Astrophys. J. 447, L143.
- Wenzler, T., Solanki, S.K., Krivova, N.A.: 2005, Astron. Astrophys. 432, 1057. 213, 87.

sp1015.tex; 4/02/2022; 0:13; p.14